

SL(2, R)-MODULE STRUCTURE OF THE EIGENSPACES OF THE CASIMIR OPERATOR

ESTHER GALINA
JORGE VARGAS

ABSTRACT. In this paper, on the space of smooth sections of a $SL(2, R)$ -homogeneous vector bundle over the upper half plane we study the $SL(2, R)$ structure for the eigenspaces of the Casimir operator. That is, we determine its Jordan-Hölder sequence and the socle filtration. We compute a suitable generalized principal series having as a quotient a given eigenspace. We also give an integral equation which characterizes the elements of a given eigenspace. Finally, we study the eigenspaces of twisted Dirac operators.

§1. Introduction

Let $G = SL(2, \mathbf{R})$ and K be a fixed maximal compact subgroup K of G . Let (τ, V) be a representation of K , we denote

$$\begin{aligned} C^\infty(G/K, V) &= \{ f : G \rightarrow V \mid f \text{ is } C^\infty \text{ and } f(gk) = \tau(k)^{-1}f(g) \text{ for all } k \in K \} \\ L^2(G/K, V) &= \{ f : G \rightarrow V \mid f(gk) = \tau(k)^{-1}f(g) \text{ for all } k \in K, \|f\|_2^2 < \infty \} \end{aligned}$$

where $\|\cdot\|_2$ is computed with respect to Haar measure. On $L^2(G/K, V)$ we fix the obvious topology. On $C^\infty(G/K, V)$ we fix the topology of uniform convergence on compacts of the functions and their derivatives. Both spaces are representations of G under the left regular action $L_g f(x) = f(g^{-1}x)$ for all $g, x \in G$.

Let Ω the Casimir element of the universal algebra $\mathcal{U}(g_o)$ of the Lie algebra g_o of G , Ω define a G -left invariant operator on $C^\infty(G/K, V)$. Here, we obtain the G -module structure of each eigenspace of the Casimir operator

$$\Omega : C^\infty(G/K, V) \rightarrow C^\infty(G/K, V)$$

whenever V is an irreducible representation of K . Actually, we prove that whenever an eigenspace is irreducible, then it is infinitesimally equivalent to a principal series representation, and when an eigenspace is reducible then we have an exact sequence

1991 *Mathematics Subject Classification*. 1980 *Mathematics Subject Classification (1985 Revision)*. Primary 22E47.

Key words and phrases. Eigenspaces of the Casimir.

This work was supported by CONICET, CONICOR and Fa.M.A.F. (Argentina), ICTP (Trieste, Italy)

$0 \rightarrow W \rightarrow A_\lambda^n \rightarrow M \rightarrow 0$, where A_λ^n is the λ -eigenspace of Ω in $C^\infty(G/K, V)$, W is a Verma module and M an irreducible Verma module.

As a corollary we obtain the eigenvalues and eigenspaces of

$$\tilde{\Omega}: L^2(G/K, V) \rightarrow L^2(G/K, V)$$

From this, it results that if λ is an eigenvalue of $\tilde{\Omega}$ the corresponding eigenspace is a proper subset of the respective one of Ω . We also compute the L^2 -eigenspaces of the Dirac operator \mathbf{D} .

Knapp-Wallach [K-W] obtained an integral operator which sends an adjusted principal series onto the K -finite vector of the L^2 -kernel of the Dirac operator \mathbf{D} . In this work we obtain a similar result for each L^2 -eigenspace of \mathbf{D} (c.f §4).

Let $\phi_{\lambda, n}$ be the Eisenstein function (cf. ***) in $C^\infty(G/K, V)$ that belongs to the λ -eigenspace of Ω , we prove:

(i) a continuous function that satisfies the integral equation

$$\int_K f(gkx)dk = f(g)\phi_{\lambda, n} \quad \text{for all } g, x \in G$$

is smooth and is an eigenfunction of Ω corresponding to the eigenvalue λ .

(ii) Any λ -eigenfunction of Ω satisfies the integral equation in (i).

Now, we establish some notations,

$$(1.2) \quad \begin{aligned} K &= \left\{ k_\theta = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} : \theta \in \mathbf{R} \right\} \\ A &= \left\{ a_t = \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} : t \in \mathbf{R}^+ \right\} \\ M &= \left\{ \pm \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right\} \\ N &= \left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} : x \in \mathbf{R} \right\} \\ A^+ &= \{a_t \in A : 1 < t\} \\ A^- &= \{a_t \in A : 0 < t < 1\} \end{aligned}$$

We will use the decompositions $G = KAN$ and $G = KAK = K\overline{A^+}K = K\overline{A^-}K$ [K]. If we denote by

$$(1.3) \quad X = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad Y = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \quad H = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

the Iwasawa decomposition of the Lie algebra \mathfrak{g}_o of G is $\mathfrak{g}_o = \mathfrak{k}_o \oplus \mathfrak{a}_o \oplus \mathfrak{n}_o$ where $\mathfrak{k}_o = \mathbf{R}X$, $\mathfrak{a}_o = \mathbf{R}H$, $\mathfrak{n}_o = \mathbf{R}Y$. We denote by g, k, a, n their complexifications.

The Casimir operator Ω is an element of the universal algebra $\mathcal{U}(g)$ of g , moreover, the center of $\mathcal{U}(g)$ is $\mathbf{C}[\Omega]$ [L]. It is defined by

$$(1.4) \quad \Omega = \frac{1}{2} (H^2 - H - YX)$$

If

$$(1.5) \quad W = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad E_+ = \frac{1}{2} \begin{pmatrix} 1 & i \\ i & -1 \end{pmatrix} \quad E_- = \frac{1}{2} \begin{pmatrix} 1 & -i \\ -i & -1 \end{pmatrix}$$

another expression of Casimir operator is

$$(1.6) \quad \Omega = \frac{1}{8} (W^2 + 2W + 4E_-E_+)$$

W , E_+ and E_- satisfy the relations

$$\overline{W} = -W \quad \overline{E_{\pm}} = E_{\mp} \quad [E_+, E_-] = W \quad [W, E_{\pm}] = \pm 2E_{\pm}$$

Let θ be the usual Cartan involution on g_o . Therefore, k_o is the subspace of fix points of θ . Let p_o be the (-1) -eigenspace of θ .

The Killing form in g_o is

$$B(X, Y) = 4\text{Trace}(XY).$$

Thus $\{\frac{1}{2}E_+, \frac{1}{2}E_-\}$ is an orthonormal base of p with respect to the hermitian form

$$-B(X, \theta\overline{Y})$$

The Iwasawa decomposition for E_+ and E_- is

$$(1.7) \quad \begin{aligned} \frac{1}{2}E_+ &= \frac{1}{4}W + \frac{1}{4} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 0 & i \\ 0 & 0 \end{pmatrix} \\ \frac{1}{2}E_- &= -\frac{1}{4}W + \frac{1}{4} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 0 & i \\ 0 & 0 \end{pmatrix} \end{aligned}$$

§2. Eigenspaces of Ω

Since K is abelian, the irreducible representations of K are onedimensional. They are (τ_n, V_n) with $n \in \mathbf{Z}$, where

$$\dim V_n = 1 \text{ and } \tau_n(k_\theta)v = e^{in\theta}v \quad \text{for all } v \in V_n$$

Given $n \in \mathbf{Z}$, the elements of the center of the universal enveloping algebra of g will be considered acting on $C^\infty(G/K, V_n)$ as left invariant operators.

For all $\lambda \in \mathbf{C}$ define

$$(2.1) \quad A_\lambda^n = \left\{ f \in C^\infty(G/K, V_n) \quad \middle/ \quad \Omega f = \frac{\lambda^2 - 1}{8} f \right\}$$

Since Ω is a continuous linear operator on $C^\infty(G/K, V_n)$, it follows that A_λ^n is a closed subspace of $C^\infty(G/K, V_n)$. Thus, A_λ^n is a subrepresentation of $C^\infty(G/K, V_n)$ with infinitesimal character $\chi_{\lambda\delta}$, where δ is the linear functional of a_o such that $\delta(H) = \frac{1}{2}$ and $\chi_{\lambda\delta}$ is the character of \mathbf{C} multiplication by $\frac{\lambda^2 - 1}{8}$.

We denote by $A_\lambda^n[m]$ the K -type τ_m of A_λ^n .

PROPOSITION 2.1.

Given $n \in \mathbf{Z}$, $\lambda \in \mathbf{C}$, the representation A_λ^n of G is admissible and finitely generated. Moreover,

- (i) $\dim A_\lambda^n[m] \leq 1$ for all $m \in \mathbf{Z}$
- (ii) If $A_\lambda^n[m] \neq \{0\}$, then n and m have the same parity.

Remark: The converse of (ii) is also true. It follows from proposition 2.4.

We need some results to prove the proposition 2.1

Let $f \in A_\lambda^n[m]$, f is a spherical function of type (m, n) because

$$f(k_\theta g k_\psi) = e^{-im\theta} f(g) e^{-in\psi} \quad \text{for all } g \in G, k_\theta, k_\psi \in K$$

Since $G = KAK$, the values of f are determined by its values on A . Besides, if $m \neq n$ then $f|_K \equiv 0$. In fact, the equality $f(k_\theta) = f(k_\theta.1) = e^{-im\theta} f(1)$, implies that $f|_K \neq 0 \Leftrightarrow f(1) \neq 0$, now since f is spherical of type (m, n) we have that $f(k_\theta) = f(1.k_\theta) = f(1)e^{-in\theta} = f(1)e^{-im\theta}$, therefore if $f|_K$ were nonzero we would have that $m = n$.

The subgroup A is Lie isomorphic to \mathbf{R}^+ (positive real numbers with the usual product) by the isomorphism $\alpha(a_t) = t^2$.

Lemma 2.2.

If $f \in A_\lambda^n[m]$, the function $F : \mathbf{R}^+ \rightarrow \mathbf{C}$ associated to f given by $F(\alpha(a)) = f(a)$ for all $a \in A$ satisfy the differential equation

$$(2.2) \quad z^2 \frac{d^2}{dz^2} - \frac{2z^3}{1-z^2} \frac{d}{dz} - \frac{z^2}{(1-z^2)^2} (m^2 + n^2) + \frac{z(1+z^2)}{(1-z^2)^2} nm - \frac{\lambda^2 - 1}{4} = 0$$

The equation has regular singularities at the points $0, \pm 1, \infty$.

A proof of this lemma is in [Ca-M].

Proof of the Proposition 2.1. Since Ω is an elliptic operator in $C^\infty(G/K, V_n)$, if $f \in A_\lambda^n$, $f|_A$ is real analytic. Therefore, the function $F : \mathbf{R}^+ \rightarrow \mathbf{C}$ defined in (2.2) is a real analytic function. Hence there is a holomorphic extension of F to a neighborhood of \mathbf{R}^+ in the right half plane.

On the other hand by the Frobenius theory for differential equations with regular singular points [C-page 132] the equation (2.2) has an analytic solution on a neighborhood of 1 if and only if m and n have the same parity. Moreover, any holomorphic solution of (2.2) is a multiple of the power series

$$(2.3) \quad (z-1)^{\frac{1}{2}|m-n|} \sum_{j=0}^{\infty} c_j (z-1)^j \quad c_0 = 1$$

In fact, the indicial equation of (2.2) is

$$s(s-1) + s - \frac{1}{4}(m-n)^2 = 0$$

and its roots are $\pm\frac{1}{2}(m-n)$. Thus, as the roots differ by an integer, the exponent of the first term of (2.3) is $\frac{1}{2}|m-n|$, if this number were not an integer the function (2.3) would not be analytic on a neighborhood of 1, this forces the same parity for n and m .

As the other singularities of (2.2) are $0, -1, \infty$, there is an extension of the analytic solution on a neighborhood of 1 to an analytic solution on a neighborhood of \mathbf{R}^+ . So (i) and (ii) holds. \square

Remark. Since A_λ^n has infinitesimal character $\chi_{\lambda\delta}$ and A_λ^n is admissible by Proposition 2.1, A_λ^n has finite length by a known result of Harish-Chandra [V, Corollary 5.4.16].

Corollary 2.3.

Given $n \in \mathbf{Z}$, $\lambda \in \mathbf{C}$, the K -type τ_n occurs in any subrepresentation of A_λ^n . Moreover, A_λ^n has a unique irreducible G -submodule.

Proof. Let W be a nontrivial closed submodule of A_λ^n and denote by W_K the set of K -finite elements in W , we consider the map

$$(*) \quad \begin{array}{ccc} \text{Hom}_G(W, A_\lambda^n) & \longrightarrow & \text{Hom}_K(W_K, V_n) \\ T & \longrightarrow & (v \rightarrow \tilde{T}v = Tv(1)) \end{array}$$

This map is well defined. In fact, if $v \in W_K$,

$$\tilde{T}(kv) = T(kv)(1) = (L_k.Tv)(1) = Tv(k^{-1}) = \tau_n(k)Tv(1)$$

Moreover, it is injective. In fact, suppose that $\tilde{T} \equiv 0$, so $Tv(1) = 0$ for all $v \in W_K$. As T is a continuous linear transformation, W_K is a dense subset of W [L-page 24], and there exists a sequence $\{v_m\}$ in W_K such that $v_m \rightarrow w$ for each $w \in W$, then

$$Tv_m \rightarrow Tw \implies 0 = Tv_m(1) \rightarrow Tw(1)$$

that is, $Tw(1) = 0$ for all w . Now, for $w \in W$,

$$Tw(g) = (L_{g^{-1}}.Tw)(1) = T(g^{-1}w)(1) = 0 \quad \text{for all } g \in G,$$

so $T \equiv 0$. If W is a closed submodule of A_λ^n , by (*) $W[n] \neq 0$, and by (i) $W[n] = A_\lambda^n[n]$. This concludes the first statement of the corollary. The second follows from the equality $W[n] = A_\lambda^n[n]$. \square

Fix $n \in \mathbf{Z}$, $\lambda \in \mathbf{C}$, let δ be the linear functional on \mathfrak{a}_o such that $\delta(H) = \frac{1}{2}$, $\log a_t = tH$, and denote by $(-1)^n$ the character of M such that $\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \rightarrow (-1)^n$. As usual, define

$$(2.4) \quad \begin{aligned} \text{I}_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1) &= \\ &= \{f : G \rightarrow \mathbf{C} \mid C^\infty \text{ such that} \\ & f(xman) = e^{-(\lambda+1)\delta(\log a)}(-1)^n(m^{-1})f(x) \text{ for all } x \in G, man \in MAN\} \end{aligned}$$

the representation of G induced by the representation $(-1)^n \otimes e^{\lambda\delta} \otimes 1$ of MAN . G acts by left translation. Recall that $I_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$ has infinitesimal character $\chi_{\lambda\delta}$ and $I_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$ is irreducible if and only if $\lambda \not\equiv (n+1) \pmod{2}$ [B].

Define linear transformations

$$(2.5) \quad \begin{array}{ccc} I_{MAN}^G((-1)^n \otimes e^{\pm\lambda\delta} \otimes 1) & \xrightarrow{T} & A_\lambda^n \\ f & \longrightarrow & (x \rightarrow Tf(x) = \int_K f(xk)\tau_n(k)dk) \end{array}$$

Whenever it becomes necessary to see which is the domain of the operators, we will write T_\pm , otherwise we will write T .

The linear transformation T is well defined because

$$Tf(xk') = \int_K f(xk'k)\tau_n(k)dk = \tau(k')^{-1} \int_K f(xk)\tau_n(k)dk.$$

Besides, since $I_{MAN}^G((-1)^n \otimes e^{\pm\lambda\delta} \otimes 1)$ has infinitesimal character $\chi_{\lambda\delta}$, T is a left G -morphism and left infinitesimal translation by Ω agrees with right infinitesimal translation, ($L_\Omega.f = R_\Omega.f$ for all $f \in C^\infty(G/K, V_n)$). Hence the image of T is contained in A_λ^n .

T is not zero because

$$T\tau_{-n}(1) = \int_K \tau_{-n}(k)\tau_n(k)dk = \int_K dk \neq 0$$

Note that A_λ^n and $A_{\lambda'}^n$ is the same eigenspace of Ω if $\lambda^2 = (\lambda')^2$. So, if $\lambda \in \mathbf{Z}$ we will always assume that $\lambda \geq 0$.

PROPOSITION 2.4.

Given $n \in \mathbf{Z}$,

- (i) If $\lambda \in \mathbf{C} \setminus \mathbf{Z}$, or $\lambda \in \mathbf{Z}$ and $\lambda \not\equiv (n+1) \pmod{2}$, A_λ^n is infinitesimally equivalent to $I_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$.
- (ii) If $\lambda \in \mathbf{Z}_{\geq 0}$, $\lambda+1 \equiv n \pmod{2}$ and $\lambda > |n|$, A_λ^n is infinitesimally equivalent to $I_{MAN}^G((-1)^n \otimes e^{-\lambda\delta} \otimes 1)$.
- (iii) If $\lambda \in \mathbf{Z}_{\geq 0}$, $\lambda+1 \equiv n \pmod{2}$ and $\lambda < n$, the (g, K) -module structure of A_λ^n is the following

$$\begin{aligned} E_+ A_\lambda^n[m] &\neq 0 && \text{for all } m \text{ such that } A_\lambda^n[m] \neq 0 \\ E_- A_\lambda^n[m] &\neq 0 && \text{for all } m \neq \pm\lambda \text{ such that } A_\lambda^n[m] \neq 0 \\ E_- A_\lambda^n[\pm\lambda+1] &= 0 \end{aligned}$$

- (iv) If $\lambda \in \mathbf{Z}_{\geq 0}$, $\lambda+1 \equiv n \pmod{2}$, $n < 0$ and $\lambda < -n$, the (g, K) -module structure of A_λ^n is the following

$$\begin{aligned} E_- A_\lambda^n[m] &\neq 0 && \text{for all } m \text{ such that } A_\lambda^n[m] \neq 0 \\ E_+ A_\lambda^n[m] &\neq 0 && \text{for all } m \neq \pm\lambda+1 \text{ such that } A_\lambda^n[m] \neq 0 \\ E_+ A_\lambda^n[\pm\lambda+1] &= 0. \end{aligned}$$

Remark 1: Under the hypothesis (iii) or (iv) we have that A_λ^n is not a quotient of $I_{MAN}^G((-1)^n \otimes e^{\pm\lambda\delta} \otimes 1)$.

Remark 2: A_λ^n is irreducible if and only if $\lambda \not\equiv (n+1) \pmod{2}$.

We need the following lemma to prove (iii) of proposition 2.4.

Lemma 2.5.

Given $n \in \mathbf{Z}$, let $\lambda \in \mathbf{Z}_{\geq 0}$, $\lambda + 1 \equiv n \pmod{2}$ and $\lambda < n$, there exist $m \in \mathbf{Z}$, $m < -\lambda$ such that $A_\lambda^n[m]$ is not zero.

Proof of Lemma 2.5. Let m be an integer such that

$$(2.6) \quad m \equiv n \pmod{2} \quad m < -\lambda \quad \frac{1}{2}(n - m) \text{ is even}$$

The conditions on m and n ensure the existence of a smooth solution F of (2.2) on the interval $(0, \infty)$. In fact, using the Frobenius method for differential equations with regular singularities, that (2.2) has a analytic solution in a neighborhood of 1 if and only if m and n have the same parity. Besides, the singularities of (2.2) are $0, \pm 1, \infty$. Therefore, this solution extends to a solution on the interval $(0, \infty)$. Moreover, any smooth solution of (2.2) in the interval $(0, \infty)$ is a multiple of the power series

$$(z - 1)^{\frac{1}{2}|m-n|} \sum_{j=0}^{\infty} c_j (z - 1)^j \quad c_0 = 1$$

Therefore, F has a zero of order $\frac{1}{2}|m - n|$ at 1.

We have to prove that F extends to an element of $A_\lambda^n[m]$. This will take some work.

Let $N_K(A)$ be the normalizer of A on K .

Consider $C_{\tau_{n-m}}^\infty(A)$ to be the set of smooth funtions on A such that

$$(j) \quad \phi(kak^{-1}) = \tau_{n-m}(k) \phi(a) \quad \text{for all } a \in A, k \in N_K(A)$$

$$(jj) \quad \frac{\phi(a)}{\delta(\log a)^{\frac{1}{2}(n-m)}} \text{ is a smooth function and even on } A.$$

Let $f: A \rightarrow \mathbf{C}$ given by $f(a) = F(\alpha(a))$, with α the isomorphism between A and \mathbf{R}^+ defined in (2.2). Let's prove that the function f is in $C_{\tau_{n-m}}^\infty(A)$. In fact, the normalizer of A on K , is exactly

$$N_K(A) = \{\pm I\} = \left\{ k_{\frac{\pi}{2}}, k_{-\frac{\pi}{2}} \right\}$$

As $n - m$ and $\frac{1}{2}(n - m)$ are even numbers,

$$\tau_{n-m}(\pm I) = \tau_{n-m}(k_{\pm \frac{\pi}{2}}) = e^{\pm i(n-m)\frac{\pi}{2}} = 1$$

So, f satisfy (j) if and only if $f(a) = f(a^{-1})$ for all $a \in A$, or equivalently $F(x) = F(x^{-1})$ for all $x \in \mathbf{R}^+$. Let's prove that $F(x) = F(x^{-1})$. Let h be the function given by $h(z) = F(z^{-1})$, we want to prove that $h = F$. We claim that h satisfies the same differential equation that F does. In fact, let $w = z^{-1}$, then

$$\begin{aligned} \frac{dh}{dz}(z) &= \frac{dF}{dw}(w) w' \\ &= -w^2 \frac{dF}{dw}(w) \end{aligned}$$

$$\begin{aligned} \frac{d^2 F}{dz^2}(z) &= -2ww' \frac{dF}{dw}(w) + w^4 \frac{d^2 F}{dw^2}(w) \\ &= 2w^3 \frac{dF}{dw}(w) + w^4 \frac{d^2 F}{dw^2}(w) \end{aligned}$$

and

$$\begin{aligned} -\frac{2z^3}{1-z^2} &= -\frac{2w^{-3}}{1-w^{-2}} = \frac{2w^{-1}}{1-w^2} \\ -\frac{z^2}{(1-z^2)^2} &= -\frac{w^{-2}}{(1-w^{-2})^2} = -\frac{w^2}{(1-w^2)^2} \\ \frac{z(1+z)}{(1-z^2)^2} &= \frac{w^{-1}(1+w^{-2})}{(1-w^{-2})^2} = \frac{w(w^2+1)}{(1-w^2)^2} \end{aligned}$$

So,

$$\begin{aligned} z^2 \frac{d^2 h}{dz^2}(z) - \frac{2z^3}{1-z^2} \frac{dh}{dz}(z) + \\ \left(-\frac{z^2}{(1-z^2)^2} (m^2 + n^2) + \frac{z(1+z^2)}{(1-z^2)^2} nm - \frac{\lambda^2 - 1}{4} \right) h(z) = \\ = w^2 \frac{d^2 F}{dw^2}(w) + \left(2w - \frac{2w^{-1}}{1-w^2} w^2 \right) \frac{dF}{dw}(w) + \\ + \left(-\frac{w^2}{(1-w^2)^2} (m^2 + n^2) + \frac{w(1+w^2)}{(1-w^2)^2} nm - \frac{\lambda^2 - 1}{4} \right) F(w) \end{aligned}$$

The right hand side is exactly the equation(2.2) on F , so it is zero. Both h and F are smooth functions on $(0, \infty)$ and solutions of the differential equation (2.2). So, by (2.6) they are multiple of each other in a neighborhood of 1. Hence, we write,

$$\begin{aligned} h(z) &= (z-1)^{\frac{1}{2}|n-m|} \psi_h(z) \\ F(z) &= (z-1)^{\frac{1}{2}|n-m|} \psi_F(z) \end{aligned}$$

with ψ_h and ψ_F power series, such that $c\psi_h(z) = \psi_F(z)$ for a suitable nonzero complex number. Therefore,

$$h(z) = F(z^{-1}) = (z^{-1} - 1)^{\frac{1}{2}|n-m|} \psi_F(z^{-1}) = (z-1)^{\frac{1}{2}(n-m)} z^{-\frac{1}{2}|n-m|} \psi_F(z^{-1})$$

Thus, $\psi_h(z) = (z-1)^{-\frac{1}{2}(n-m)} \psi_F(z^{-1})$. This imply that

$$c\psi_h(z) = (z-1)^{-\frac{1}{2}(n-m)} \psi_F(z^{-1})$$

Hence, $F(z) = F(z^{-1})$ in a neighborhood of 1. As F is real analytic in $(0, \infty)$, $F(z) = F(z^{-1})$ for all $z \in \mathbf{R}^+$. Equivalently, $f(a) = f(a^{-1})$ for all $a \in A$. Thus, f satisfies (j).

We want to prove that f satisfies (jj). The function $\delta(\log a)^{-\frac{1}{2}(n-m)}$ is even on A because

$$\begin{aligned} \delta(\log a_t)^{-\frac{1}{2}(n-m)} &= (t \delta(H))^{-\frac{1}{2}(n-m)} \\ &= (-t \delta(H))^{-\frac{1}{2}(n-m)} \quad \text{by (2.6)} \\ &= \delta(\log a_t^{-1})^{-\frac{1}{2}(n-m)} \end{aligned}$$

Thus, the function $f(a)\delta(\log a)^{-\frac{1}{2}(n-m)}$ is even. The function $f(a)\delta(\log a)^{-\frac{1}{2}(n-m)}$ is smooth because f is real analytic and has a zero of order $\frac{1}{2}(n-m)$ at 1. Therefore, we have proved that $f \in C_{\tau_{n-m}}^{\infty}(A)$. We want to extend f to an element of $A_{\lambda}^n[m]$

Let $C^{\infty}(G/K)[\tau_{n-m}]$ be the space of smooth complex valued functions on G/K such that $f(kx) = \tau_{n-m}(k)f(x)$ for all $k \in K, x \in G$.

We need to prove:

Sublemma 2.6.

The restriction map from $C^{\infty}(G/K)[\tau_{n-m}]$ to $C_{\tau_{n-m}}^{\infty}(A)$ is bijective.

Proof of sublemma 2.6. : The equality $G = KAK$ implies that the restriction map is injective. To prove that is surjective we appeal to a theorem of Helgason. Let \mathcal{H} be the set of harmonic polynomial functions on p_o . We consider the usual action of K on \mathcal{H} . That is, the one determined by the isotropy representation of K in p_o . We now set ourselves in §10 of [H-1], with $\delta = \tau_{n-m}$. Since $n \equiv m \pmod{2}$, we have that $\tau_{n-m} \in \hat{K}_o$. Let $\deg Q^{\delta}(\lambda) = p(\delta)$. A formula due to Kostant and cited on pag 203 of [H-1] says that $p(\delta) = d(\delta) = \text{degree of the harmonic homogeneous polynomials in the } \delta\text{-isotypic component of } \mathcal{H}$. To compute $d(\delta)$ we proceed as follow: If e_1, e_2 is an orthonormal basis for p_o , we know that $k(\theta)\dot{e}_1 = \cos(2\theta)e_1 - \sin(2\theta)e_2, k(\theta)\dot{e}_2 = \sin(2\theta)e_1 + \cos(2\theta)e_2$. Since $(n-m)/2$ is a whole number the polynomial function on $p_o, (e_1 + ie_2)^{(n-m)/2}$ is harmonic and has degree $(n-m)/2$, moreover $k(\theta)(e_1 + ie_2)^{(n-m)/2} = e^{i(n-m)\theta}(e_1 + ie_2)^{(n-m)/2}$. Thus, we have that $p(\delta) = (n-m)/2$. Therefore, our space $C_{\tau_{n-m}}^{\infty}(A)$ contains the space $\mathcal{D}^{\tau_{n-m}}(A)$ of page 211 in [H-1]. Hence, lemma 10.1 of [H-1] implies that the restriction map from $\mathcal{D}^{\tau_{n-m}}(G/K)$ into $\mathcal{D}^{\tau_{n-m}}(A)$ is a linear homeomorphism. We remark that $\mathcal{D}^{\tau_{n-m}}(G/K) \subset C^{\infty}(G/K)[\tau_{n-m}]$. A density argument together with the fact that K is compact imply sublemma 2.6. \square

We proceed with the proof of lemma 2.5. For this end, we now have that the function f admits a smooth extension $\tilde{f}: \exp p_o \rightarrow \mathbf{C}$ which satisfies

$$(2.7) \quad \begin{aligned} \tilde{f}(kak^{-1}) &= \tau_{n-m}(k)\tilde{f}(a) \\ &= \tau_m(k)^{-1}\tilde{f}(a)\tau_n(k) \end{aligned}$$

The diffeomorphism between G and $\exp p_o K$ ensures that the function $\hat{f}: G \rightarrow \mathbf{C}$ given by

$$\hat{f}(pk) = \tilde{f}(p)\tau_n(k)^{-1} \quad \text{for all } p \in \exp p_o, k \in K$$

is well defined and it is smooth. Also, \hat{f} is in the K -type τ_m of $C^{\infty}(G/K, V_n)$. In fact, for $x \in G$ we write $x = k_2ak_2^{-1}k_1$ with $k_1, k_2 \in K$, and $a \in A$, hence

$$\begin{aligned} (L_k\hat{f})(x) &= \hat{f}(k^{-1}k_2ak_2^{-1}k_1) = \tilde{f}(k^{-1}k_2ak_2^{-1}k)\tau_n(k^{-1}k_1)^{-1} \\ &= \tau_{n-m}(k^{-1}k_2)f(a)\tau_n(k^{-1}k_1)^{-1} \\ &= \tau_{n-m}(k^{-1})\tau_{n-m}(k_2)f(a)\tau_n(k^{-1}k_1)^{-1} \\ &= \tau_{n-m}(k^{-1})\tilde{f}(k_2ak_2^{-1})\tau_n(k^{-1})^{-1}\tau_n(k_1)^{-1} \\ &= \tau_n(k^{-1})\tau_m(k)\tilde{f}(p)\tau_n(k^{-1})^{-1}\tau_n(k_1)^{-1} \\ &= \tau_m(k)\tilde{f}(p)\tau_n(k_1)^{-1} \\ &= \tau_m(k)\hat{f}(x) \quad \square \end{aligned}$$

A computation like the one in [Wa] page 280, implies that

$$(\Omega \hat{f})(x) = \tau_m(k_2^{-1})\tau_n(k_2^{-1}k_1)(z^2 \frac{d^2 F}{d^2 z} + \dots) = 0$$

because F satisfies the equation 2.2.

This concludes the proof of lemma 2.5

Proof of the Proposition 2.4. (i) As T is not the zero function and since $\lambda \not\equiv n+1 \pmod{2}$ the module $\mathbf{I}_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$ is irreducible. Thus T is injective. The K -types τ_m which occur in $\mathbf{I}_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$ are indexed by all the m with the same parity as n . Since T is one-to-one they must occur in A_λ^n . By proposition 2.1 (i), (ii), they are exactly the K -types of A_λ^n . Thus, T is surjective at the level of (g, K) -modules.

(ii) Since $\lambda \geq 0$, $\mathbf{I}_{MAN}^G((-1)^n \otimes e^{-\lambda\delta} \otimes 1)$ has only one irreducible submodule F which is finite dimensional and whose K -types are parametrized by $\{m : -(\lambda-1) \leq m \leq \lambda-1, m \equiv n(2)\}$. The structure of $\mathbf{I}_{MAN}^G((-1)^n \otimes e^{-\lambda\delta} \otimes 1)$ is

$$\begin{array}{ccccccc} \mathbf{I}_{MAN}^G((-1)^n \otimes e^{-\lambda\delta} \otimes 1) & & \supset W_+ & & & & \\ & & \supset W_- & \supset F & \supset & 0 & \end{array}$$

where W_+ is the G -submodule spanned by the K -types $\{-(\lambda-1), -(\lambda-3), \dots, \lambda-1, \lambda+1, \dots\}$ and W_- is the one spanned by the K -types $\{\dots, \lambda-3, \lambda-1\}$. As $\lambda > |n|$ the K -type τ_n occur in F . On the other hand, we have verified that T maps non trivially the K -type τ_n , so F is not a submodule of $\text{Ker}T$. Since F is contained in every nonzero submodule of $\mathbf{I}_{MAN}^G((-1)^n \otimes e^{-\lambda\delta} \otimes 1)$. T is 1:1; by a similar argument to the one used on (i) we get that T is surjective.

(iii) Suppose that $n, \lambda > 0$ $\lambda < n, \lambda \not\equiv n+1(2)$. Then the image of T_- is the discrete serie $H_{\lambda\delta}$ of infinitesimal character $\chi_{\lambda\delta}$. We recall that the K -types of $H_{\lambda\delta}$ are parametrized by $\{\lambda+1, \lambda+3, \dots\}$. In fact, the nonzero quotients of $\mathbf{I}_{MAN}^G((-1)^n \otimes e^{-\lambda\delta} \otimes 1)$ are $H_{\lambda\delta}, H_{-\lambda\delta}, H_{\lambda\delta} \oplus H_{-\lambda\delta}$ or itself. Now, the irreducible finite-dimensional submodule occurs in $\text{Ker}T_-$, otherwise $T_-(F)$ would be an irreducible submodule of A_λ^n and do not have the K -type τ_n ($\lambda < |n|!$), that contradicts corollary 2.3. This contradiction ensures that T_- is not injective. By corollary 2.3, A_λ^n has only one irreducible submodule, $\text{Im}T_- \neq H_{\lambda\delta} \oplus H_{-\lambda\delta}$. Furthermore, since the irreducible submodule contains the K -type τ_n , so $\text{Im}T_- = H_{\lambda\delta}$. Therefore $H_{\lambda\delta}$ is the irreducible submodule of A_λ^n .

The structure of $\mathbf{I}_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$ is the following

$$\begin{array}{ccccccc} \mathbf{I}_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1) & \supset H_{\lambda\delta} \oplus H_{-\lambda\delta} & & \supset H_{\lambda\delta} & & & \\ & & & \supset H_{-\lambda\delta} & \supset & 0 & \end{array}$$

T_+ is not injective; otherwise $T_+(H_{-\lambda\delta})$ is an irreducible submodule of A_λ^n and does not have the K -type τ_n . Also $\text{Ker} T_+ \neq H_{\lambda\delta} \oplus H_{-\lambda\delta}$; otherwise, the finite dimensional representation F is a subrepresentation of A_λ^n , contradicting corollary 2.3. Thus,

$$\text{Im}T_+ \cong \mathbf{I}_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1) / H_{-\lambda\delta}$$

This implies that

$$(\mathrm{Im}T_+)_K = \bigcup_{\substack{m \geq -(\lambda-1) \\ m \equiv n(2)}} A_\lambda^n[m]$$

which is the Verma module of lowest weight $-(\lambda - 1)$. Thus,

$$\begin{aligned} E_+ A_\lambda^n[m] &\neq 0 && \text{for all } m \geq -(\lambda - 1) \\ E_- A_\lambda^n[m] &\neq 0 && \text{for all } m \geq -(\lambda - 1) \text{ and } m \neq -\lambda + 1 \end{aligned}$$

By lemma 2.5 there exists a K -type $A_\lambda^n[m] \neq 0$ for some $m < -\lambda$. This ensure that $A_\lambda^n[m] \neq 0$ for all $m < -\lambda$ and $m \equiv n \pmod{2}$, on the other hand, A_λ^n would have a lowest weight submodule with lowest weight less than $-\lambda\delta$. The infinitesimal character of this lowest weight submodule would be different from $\chi_{\lambda\delta}$, giving a contradiction. Following the same argument, E_+ acts nontrivially on each $A_\lambda^n[m]$, $m < -\lambda$.

For the case $\lambda = 0$ and $\lambda + 1 \equiv n \pmod{2}$ the proof is easier.

(iv) It has the same proof of (iii). This concludes the proof of proposition 2.4. \square

Remark 1: Given $n \in \mathbf{Z}$ and $\lambda \in \mathbf{C}$, the K -types $A_\lambda^n[m]$ are not zero for all m with the same parity of n .

Remark 2: In view of [S], in cases (i) and (ii) A_λ^n is equivalent to the maximal model of I_{MAN}^G which is the induced representation with hiperfunctions coefficients. In case (iii) A_λ^n is a quotient of the maximal model of a generalized principal series.

Remark 3: Given $n \in \mathbf{Z}_{\geq 0}$ and $\lambda \geq 0$ as in (iii) of proposition 2.4, the G -module structure of A_λ^n is

$$\cdots \quad \bullet \quad -(\lambda+1) \bullet \quad \xrightarrow[\leftarrow 0]{\neq 0} \bullet \quad -(\lambda-1) \quad \cdots \quad \lambda-1 \bullet \quad \xrightarrow[\leftarrow 0]{\neq 0} \bullet \quad \lambda+1 \quad \bullet \quad \cdots$$

the right arrows represent the action of E_+ and the left ones the action of E_- . That is, we have proved

Corollary 2.6.

Let $\lambda \in \mathbf{Z}_{\geq 0}$ and $\lambda \equiv n + 1 \pmod{2}$. A composition series for A_λ^n is

$$0 \rightarrow V \rightarrow A_\lambda^n \rightarrow M \rightarrow 0$$

where V is the Verma module of lowest weight $-(\lambda - 1)$ and M is the irreducible Verma module of highest weight $-(\lambda + 1)$.

PROPOSITION 2.7.

Given $n \in \mathbf{Z}$ and λ as in (iii) of proposition 2.4 (i.e. $\lambda \equiv n + 1 \pmod{2}$ and $\lambda \geq 0$ an integer), then A_λ^n is quotient of a generalized principal series $I_{MAN}^G(W_0)$ where $W_0 = \mathbf{R}^2$ and the representation of MAN is

$$\pm \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} e^t & 0 \\ 0 & e^t \end{pmatrix} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \rightarrow (-1)^n \exp t \begin{pmatrix} \lambda & 1 \\ 0 & -\lambda \end{pmatrix}$$

Proof. For $f = (f_1, f_2) \in \mathbf{I}_{MAN}^G(W_0)$ let

$$S : \mathbf{I}_{MAN}^G(W_0) \rightarrow C^\infty(G/K, V_n)$$

defined by

$$(Sf)(x) = \int_K f_1(xk)\tau_n(k) dk + \int_K f_2(xk)\tau_n(k) dk$$

Since $\mathbf{I}_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$ is contained in $\mathbf{I}_{MAN}^G(W_0)$ via the map $f \rightarrow F = (f, 0)$ and S restricted to $\mathbf{I}_{MAN}^G(W_0)$ is equal to T_+ , hence $\text{Im}(S)$ contains $\text{Im}(T_+)$. An easy calculation shows that $\text{Im}(S)$ contains properly $\text{Im}(T_+)$. Now, corollary 2.6 implies that any K -finite vector in A_λ^n outside of $\text{Im}(T_+)$ is cyclic in $A_\lambda^n/\text{Im}(T_+)$. Therefore, S is onto. \square

Now, consider the Casimir operator acting on the subspace of compactly supported functions in $C^\infty(G/K, V_n)$. We denote by $\tilde{\Omega}$ the unique essentially self-adjoint extension of Ω to a dense subspace of

$$L^2(G, V_n) = \left\{ f : G \rightarrow \mathbf{C} \quad \left/ \quad \begin{array}{l} f(xk) = \tau_n(k)^{-1} f(x) \\ \int_G |f(x)|^2 dx < \infty \end{array} \right. \right\}$$

(cf [A-S]).

PROPOSITION 2.8.

If $W_\lambda^n = \{f \in L^2(G/K, V_n) / \tilde{\Omega}f = \frac{\lambda^2-1}{8}f\}$, then W_λ^n is non zero if and only if $\lambda \in \mathbf{Z} - \{0\}$, $\lambda + 1 \equiv n \pmod{2}$ and $|\lambda| < |n|$. Moreover, $W_\lambda^n = W_{-\lambda}^n$ is isomorphic to the discrete series of Harish-Chandra parameter $\lambda\delta$.

Proof. Suppose that $\lambda \in \mathbf{Z} - \{0\}$, $\lambda + 1 \equiv n \pmod{2}$ and $|\lambda| < |n|$. As $\tilde{\Omega}$ is elliptic, a Connes-Moscovici result [C-M] ensure that W_λ^n is a sum of discrete series, actually, it is irreducible by the Frobenius Reciprocity. The K -finite elements of $L^2(G/K, V_n)$ are in the set of K -finite elements of $C^\infty(G/K, V_n)$, so $W_\lambda^n[m] \subset A_\lambda^n[m]$ for all $m \in \mathbf{Z}$. By proposition 2.4, A_λ^n has subspaces infinitesimally equivalent to a discrete series for λ such that

$$\lambda \in \mathbf{Z} \quad \lambda \equiv n + 1 \pmod{2}, \quad 0 < |\lambda| < |n|$$

This "discrete series" subspaces are really contained in $L^2(G/K, V_n)$. In fact, if $f \in A_\lambda^n[m]$ and it belongs to a "discrete series", then f satisfies the differential equation (2.2) or the one which results from the identification of A^+ with $\mathbf{R}_{>0}$ via $a_t \leftrightarrow t$. Then the theory of leading exponents as in [K] says that $f(a_t) e^{-(\lambda-1)t}$ at $t = \infty$. Now, the integral formula for the Cartan decomposition together with $\lambda > 0$ imply that f is square integrable. For negative λ we have a similar proof.

For the converse we use the structure of the discrete series, Frobenius Reciprocity together with proposition 2.4. This concludes proposition 2.8. \square

§3. L^2 and C^∞ -eigenspaces of the Dirac operator

Let $g_o = k_o \oplus p_o$ be the Cartan decomposition of g_o , then p_o is the subspace of symmetric matrix of g_o .

If we fix a minimal left ideal S in the Clifford algebra of p_o , the resulting representation of $so(p_o)$ brakes down in two irreducible representations. Such representation composed with the adjoint representation of k_o restricted to p_o lift up at a representation of K called the spin representation of K . Let $\{X_1, X_2\}$ be an orthonormal base of p_o , let c be the Clifford multiplication and fix an integer n . The Dirac operator

$$\mathbf{D}: C^\infty(G/K, V_{n+1} \otimes S) \rightarrow C^\infty(G/K, V_{n+1} \otimes S)$$

is defined by

$$(3.1) \quad \mathbf{D} = \sum_{i=1}^2 (1 \otimes c(X_i)) X_i$$

where X_i act as left invariant operators for all i . The spin representation S decompose into a sum of two irreducible subrepresentations $S = S^+ \oplus S^-$ (c.f. 4.2 bellow). If $X \in p_o$, then $c(X)S^\pm = S^\mp$, so

$$(3.2) \quad \mathbf{D}^\pm : C^\infty(G/K, V_n \otimes S^\pm) \rightarrow C^\infty(G/K, V_n \otimes S^\mp)$$

are well defined.

We also consider

$$\tilde{\mathbf{D}}: L^2(G/K, V_{n+1} \otimes S) \rightarrow L^2(G/K, V_{n+1} \otimes S)$$

Some properties of the Dirac operators \mathbf{D} and $\tilde{\mathbf{D}}$ are: both are elliptic G -invariant differential operator. As the Riemannian metric of G/K is complete, $\tilde{\mathbf{D}}$ and $\tilde{\mathbf{D}}^2$ are essentially selfadjoint in $L^2(G/K, V_{n+1} \otimes S)$ [W], that is, the minimal extension is the unique selfadjoint closed extension over the set of smooth compactly supported functions. Thus, we consider $\tilde{\mathbf{D}}$ equal to this extension which coincides with the maximal one [A]. The eigenvalues of $\tilde{\mathbf{D}}$ are defined as the eigenvalues of the unique selfadjoint extension.

The following proposition is a corollary to proposition 2.8.

PROPOSITION 3.1.

If α is an eigenvalue of $\tilde{\mathbf{D}}$, then the α -eigenspace $W_\alpha(\tilde{\mathbf{D}})$ is irreducible and it is a proper subspace of the α -eigenspace $W_\alpha(\mathbf{D})$ of \mathbf{D} . The eigenvalues of $\tilde{\mathbf{D}}$ are $\alpha \in \mathbf{R}$ such that $\alpha^2 = \frac{1}{8}(n+2)^2 - \lambda^2$ with λ integer and $0 < |\lambda| \leq n+1$.

Proof. For $G = SL(2, R)$ The Parthasarathy equality [A-S] is

$$(3.3) \quad \begin{aligned} \mathbf{D}^2 &= -\Omega + \frac{(n+1)^2 - 1}{8} Id \\ \tilde{\mathbf{D}}^2 &= -\tilde{\Omega} + \frac{(n+1)^2 - 1}{8} Id \end{aligned}$$

If α is a non-zero eigenvalue of $\tilde{\mathbf{D}}$,

$$(3.4) \quad W_{\alpha^2}(\tilde{\mathbf{D}}^2) = W_{\alpha}(\tilde{\mathbf{D}}) \oplus W_{-\alpha}(\tilde{\mathbf{D}})$$

(cf [G-V]). Because of (3.3), the left hand side of (3.4) is the $-\alpha^2 + (n+1)^2 - 1 = \frac{1}{8}(\lambda^2 - 1)$ eigenspace of the Casimir operator. Now, since $S = V_{-1} \oplus V_1$,

$$\mathbb{L}^2(G/K, V_{n+1} \otimes S) = \mathbb{L}^2(G/K, V_n) \oplus \mathbb{L}^2(G/K, V_{n+2})$$

Hence proposition 2.8 implies that $0 \leq \lambda \leq n+1$ and

$$\alpha^2 = \frac{(n+1)^2 - \lambda^2}{8}$$

Moreover,

$$W_{\alpha^2}(\tilde{\mathbf{D}}^2) = A_{\lambda}^n \cap L^2(G/K, V_n) \oplus A_{\lambda}^{n+1} \cap L^2(G/K, V_{n+2})$$

Thus, $W_{\alpha^2}(\tilde{\mathbf{D}}^2)$ is equal to the sum of two copies of the discrete series $H_{\lambda\delta}$. Since, $W_{\alpha}(\tilde{\mathbf{D}})$ is isomorphic to $H_{\lambda\delta}$ we get that $W_{\alpha}(\tilde{\mathbf{D}})$ is properly contained in $W_{\alpha}(\mathbf{D})$. \square

Corollary 3.2.

(τ_n, V_n) and (τ_{n+2}, V_{n+2}) are K -types of $W_{\alpha}(\tilde{\mathbf{D}})$ for every non-zero eigenvalue α of $\tilde{\mathbf{D}}$. For the case $\alpha = 0$, (τ_{n+2}, V_{n+2}) is contained in $\text{Ker } \tilde{\mathbf{D}}$ and (τ_n, V_n) is not.

§4. Szegő kernels associated to the eigenspaces of $\tilde{\mathbf{D}}$

In [K-W] Knapp and Wallach gave an integral operator to explicitly obtain a discrete serie as the image of a nonunitary principal serie when the discrete serie is realized as the kernel of Schmid operator. In §3 we have obtained that each eigenspace of the Dirac operator

$$\tilde{\mathbf{D}}: L^2(G/K, V_{n+1} \otimes S) \rightarrow L^2(G/K, V_{n+1} \otimes S)$$

is a discrete serie. The purpose of this section is to give an integral operator for each non zero eigenvalue α of $\tilde{\mathbf{D}}$ which will realize the eigenspace $W_{\alpha}(\tilde{\mathbf{D}})$ as a quotient of an appropriated principal serie. From §3 it is easy to deduce which will be the principal serie corresponding to each eigenspace $W_{\alpha}(\tilde{\mathbf{D}})$, the problem is to obtain the G -invariant integral operator onto $W_{\alpha}(\tilde{\mathbf{D}})$. Let $G = SL(2, \mathbf{R})$ and K the maximal compact subgroup defined as in (1.2).

Let V_{n+1} be the $n+1$ irreducible representation of K , we assume that $n+1 > 0$. In §3, given an orthonormal base of p_o it was defined the Dirac operator $\tilde{\mathbf{D}}$. If we take $\{X_i\}_{i=1}^2$ an orthonormal base of the complexification p of p_o , another expresion of $\tilde{\mathbf{D}}$ is

$$(4.1) \quad \tilde{\mathbf{D}} = \sum_{i=1}^2 (1 \otimes c(X_i)) \bar{X}_i$$

where bar is conjugation with respect to g_o .

One form to obtain the representations S^\pm is choosing the left minimal ideals of the Clifford algebra of p ,

$$S^+ = \mathbf{C}E_+ \quad S^- = \mathbf{C}E_-E_+$$

where the product is Clifford multiplication. In $Cliff(p)$ the following set of relations holds:

$$(4.2) \quad E_+^2 = E_-^2 = 0 \quad E_+E_-E_+ = -E_+$$

Hence $S = V_{-1} \oplus V_1$. Thus, we have that

$$V_{n+1} \otimes S = V_n \oplus V_{n+2}$$

The set of K -finite elements of a principal serie $I_{MAN}^G(\epsilon \otimes e^{\lambda\delta} \otimes 1)$ defined in (2.4), is the representation of K induced by ϵ of M , hence

$$I_M^K(\epsilon) = \bigoplus_{i \in \hat{K}} V_i \otimes \text{Hom}_M(V_i, \epsilon)$$

So, if the representation ϵ occur at V_n and V_{n+2} as M -submodule, then $\epsilon = (-1)^n$. We denote by i_j the inclusions

$$i_j: (\epsilon, W_\epsilon) \rightarrow (\tau_j, V_j) \quad j = n, n+2$$

As W_ϵ and V_j are one dimensional

$$W_\epsilon = \mathbf{C}w \quad V_j = \mathbf{C}v \otimes u$$

where $w \in W_\epsilon$, $v \in V_{n+1}$ and $u \in S^\pm$.

Then the inclusions i_j are determined by the constants a_j such that

$$(4.3) \quad i_j(w) = a_j v \otimes u \quad \text{where } u = \begin{cases} E_+ & j = n \\ E_-E_+ & j = n+2 \end{cases}$$

If $sg \alpha$ is the sign of the real number α , fix

$$a_n = \left(\frac{\lambda + n + 1}{-\lambda + n + 1} \right)^{\frac{1}{2}} sg \alpha \quad \text{con } 0 \neq \lambda \in \mathbf{Z}, |\lambda| \leq n$$

$$a_{n+2} = 1$$

Let $G = KAN$ be the Iwasawa decomposition of G . According to this decomposition we write an element of G by

$$x = \kappa(x)e^{H(x)}n(x)$$

Let $S(x, t)$ be the function on $G \times K$ defined by

$$(4.4) \quad S(x, t) = e^{(\lambda-1)\delta H(x^{-1}t)} (\tau_n(\kappa(x^{-1}t))i_n + \tau_{n+2}(\kappa(x^{-1}t))i_{n+2})$$

Let $\tau = \tau_n + \tau_{n+2}$ on $V_n \oplus V_{n+2}$, so (4.4) implies

$$(4.5) \quad S(xk, t) = \tau(k)^{-1} S(x, t) \quad \text{for all } k \in K$$

We will call $S(x, t)$ the Szegö kernel associated to the parameters $(\lambda, n+1)$. If $f \in \mathbf{I}_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$, the Szegö map associated to the parameters $(\lambda, n+1)$ is

$$(4.6) \quad \begin{aligned} S(f)(x) &= \int_K S(x, t) f(t) dt \\ &= \int_K e^{(\lambda-1)\delta H(x^{-1}t)} \tau(\kappa(x^{-1}t))(i_n + i_{n+2}) f(t) dt \end{aligned}$$

The equation (4.5) ensure that the image of the Szegö map is in $C^\infty(G/K, V_n \oplus V_{n+2})$.

Let $\tilde{\mathbf{D}}$ defined as in §3

PROPOSITION 4.1.

Given $n \in \mathbf{Z}$, α a non zero eigenvalue of $\tilde{\mathbf{D}}$, and λ a negative integer which satisfies the equality

$$\alpha = \frac{1}{8} (-\lambda^2 + (n+1)^2)^{\frac{1}{2}} \text{sg } \alpha$$

Then, the Szegö map of parameters $(\lambda, n+1)$ is a G -invariant operator onto the eigenspace $W_\alpha(\tilde{\mathbf{D}})$.

Before proving this result we will see that Szegö map is not the zero map. Let $f \in C^\infty(K/M, W_\epsilon)$ where $\epsilon = (-1)^n$, given by

$$f(k) = i^{-1} \tau_n(k)^{-1} i_n w$$

Extend f to G so that $f \in \mathbf{I}_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$.

$$\begin{aligned} (S(f)(1), i_n w) &= \int_K (\tau(t)(i_n + i_{n+2}) (i_n^{-1} \tau_n(t)^{-1} i_n w), i_n w) dt \\ &= \int_K (i_n w + \tau_{n+2}(t) i_{n+2} (i^{-1} \tau_n(t)^{-1} i_n w), i_n w) dt \\ &= \int_K \|i_n w\|^2 dt \\ &\neq 0 \end{aligned}$$

because $\tau_{n+2}(t) i_{n+2} (i^{-1} \tau_n(t)^{-1} i_n w) \in V_{n+2}$ which is orthogonal to V_n .

To see that the Szegö map is G -invariant we need next lemma

Lemma 4.2.

Let S be the Szegö map with parameters $(\lambda, n+1)$. If $f \in \mathbf{I}_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$ then

$$S(f)(x) = \int_K \tau(t)(i_n + i_{n+2}) f(xt) dt$$

Proof of Lemma 4.2. Using the change of variable

$$\int_K h(k) dk = \int_K h(\kappa(x^{-1}t))e^{-2\delta H(x^{-1}t)} dt$$

for $h(k) = \tau(k)(i_n + i_{n+2}) f(xk)$ the following equality holds

$$\begin{aligned} \int_K \tau(k)(i_n + i_{n+2}) f(xk) dk &= \\ &= \int_K \tau(\kappa(x^{-1}t))e^{-2\delta H(x^{-1}t)}(i_n + i_{n+2}) f(x\kappa(x^{-1}t)) dt \end{aligned}$$

As A normalize N ,

$$\begin{aligned} x^{-1}t &= \kappa(x^{-1}t)e^{H(x^{-1}t)}n(x^{-1}t) \\ x\kappa(x^{-1}t) &= tn(x^{-1}t)^{-1}e^{-H(x^{-1}t)} \\ &= te^{-H(x^{-1}t)}n' \quad \text{with } n' \in N \end{aligned}$$

So, $f(x\kappa(x^{-1}t)) = f(te^{-H(x^{-1}t)}n') = e^{(\lambda+1)\delta H(x^{-1}t)} f(t)$. And

$$\begin{aligned} \int_K \tau(k)(i_n + i_{n+2}) f(xk) dk &= \int_K \tau(\kappa(x^{-1}t))e^{(\lambda-1)\delta H(x^{-1}t)}(i_n + i_{n+2}) f(t) dt \\ &= \int_K S(x, t) f(t) dt \quad \square \end{aligned}$$

Proof of the Proposition 4.1. By the lemma 4.2 the Szegö map is G -equivariant for left regular actions. As $\tilde{\mathbf{D}}$ also commute with the action of G , it is enough to see that if $f \in \mathbf{I}_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$

$$\tilde{\mathbf{D}}(Sf)(1) = \alpha Sf(1)$$

If $f \in \mathbf{I}_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$, the image of f is in $W_\epsilon = \mathbf{C}w$ with $\epsilon = (-1)^n$, then $f(t) = h(t)w$ with h a complex valued function. So,

$$\begin{aligned} Sf(x) &= \int_K S(x, t)wh(t) dt \\ \tilde{\mathbf{D}}Sf(1) &= \int_K \tilde{\mathbf{D}}(S(x, t)w)_{x=1}h(t) dt \end{aligned}$$

from which we only need prove that

$$\begin{aligned} D(S(x, t)w)_{x=1} &= \alpha S(1, t)w \\ &= \alpha \tau(t)(i_n w + i_{n+2}w) \end{aligned}$$

Let X_1, X_2 be an orthonormal base of p . Then,

$$\begin{aligned}
& \tilde{\mathbf{D}}(S(x, t)w)_{x=1} = \\
& = (I \otimes c) \left(\sum_{i=1}^2 (X_i S(x, t)w)_{x=1} \otimes \bar{X}_i \right) \\
& = (I \otimes c) \left(\sum_{i=1}^2 \frac{d}{du} \Big|_{u=0} e^{(\lambda-1)\delta H(\exp(-uX_i)t)} \tau(\kappa(\exp(-uX_i)t)) (i_n + i_{n+2})w \otimes \bar{X}_i \right) \\
& = (I \otimes c) \left(\sum_{i=1}^2 \frac{d}{du} \Big|_{u=0} e^{(\lambda-1)\delta H(\exp(-u\text{Ad}(t^{-1})X_i)t)} \tau(\kappa(\exp(-u\text{Ad}(t^{-1})X_i)t)) (i_n + i_{n+2})w \otimes \bar{X}_i \right) \\
& = (I \otimes c) \left(\tau(t) \otimes \text{Ad}(t) \sum_{i=1}^2 (\text{Ad}(t^{-1})X_i) S(1, 1)w \otimes \overline{\text{Ad}(t^{-1})X_i} \right)
\end{aligned}$$

As $\{\text{Ad}(t^{-1})X_i\}_{i=1,2}$ is another orthonormal base of p , and

$$\tau(t)(I \otimes c) = (I \otimes c)(\tau(t) \otimes \text{Ad}(t))$$

then

$$\tilde{\mathbf{D}}(S(x, t)w)_{x=1} = \tau(t)\tilde{\mathbf{D}}(S(x, 1)w)_{x=1}$$

So we must prove

$$\begin{aligned}
\tilde{\mathbf{D}}(S(x, 1)w)_{x=1} & = \alpha S(1, 1)w \\
& = \alpha(i_n + i_{n+2})w
\end{aligned}$$

Let $\frac{1}{2}E_-, \frac{1}{2}E_+$ be the orthonormal base of p given in §1, then

$$\begin{aligned}
& \tilde{\mathbf{D}}(S(x, t)w)_{x=1} = \\
& = (I \otimes c) \left(\frac{d}{du} \Big|_{u=0} e^{(\lambda-1)\delta H(\exp(-u\frac{1}{2}E_-))} \tau(\kappa(\exp(-u\frac{1}{2}E_-))) (i_n + i_{n+2})w \otimes \frac{1}{2}E_+ \right. \\
& \quad \left. + \frac{d}{du} \Big|_{u=0} e^{(\lambda-1)\delta H(\exp(-u\frac{1}{2}E_+))} \tau(\kappa(\exp(-u\frac{1}{2}E_+))) (i_n + i_{n+2})w \otimes \frac{1}{2}E_+ \right)
\end{aligned}$$

By (1.7)

$$\begin{aligned}
\tilde{\mathbf{D}}(S(x, t)w)_{x=1} & = (I \otimes c) \left(-(\lambda-1)\delta\frac{1}{4} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} (i_n + i_{n+2})w \otimes \frac{1}{2}E_+ - \right. \\
& \quad - (\lambda-1)\delta\frac{1}{4} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} (i_n + i_{n+2})w \otimes \frac{1}{2}E_+ - \\
& \quad - \tau \left(\frac{1}{4} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \right) (i_n + i_{n+2})w \otimes \frac{1}{2}E_+ - \\
& \quad \left. - \tau \left(-\frac{1}{4} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \right) (i_n + i_{n+2})w \otimes \frac{1}{2}E_- \right)
\end{aligned}$$

By (4.2) and (4.3) applying $I \otimes c$, the following holds

$$c(\frac{1}{2}E_+)i_n w = c(\frac{1}{2}E_-)i_{n+2}w = 0$$

and by (4.4)

$$\begin{aligned} c(\tfrac{1}{2}E_-)i_n w &= \tfrac{1}{2}a_n i_{n+2} w \\ c(\tfrac{1}{2}E_+)i_{n+2} w &= -\tfrac{1}{2}\tfrac{1}{a_n} i_w \end{aligned}$$

So that

$$\begin{aligned} \tilde{\mathbf{D}}(S(x, t)w)_{x=1} &= \\ &= -\tfrac{1}{8}(-\lambda + 1)\tfrac{1}{a_n} i_n w + \tfrac{1}{8}(-\lambda + 1)a_n i_{n+2} w + \tfrac{1}{8}(n + 2)\tfrac{1}{a_n} i_n w + \tfrac{1}{8}n a_n i_{n+2} w \\ &= \tfrac{1}{8}(\lambda + n + 1)\tfrac{1}{a_n} i_n w + \tfrac{1}{8}(-\lambda + n + 1)a_n i_{n+2} w \end{aligned}$$

because

$$\begin{aligned} \delta \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} &= 1 \\ \tau_j \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} v &= jv \quad \text{si } v \in V_{j\delta} \quad j = n, n + 2 \end{aligned}$$

The coefficients of $i_n w$ and $i_{n+2} w$ are

$$\begin{aligned} \tfrac{1}{8}(\lambda + n + 1)\tfrac{1}{a_n} &= \tfrac{1}{8}(\lambda + n + 1) \left(\frac{-\lambda + n + 1}{\lambda + n + 1} \right)^{\frac{1}{2}} sg \alpha \\ &= \tfrac{1}{8}(-\lambda^2 + (n + 1)^2)^{\frac{1}{2}} sg \alpha \\ &= \alpha \end{aligned}$$

$$\begin{aligned} \tfrac{1}{8}(-\lambda + n + 1)a_n &= \tfrac{1}{8}(-\lambda^2 + (n + 1)^2)^{\frac{1}{2}} sg \alpha \\ &= \alpha \end{aligned}$$

That is,

$$\tilde{\mathbf{D}}(S(x, 1)w)_{x=1} = \alpha S(1, 1)w$$

Now, we will prove that the Sezgö map of parameters $(\lambda, n + 1)$ for negative λ maps onto $W_\alpha(\tilde{\mathbf{D}})$. We know by proposition 3.1 that $W_\alpha(\tilde{\mathbf{D}})$ is irreducible. As S is non zero, if $\text{Im}(S)$ is square integrable, then $\text{Im}(S) = W_\alpha(\tilde{\mathbf{D}})$. $\text{Im}(S)$ is a subset of the eigenspace $W_\alpha(\tilde{\mathbf{D}})$ of the Dirac operator $\tilde{\mathbf{D}}$. But $W_\alpha(\tilde{\mathbf{D}})$ is a subset of $W_{\alpha^2}(\tilde{\mathbf{D}}^2)$. According with the notation of §2, as $\tilde{\mathbf{D}}^2$ differ with the Casimir operator Ω by a constant, $W_{\alpha^2}(\tilde{\mathbf{D}}^2)$ is isomorphic to $A_\lambda^n \oplus A_\lambda^{n+2}$. But the only quotient of $I_{MAN}^G((-1)^n \otimes e^{\lambda\delta} \otimes 1)$ isomorphic to a subspace of $A_\lambda^n \oplus A_\lambda^{n+2}$ is infinitesimally equivalent to a discrete serie. Let $\phi \in \text{Im}(S)$ in a non zero K -type, as the action of this K -type is one and the set of K -finite elements of the square integrable function space is a subset of the K -finite elements of the C^∞ , then ϕ is square integrable. So $\text{Im}(S)$ is a subset of $W_\alpha(\tilde{\mathbf{D}})$. The irreducibility concludes the proof. \square

REFERENCES.

[A] M.Atiyah - "Elliptic Operators, Discrete Groups and von Neumann Algebras" Astérisque - Vol.32-33, 1976.

[A-S] M.Atiyah and W.Schmid - "A Geometric Construction of the Discrete Series for Semisimple Lie groups" Inventiones Mathematicae - Vol.42, 1977.

[B] W.Barker - " L^p Harmonic Analysis in $SL(2, \mathbf{L})$ " Memoirs of the American Mathematical Society.

[C] Coddington y Levinson - "Theory of Ordinary Differential Equations" Mc Graw Hill, New York, 1955.

[C-M] A.Connes and H.Moscovici - "The L^2 -index Theorem for Homogeneous Spaces of Lie groups" Annals of Mathematics -Vol. 115, N.2, 1982.

[Ca-M] W.Casselmann and D.Milicic - "Asymptotic Behavior of Matrix Coefficients of Admissible Representations" Duke Math.- J.49, 1982,869-930.

[G-V] E.Galina and J.Vargas - Eigenvalues and eigenvectors for the twisted Dirac operator over $SU(n, 1)$ and $Spin(2n, 1)$ accepted by Transactions of the American Math. Soc.

[H-1] S.Helgason - "A Duality for Simetric Spaces with Applications to Group Representations, II. Differential Equations and Eigenspace Representations" Advances in Mathematics - Vol.22, N.2, 1976, 187-219.

[H-2] S.Helgason - "Group and Geometric Analysis" Academic Press,1984. Inc.

[K] A.Knapp - "Representation Theory of Real Reductive Groups" Princeton University Press. 1986.

[K-W] A.W.Knapp and N.R.Wallach - "Sezgo Kernels Associated with Discrete Series" Inventiones Mathematicae - Vol.34, F.3, 1976, 163-200.

[S] W.Schmid - "Boudary Value Problems for Group Invariant Differential Equations" Astérisque, hors serie, 1985.

[V] D.Vogan - "Representations of Real Reductive Lie Groups" Birkhäuser, Boston, 1981.

[Wa] Wallach, Harmonic analysis on homogeneous spaces, Marcell Deker, 1974.

[W] J.A.Wolf - "Essential Self Adjointness for the Dirac Operator and its Square" Indiana University Mathematical Journal-Vol 22, N.7 , Jan. 1973.