

# ON THE CLASSIFICATION OF FINITE-DIMENSIONAL POINTED HOPF ALGEBRAS

NICOLÁS ANDRUSKIEWITSCH AND HANS-JÜRGEN SCHNEIDER

ABSTRACT. We classify finite-dimensional complex Hopf algebras  $A$  which are pointed, that is, all of whose irreducible comodules are one-dimensional, and whose group of group-like elements  $G(A)$  is abelian such that all prime divisors of the order of  $G(A)$  are  $> 7$ . Since these Hopf algebras turn out to be deformations of a natural class of generalized small quantum groups, our result can be read as an axiomatic description of generalized small quantum groups.

## INTRODUCTION

One of the very few general classification results for Hopf algebras is due to Milnor, Moore, Cartier and Kostant around 1963. It says that any cocommutative Hopf algebra over the complex numbers is a semidirect product of the universal enveloping algebra of a Lie algebra and a group algebra.

In the terminology of Sweedler's book [S] from 1969, cocommutative Hopf algebras (over an algebraically closed field) are examples of *pointed Hopf algebras*, that is, all their simple subcoalgebras are one-dimensional, or equivalently, all their simple comodules are one-dimensional. Thus duals of finite-dimensional pointed Hopf algebras are analogs of basic algebras in the theory of finite-dimensional algebras. A rich supply of examples of non-cocommutative pointed Hopf algebra was only found in the mid-eighties of the last century: The Drinfeld-Jimbo quantum groups  $U_q(\mathfrak{g})$ ,  $\mathfrak{g}$  a semisimple Lie algebra, and their multiparameter versions, as well as the finite-dimensional small quantum groups introduced by Lusztig a bit later are all pointed.

In the present paper we assume that the ground-field  $k$  is algebraically closed of characteristic zero. If  $A$  is a Hopf algebra, we denote the group of group-like elements of  $A$  by

$$G(A) = \{g \in A \mid \Delta(g) = g \otimes g, \varepsilon(g) = 1\}.$$

---


Results of this paper were obtained during a visit of H.-J. S. at the University of Córdoba, partially supported through a grant of CONICET. The work of N. A. was partially supported by CONICET, Fundación Antorchas, Agencia Córdoba Ciencia, ANPCyT and Secyt (UNC).

We classify all finite-dimensional pointed Hopf algebras  $A$  over  $k$  with abelian group  $G(A)$  such that the prime divisors of the order of  $G(A)$  are  $> 7$ . We describe these Hopf algebras by generators and relations, and we show that they are the small quantum groups discovered by Lusztig and variations of them. Thus our results can be viewed as an axiomatic description of generalized small quantum groups.

Other types of Hopf algebras occur if prime divisors  $\leq 7$  of the order of the abelian group  $G(A)$  are allowed. However, extending the methods of this paper it should be possible to describe their structure by some generalization of Cartan matrices. Finite-dimensional pointed Hopf algebras with non-abelian group  $G(A)$  seem to be of a very different nature. Their structure is not understood.

We now give a brief overview of the classification program for finite-dimensional Hopf algebras. We remark that these Hopf algebras give rise to finite tensor categories in the sense of [EO] and thus classification results on finite-dimensional Hopf algebras should have applications in conformal field theory [Ga].

The classification splits into several very different parts according to the behaviour of the coradical. Recall that the coradical  $A_0$  of a Hopf algebra  $A$  is the sum of all its simple subcoalgebras.



If  $A = A_0$  then  $A$  is semisimple as an algebra. Semisimple Hopf algebras define examples of fusion categories. There are various important results on semisimple Hopf algebras, but there is at present no general strategy to classify these algebras.

Next assume that  $A_0 \neq A$  and that the coradical  $A_0$  is a Hopf subalgebra. If  $A$  is pointed then  $A_0$  is a Hopf subalgebra, namely the group algebra  $k[G(A)]$ . The only general method for the classification of a class of Hopf algebras is the Lifting Method, developed in [AS1] and

which works for Hopf algebras whose coradical is a Hopf subalgebra. Since the coradical is a semisimple Hopf algebra and the classification of semisimple Hopf algebras is still widely open, it is natural to concentrate on pointed Hopf algebras. The starting point of this method is an analog of the Milnor-Moore-Cartier-Kostant decomposition theorem on the level of the associated graded Hopf algebras. The enveloping algebra of a Lie algebra is replaced by a braided Hopf algebra which is generated by primitive elements.

In the case when the coradical  $A_0$  of  $A$  is not a Hopf subalgebra very little is known. There are a few results on the classification of arbitrary Hopf algebras of a given dimension such as for dimension  $p$  or  $p^2$  with prime  $p$ . For their proof only difficult ad hoc methods are used.

To formulate our main result we first describe the data  $\mathcal{D}, \lambda, \mu$  we need to define the Hopf algebras of the class we are considering. We fix a finite abelian group  $\Gamma$ .

**The datum  $\mathcal{D}$ .** A datum  $\mathcal{D}$  of finite Cartan type for  $\Gamma$ ,

$$\mathcal{D} = \mathcal{D}(\Gamma, (g_i)_{1 \leq i \leq \theta}, (\chi_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta}),$$

consists of elements  $g_i \in \Gamma, \chi_i \in \widehat{\Gamma}, 1 \leq i \leq \theta$ , and a Cartan matrix  $(a_{ij})_{1 \leq i, j \leq \theta}$  of finite type satisfying

$$(0.1) \quad q_{ij}q_{ji} = q_{ii}^{a_{ij}}, \quad q_{ii} \neq 1, \quad \text{with } q_{ij} = \chi_j(g_i) \text{ for all } 1 \leq i, j \leq \theta.$$

The Cartan condition (0.1) implies in particular,

$$(0.2) \quad q_{ii}^{a_{ij}} = q_{jj}^{a_{ji}} \text{ for all } 1 \leq i, j \leq \theta.$$

The explicit classification of all data of finite Cartan type for a given finite abelian group  $\Gamma$  is a computational problem. But at least it is a finite problem since the size  $\theta$  of the Cartan matrix is bounded by  $2(\text{ord}(\Gamma))^2$  by [AS2, 8.1], if  $\Gamma$  is an abelian group of odd order. For groups of prime order, all possibilities for  $\mathcal{D}$  are listed in [AS2].

Let  $\Phi$  be the root system of the Cartan matrix  $(a_{ij})_{1 \leq i, j \leq \theta}$ ,  $\alpha_1, \dots, \alpha_\theta$  a system of simple roots, and  $\mathcal{X}$  the set of connected components of the Dynkin diagram of  $\Phi$ . Let  $\Phi_J, J \in \mathcal{X}$ , be the root system of the component  $J$ . We write  $i \sim j$ , if  $\alpha_i$  and  $\alpha_j$  are in the same connected component of the Dynkin diagram of  $\Phi$ . For a positive root  $\alpha = \sum_{i=1}^{\theta} n_i \alpha_i, n_i \in \mathbb{N} = \{0, 1, 2, \dots\}$ , for all  $i$ , we define

$$g_\alpha = \prod_{i=1}^{\theta} g_i^{n_i}, \quad \chi_\alpha = \prod_{i=1}^{\theta} \chi_i^{n_i}.$$

We assume that the order of  $q_{ii}$  is odd for all  $i$ , and that the order of  $q_{ii}$  is prime to 3 for all  $i$  in a connected component of type  $G_2$ . Then it

follows from (0.2) that the order  $N_i$  of  $q_{ii}$  is constant in each connected component  $J$ , and we define  $N_J = N_i$  for all  $i \in J$ .

**The parameter  $\lambda$ .** Let  $\lambda = (\lambda_{ij})_{1 \leq i < j \leq \theta, i \not\sim j}$  be a family of elements in  $k$  satisfying the following condition for all  $1 \leq i < j \leq \theta, i \not\sim j$ : If  $g_i g_j = 1$  or  $\chi_i \chi_j \neq \varepsilon$ , then  $\lambda_{ij} = 0$ .

**The parameter  $\mu$ .** Let  $\mu = (\mu_\alpha)_{\alpha \in \Phi^+}$  be a family of elements in  $k$  such that for all  $\alpha \in \Phi_J^+, J \in \mathcal{X}$ , if  $g_\alpha^{N_J} = 1$  or  $\chi_\alpha^{N_J} \neq \varepsilon$ , then  $\mu_\alpha = 0$ .

Thus  $\lambda$  and  $\mu$  are finite families of free parameters in  $k$ . We can normalize  $\lambda$  and assume that  $\lambda_{ij} = 1$ , if  $\lambda_{ij} \neq 0$ .

**The Hopf algebra  $u(\mathcal{D}, \lambda, \mu)$ .** The definition of  $u(\mathcal{D}, \lambda, \mu)$  in Section 4.2 can be summarized as follows. In Definition 2.13 we associate to any  $\mu$  and  $\alpha \in \Phi^+$  an element  $u_\alpha(\mu)$  in the group algebra  $k[\Gamma]$ . By construction,  $u_\alpha(\mu)$  lies in the augmentation ideal of  $k[g_i^{N_i} \mid 1 \leq i \leq \theta]$ . The braided adjoint action  $\text{ad}_c(x_i)$  of  $x_i$  is defined in (1.12), and the root vectors  $x_\alpha$  are explained in Section 2.1.

The Hopf algebra  $u(\mathcal{D}, \lambda, \mu)$  is generated as an algebra by the group  $\Gamma$ , that is, by generators of  $\Gamma$  satisfying the relations of the group, and  $x_1, \dots, x_\theta$ , with the relations:

$$\begin{aligned} (\text{Action of the group}) \quad & g x_i g^{-1} = \chi_i(g) x_i, \text{ for all } i, \text{ and all } g \in \Gamma, \\ (\text{Serre relations}) \quad & \text{ad}_c(x_i)^{1-a_{ij}}(x_j) = 0, \text{ for all } i \neq j, i \sim j, \\ (\text{Linking relations}) \quad & \text{ad}_c(x_i)(x_j) = \lambda_{ij}(1 - g_i g_j), \text{ for all } i < j, i \not\sim j, \\ (\text{Root vector relations}) \quad & x_\alpha^{N_J} = u_\alpha(\mu), \text{ for all } \alpha \in \Phi_J^+, J \in \mathcal{X}. \end{aligned}$$

The coalgebra structure is given by

$$\Delta(x_i) = g_i \otimes x_i + x_i \otimes 1, \quad \Delta(g) = g \otimes g, \text{ for all } 1 \leq i \leq \theta, g \in \Gamma.$$

Now we can formulate our main result.

**Classification Theorem 0.1.** (1) Let  $\mathcal{D}, \lambda$  and  $\mu$  as above. Assume that  $q_{ij}$  has odd order for all  $i, j$ , and that the order of  $q_{ii}$  is prime to 3 for all  $i$  in a connected component of type  $G_2$ . Then  $u(\mathcal{D}, \lambda, \mu)$  is a pointed Hopf algebra of dimension  $\prod_{J \in \mathcal{X}} N_J^{|\Phi_J^+|} |\Gamma|$ , and  $G(u(\mathcal{D}, \lambda, \mu)) = \Gamma$ .

(2) Let  $A$  be a finite-dimensional pointed Hopf algebra with abelian group  $\Gamma = G(A)$ . Assume that all prime divisors of the order of  $\Gamma$  are  $> 7$ . Then  $A \cong u(\mathcal{D}, \lambda, \mu)$  for some  $\mathcal{D}, \lambda, \mu$ .

Moreover, in Theorem 7.2 we determine all isomorphisms between the Hopf algebras  $u(\mathcal{D}, \lambda, \mu)$ .

Part (1) of Theorem 0.1 is shown in Theorem 4.4, and part (2) is a special case of Theorem 6.2.

In [AS4] we proved the Classification Theorem for groups of the form  $(\mathbb{Z}/(p))^s$ ,  $s \geq 1$ , where  $p$  is a prime number  $> 17$ . In this special case, all the elements  $\mu$  and  $u_\alpha(\mu)$  are zero. In [AS1] we proved part (1) of Theorem 0.1 for Dynkin diagrams whose connected components are of type  $A_1$ , and in [AS5] for Dynkin diagrams of type  $A_n$ ; in [D2] our construction was extended to Dynkin diagrams whose connected components are of type  $A_n$  for various  $n$ . In [BDR] the Hopf algebra  $u(\mathcal{D}, \lambda, \mu)$  was introduced for type  $B_2$ .

Our proof of Theorem 0.1 is based on [AS1, AS2, AS3, AS4, AS5], and on previous work on quantum groups in [dCK, dCP, L1, L2, L3, M1, Ro], in particular on Lusztig's theory of the small quantum groups. Another essential ingredient of our proof are the recent results of Heckenberger on Nichols algebras of diagonal type in [H1, H2, H3] which use Kharchenko's theory [K] of PBW-bases in braided Hopf algebras of diagonal type.

In [AS2, 1.4] we conjectured that any finite-dimensional pointed Hopf algebra (over an algebraically closed field of characteristic 0) is generated by group-like and skew-primitive elements. Our Classification Theorem and Theorem 6.2 confirm this conjecture for a large class of Hopf algebras.

Finally we note that the following analog of Cauchy's Theorem from group theory holds for the Hopf algebras  $A = u(\mathcal{D}, \lambda, \mu)$ : If  $p$  is a prime divisor of the dimension of  $A$ , then  $A$  contains a group-like element of order  $p$ . We conjecture that Cauchy's Theorem holds for all finite-dimensional pointed Hopf algebras.

## 1. BRAIDED HOPF ALGEBRAS

**1.1. Yetter-Drinfeld modules over abelian groups and the tensor algebra.** Let  $\Gamma$  be an abelian group, and  $\hat{\Gamma}$  the character group of all group homomorphisms from  $\Gamma$  to the multiplicative group  $k^\times$  of the field  $k$ . The braided category  ${}^\Gamma\mathcal{YD}$  of (left) Yetter-Drinfeld modules over  $\Gamma$  is the category of left  $k[\Gamma]$ -modules which are  $\Gamma$ -graded vector spaces  $V = \bigoplus_{g \in \Gamma} V_g$  such that each homogeneous component  $V_g$  is stable under the action of  $\Gamma$ . Morphisms are  $\Gamma$ -linear maps  $f : \bigoplus_{g \in \Gamma} V_g \rightarrow \bigoplus_{g \in \Gamma} W_g$  with  $f(V_g) \subset W_g$  for all  $g \in \Gamma$ . The  $\Gamma$ -grading is equivalent to a left  $k[\Gamma]$ -comodule structure  $\delta : V \rightarrow k[\Gamma] \otimes V$ , where

$\delta(v) = g \otimes v$  is equivalent to  $v \in V_g$ . We use a Sweedler notation  $\delta(v) = v_{(-1)} \otimes v_{(0)}$  for all  $v \in V$ .

If  $V = \bigoplus_{g \in \Gamma} V_g$  and  $W = \bigoplus_{g \in \Gamma} W_g$  are in  ${}^{\Gamma}\mathcal{YD}$ , the monoidal structure is given by the usual tensor product  $V \otimes W$  with  $\Gamma$ -action  $g(v \otimes w) = gv \otimes gw$ ,  $v \in V, w \in W$ , and  $\Gamma$ -grading  $(V \otimes W)_g = \bigoplus_{ab=g} V_a \otimes W_b$  for all  $g \in \Gamma$ . The braiding in  ${}^{\Gamma}\mathcal{YD}$  is the isomorphism

$$c = c_{V,W} : V \otimes W \rightarrow W \otimes V$$

defined by  $c(v \otimes w) = g \cdot w \otimes v$  for all  $g \in \Gamma, v \in V_g$ , and  $w \in W$ . Thus each Yetter-Drinfeld module  $V$  defines a braided vector space  $(V, c_{V,V})$ .

If  $\chi$  is a character of  $\Gamma$  and  $V$  a left  $\Gamma$ -module, we define

$$V^{\chi} := \{v \in V \mid g \cdot v = \chi(g)v \text{ for all } g \in \Gamma\}.$$

Let  $\theta \geq 1$  be a natural number,  $g_1, \dots, g_{\theta} \in \Gamma$ , and  $\chi_1, \dots, \chi_{\theta} \in \widehat{\Gamma}$ . Let  $V$  be a vector space with basis  $x_1, \dots, x_{\theta}$ .  $V$  is an object in  ${}^{\Gamma}\mathcal{YD}$  by defining  $x_i \in V_{g_i}^{\chi_i}$  for all  $i$ . Thus each  $x_i$  has degree  $g_i$ , and the group  $\Gamma$  acts on  $x_i$  via the character  $\chi_i$ . We define

$$q_{ij} := \chi_j(g_i) \text{ for all } 1 \leq i, j \leq \theta.$$

The braiding on  $V$  is determined by the matrix  $(q_{ij})$  since

$$c(x_i \otimes x_j) = q_{ij} x_j \otimes x_i \text{ for all } 1 \leq i, j \leq \theta.$$

We will identify the tensor algebra  $T(V)$  with the free associative algebra  $k\langle x_1, \dots, x_{\theta} \rangle$ . It is an algebra in  ${}^{\Gamma}\mathcal{YD}$ , where a monomial

$$x = x_{i_1} x_{i_2} \cdots x_{i_n}, 1 \leq i_1, \dots, i_n \leq \theta,$$

has  $\Gamma$ -degree  $g_{i_1} g_{i_2} \cdots g_{i_n}$  and the action of  $g \in \Gamma$  on  $x$  is given by  $g \cdot x = \chi_{i_1} \chi_{i_2} \cdots \chi_{i_n}(g)x$ .  $T(V)$  is a braided Hopf algebra in  ${}^{\Gamma}\mathcal{YD}$  with comultiplication

$$\Delta_{T(V)} : T(V) \rightarrow T(V) \underline{\otimes} T(V), \quad x_i \mapsto x_i \otimes 1 + 1 \otimes x_i, \quad 1 \leq i \leq \theta.$$

Here we write  $T(V) \underline{\otimes} T(V)$  to indicate the braided algebra structure on the vector space  $T(V) \otimes T(V)$ , that is

$$(x \otimes y)(x' \otimes y') = x(g \cdot x') \otimes yy',$$

for all  $x, x', y, y' \in T(V)$  and  $y \in T(V)_g, g \in \Gamma$ .

Let  $I = \{1, 2, \dots, \theta\}$ , and  $\mathbb{Z}[I]$  the free abelian group of rank  $\theta$  with basis  $\alpha_1, \dots, \alpha_{\theta}$ . Given the matrix  $(q_{ij})$ , we define the bilinear map

$$(1.1) \quad \mathbb{Z}[I] \times \mathbb{Z}[I] \rightarrow k^{\times}, \quad (\alpha, \beta) \mapsto q_{\alpha, \beta}, \text{ by } q_{\alpha_i, \alpha_j} = q_{ij}, 1 \leq i, j \leq \theta.$$

We consider  $V$  as a Yetter-Drinfeld module over  $\mathbb{Z}[I]$  by defining  $x_i \in V_{\alpha_i}^{\psi_i}$  for all  $1 \leq i \leq \theta$ , where  $\psi_j$  is the character of  $\mathbb{Z}[I]$  with

$$\psi_j(\alpha_i) = q_{ij} \text{ for all } 1 \leq i, j \leq \theta.$$

Thus  $T(V) = k\langle x_1, \dots, x_\theta \rangle$  is also a braided Hopf algebra in  ${}_{\mathbb{Z}[I]}^{\mathbb{Z}[I]}\mathcal{YD}$ . The  $\mathbb{Z}[I]$ -degree of a monomial  $x = x_{i_1}x_{i_2}\cdots x_{i_n}$ ,  $1 \leq i_1, \dots, i_n \leq \theta$ , is  $\sum_{i=1}^\theta n_i \alpha_i$ , where for all  $i$ ,  $n_i$  is the number of occurrences of  $i$  in the sequence  $(i_1, i_2, \dots, i_n)$ . The braiding on  $T(V)$  as a Yetter-Drinfeld module over  $\Gamma$  or  $\mathbb{Z}[I]$  is in both cases given by

$$(1.2) \quad c(x \otimes y) = q_{\alpha, \beta} y \otimes x, \text{ where } x \in T(V)_\alpha, y \in T(V)_\beta, \alpha, \beta \in \mathbb{Z}[I].$$

The comultiplication of  $T(V)$  as a braided Hopf algebra in  ${}_{\Gamma}\mathcal{YD}$  only depends on the matrix  $(q_{ij})$ , hence it coincides with the comultiplication of  $T(V)$  as a coalgebra in  ${}_{\mathbb{Z}[I]}^{\mathbb{Z}[I]}\mathcal{YD}$ . In particular, the comultiplication of  $T(V)$  is  $\mathbb{Z}[I]$ -graded.

**1.2. Bosonization and twisting.** Let  $R$  be a braided Hopf algebra in  ${}_{\Gamma}\mathcal{YD}$ . We will use a Sweedler notation for the comultiplication

$$\Delta_R : R \rightarrow R \otimes R, \quad \Delta_R(r) = r^{(1)} \otimes r^{(2)}.$$

For Hopf algebras  $A$  in the usual sense, we always use the Sweedler notation

$$\Delta : A \rightarrow A \otimes A, \quad \Delta(a) = a_{(1)} \otimes a_{(2)}.$$

Then the smash product  $A = R \# k[\Gamma]$  is a Hopf algebra in the usual sense (the bosonization of  $R$ ). As vector spaces,  $R \# k[\Gamma] = R \otimes k[\Gamma]$ . Multiplication and comultiplication are defined by

$$(1.3) \quad (r \# g)(s \# h) = r(g \cdot s) \# gh, \quad \Delta(r \# g) = r^{(1)} \# r^{(2)}_{(-1)} g \otimes r^{(2)}_{(0)} \# g.$$

Then the maps

$$\iota : k[\Gamma] \rightarrow R \# k[\Gamma], \text{ and } \pi : R \# k[\Gamma] \rightarrow k[\Gamma]$$

with  $\iota(g) = 1 \# g$  and  $\pi(r \# g) = r$  for all  $r \in R, g \in \Gamma$  are Hopf algebra maps with  $\pi \iota = \text{id}$ .

Conversely, if  $A$  is a Hopf algebra in the usual sense with Hopf algebra maps  $\iota : k[\Gamma] \rightarrow A$  and  $\pi : A \rightarrow k[\Gamma]$  such that  $\pi \iota = \text{id}$ , then

$$R = \{a \in A \mid (\text{id} \otimes \pi)\Delta(a) = a \otimes 1\}$$

is a braided Hopf algebra in  ${}_{\Gamma}\mathcal{YD}$  in the following way. As an algebra,  $R$  is a subalgebra of  $A$ . The  $k[\Gamma]$ -coaction,  $\Gamma$ -action and comultiplication of  $R$  are defined by

$$(1.4) \quad \delta(r) = \pi(r^{(1)}) \otimes r^{(2)}, \quad g \cdot r = \iota(g)r\iota(g^{-1})$$

and

$$(1.5) \quad \Delta_R(r) = \vartheta(r_{(1)}) \otimes r_{(2)}.$$

Here,  $\Delta_A(r) = r_{(1)} \otimes r_{(2)}$ , and  $\vartheta$  is the map

$$(1.6) \quad \vartheta : A \rightarrow R, \quad \vartheta(r) = r_{(1)}\iota(S(\pi(r_{(2)}))),$$

where  $S$  is the antipode of  $A$ . Then

$$(1.7) \quad R\#k[\Gamma] \rightarrow A, \quad r\#g \mapsto r\iota(g), \quad r \in R, g \in \Gamma,$$

is an isomorphism of Hopf algebras.

We recall the notion of *twisting* the algebra structure of an arbitrary Hopf algebra  $A$ , see for example [KS, 10.2.3]. Let  $\sigma : A \otimes A \rightarrow k$  be a convolution invertible linear map, and a normalized 2-cocycle, that is, for all  $x, y, z \in A$ ,

$$(1.8) \quad \sigma(x_{(1)}, y_{(1)})\sigma(x_{(2)}y_{(2)}, z) = \sigma(y_{(1)}, z_{(1)})\sigma(x, y_{(2)}z_{(2)}),$$

and  $\sigma(x, 1) = \varepsilon(x) = \sigma(1, x)$ . The Hopf algebra  $A_\sigma$  with twisted algebra structure is equal to  $A$  as a coalgebra, and has multiplication  $\cdot_\sigma$  with

$$(1.9) \quad x \cdot_\sigma y = \sigma(x_{(1)}, y_{(1)})x_{(2)}y_{(2)}\sigma^{-1}(x_{(3)}, y_{(3)}) \text{ for all } x, y \in A.$$

In the situation  $A = R\#k[\Gamma]$  above, let  $\sigma : \Gamma \times \Gamma \rightarrow k^\times$  be a normalized 2-cocycle of the group  $\Gamma$ . Then  $\sigma$  extends to a 2-cocycle of the group algebra  $k[\Gamma]$  and it defines a normalized and invertible 2-cocycle  $\sigma_\pi = \sigma(\pi \otimes \pi)$  of the Hopf algebra  $A$ . Since  $k[\Gamma]$  is cocommutative,  $\iota$  and  $\pi$  are Hopf algebra maps

$$\iota : k[\Gamma] \rightarrow A_{\sigma_\pi} \text{ and } \pi : A_{\sigma_\pi} \rightarrow k[\Gamma].$$

Hence the coinvariant elements

$$R_\sigma = \{a \in A_{\sigma_\pi} \mid (\text{id} \otimes \pi)\Delta(a) = a \otimes 1\}$$

form a braided Hopf algebra in  ${}^\Gamma_1\mathcal{YD}$ . As a vector space,  $R_\sigma$  coincides with  $R$ , but  $R_\sigma$  and  $R$  have different multiplication and comultiplication.

To simplify the formulas, we will treat  $\iota$  as an inclusion map.

In any braided Hopf algebra  $R$  with multiplication  $m$  and braiding  $c : R \otimes R \rightarrow R \otimes R$  we define the *braided commutator* of elements  $x, y \in R$  by

$$(1.10) \quad [x, y]_c = xy - mc(x \otimes y).$$

If  $x \in R$  is a primitive element, then

$$(1.11) \quad (\text{ad}_c x)(y) = [x, y]_c$$

denotes the *braided adjoint action* of  $x$  on  $R$ . For example, in the situation of the free algebra in Section 1.1 with braiding (1.2), we have for all  $x_i$  and  $y = x_{j_1} \cdots x_{j_n}$ ,

$$(1.12) \quad (\text{ad}_c x_i)(y) = x_i y - q_{i j_1} \cdots q_{i j_n} y x_i.$$

In the formulation of the next lemma we need one more notation. If  $V$  is a left  $C$ -comodule over a coalgebra  $C$ , then  $V$  is a right module over the dual algebra  $C^*$  by  $v \leftarrow p = p(v_{(-1)})v_{(0)}$  for all  $v \in V, p \in C^*$ .



In particular, if  $R$  is a braided Hopf algebra in  ${}^{\Gamma}\mathcal{YD}$ , then the  $k[\Gamma]$ -coaction defines a left  $k[\Gamma] \otimes k[\Gamma]$ -comodule structure on  $R \otimes R$ , hence a right  $(k[\Gamma] \otimes k[\Gamma])^*$ -module structure on  $R \otimes R$  denoted by  $\leftarrow$ .

**Lemma 1.1.** *Let  $\Gamma$  be an abelian group,  $\sigma : \Gamma \times \Gamma \rightarrow k^{\times}$  a normalized 2-cocycle,  $R$  a braided Hopf algebra in  ${}^{\Gamma}\mathcal{YD}$ ,  $g, h \in \Gamma$ , and  $x \in R_g, y \in R_h, r \in R$ .*

- (1)  $x \cdot_{\sigma} y = \sigma(g, h)xy$ .
- (2)  $\Delta_{R_{\sigma}}(r) = \Delta_R(r) \leftarrow \sigma^{-1}$ .
- (3) *If  $y \in R_h^{\eta}$  for some character  $\eta \in \widehat{\Gamma}$ , and  $R$  as an algebra is generated by primitive elements, then  $g \cdot_{\sigma} y = \sigma(g, h)\sigma^{-1}(h, g)\eta(g)y$ , and hence  $[x, y]_{c_{\sigma}} = \sigma(g, h)[x, y]_{\sigma}$ .*

*Proof.* (1) and (3) are [AS5, (2-11), (2-14)]. To prove (2), using the cocommutativity of the group algebra we compute

$$\begin{aligned} \Delta_{R_{\sigma}}(r) &= r_{(1)} \cdot_{\sigma} S(\pi(r_{(2)})) \otimes r_{(3)} \\ &= \sigma(\pi(r_{(1)}), S(\pi(r_{(2)})))\vartheta(r_{(2)})\sigma^{-1}(\pi(r_{(3)}), S(\pi(r_{(4)}))) \otimes r_{(6)}. \end{aligned}$$

On the other hand,  $\Delta_R(r) = r_{(1)}S\pi(r_{(2)}) \otimes r_{(3)}$ , hence

$$r_{(1)}^{(1)}{}_{(-1)} \otimes r_{(2)}^{(2)}{}_{(-1)} \otimes r_{(1)}^{(1)}{}_{(0)} \otimes r_{(2)}^{(2)}{}_{(0)} = \pi(r_{(1)}S(r_{(3)})) \otimes \pi(r_{(4)}) \otimes \vartheta(r_{(2)}) \otimes r_{(5)},$$

and

$\Delta_R(r) \leftarrow \sigma^{-1} = \sigma^{-1}(\pi(r_{(1)}S(r_{(3)})), \pi(r_{(4)}))\vartheta(r_{(2)}) \otimes r_{(5)}$ . Hence the claim follows from the equality

$$\sigma(a, S(b_{(3)}))\sigma^{-1}(b_{(1)}, S(b_{(2)})) = \sigma^{-1}(aS(b_{(1)}), b_{(2)})$$

for all  $a, b \in k[\Gamma]$ . It is enough to check this equation for elements  $a, b \in \Gamma$ . Then the equality follows from the group cocycle condition.  $\square$

We now apply the twisting procedure to the braided Hopf algebra  $T(V) \in {}_{\mathbb{Z}[I]}^{\mathbb{Z}[I]}\mathcal{YD}$ .

**Lemma 1.2.** *Let  $\theta \geq 1$ , and  $(q_{ij})_{1 \leq i, j \leq \theta}, (q'_{ij})_{1 \leq i, j \leq \theta}$  matrices with coefficients in  $k$ . Let  $V \in {}_{\mathbb{Z}[I]}^{\mathbb{Z}[I]}\mathcal{YD}$  with basis  $x_1, \dots, x_{\theta}$  and  $x_i \in V_{\alpha_i}^{\psi_i}, \psi_j(\alpha_i) = q_{ij}$  for all  $i, j$  as in Section 1.1, and  $V' \in {}_{\mathbb{Z}[I]}^{\mathbb{Z}[I]}\mathcal{YD}$  with basis  $x'_1, \dots, x'_{\theta}$  and  $x'_i \in V_{\alpha_i}^{\psi'_i}, \psi'_j(\alpha_i) = q'_{ij}$  for all  $i, j$ . Then  $T(V)$  and  $T(V')$  are braided Hopf algebras in  ${}_{\mathbb{Z}[I]}^{\mathbb{Z}[I]}\mathcal{YD}$  as in Section 1.1. Assume*

$$(1.13) \quad q_{ij}q_{ji} = q'_{ij}q'_{ji}, \text{ and } q_{ii} = q'_{ii} \text{ for all } 1 \leq i, j \leq \theta.$$

*Then there is a 2-cocycle  $\sigma : \mathbb{Z}[I] \times \mathbb{Z}[I] \rightarrow k^{\times}$  with*

$$(1.14) \quad \sigma(\alpha, \beta)\sigma^{-1}(\beta, \alpha) = q_{\alpha\beta}q'_{\alpha\beta}{}^{-1} \text{ for all } \alpha, \beta \in \mathbb{Z}[I],$$

*and a  $k$ -linear isomorphism  $\varphi : T(V) \rightarrow T(V')$  with  $\varphi(x_i) = x'_i$  for all  $i$  and such that for all  $\alpha, \beta \in \mathbb{Z}[I], x \in T(V)_{\alpha}, y \in T(V)_{\beta}$  and  $z \in T(V)$*

- (1)  $\varphi(xy) = \sigma(\alpha, \beta)\varphi(x)\varphi(y)$ .
- (2)  $\Delta_{T(V')}(\varphi(z)) = (\varphi \otimes \varphi)(\Delta_{T(V)}(z)) \leftarrow \sigma$ .
- (3)  $\varphi([x, y]_c) = \sigma(\alpha, \beta)[\varphi(x), \varphi(y)]_{c'}$ .

*Proof.* Define  $\sigma$  as the bilinear map with  $\sigma(\alpha_i, \alpha_j) = q_{ij}q'_{ij}{}^{-1}$  if  $i \leq j$ , and  $\sigma(\alpha_i, \alpha_j) = 1$  if  $i > j$  (see [AS5, Prop. 3.9]).

Let  $\varphi : T(V) \rightarrow T(V')_\sigma$  be the algebra map with  $\varphi(x_i) = x'_i$  for all  $i$ . Then  $\varphi$  is bijective since it follows from Lemma 1.1 (1) and the bilinearity of  $\sigma$  that for all monomials  $x = x_{i_1}x_{i_2} \cdots x_{i_n}$  of length  $n \geq 1$  with  $x' = x'_{i_1}x'_{i_2} \cdots x'_{i_n}$ ,

$$\varphi(x) = \prod_{r < s} \sigma(\alpha_{i_r}, \alpha_{i_s}) x'.$$

In particular,  $\varphi$  is  $\mathbb{Z}[I]$ -graded. To see that  $\varphi$  is  $\mathbb{Z}[I]$ -linear, let  $\alpha, \beta \in \mathbb{Z}[I]$  and  $x \in T(V)_\beta$ . Then by Lemma 1.1 (3),

$$\alpha \cdot x = q_{\alpha\beta}x, \text{ and } \alpha \cdot_\sigma \varphi(x) = \sigma(\alpha, \beta)\sigma^{-1}(\beta, \alpha)q'_{\alpha\beta}\varphi(x),$$

and  $\varphi(\alpha \cdot x) = \alpha \cdot_\sigma \varphi(x)$  follows by (1.14). Since the elements  $x_i$  and  $x'_i$  are primitive we now see that  $\varphi : T(V) \rightarrow T(V')_\sigma$  is an isomorphism of braided Hopf algebras. Then the claim follows from Lemma 1.1.  $\square$

## 2. SERRE RELATIONS AND ROOT VECTORS

### 2.1. Datum of finite Cartan type and root vectors.

**Definition 2.1.** A *datum of Cartan type*

$$\mathcal{D} = \mathcal{D}(\Gamma, (g_i)_{1 \leq i \leq \theta}, (\chi_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta})$$

consists of an abelian group  $\Gamma$ , elements  $g_i \in \Gamma$ ,  $\chi_i \in \widehat{\Gamma}$ ,  $1 \leq i \leq \theta$ , and a Cartan matrix  $(a_{ij})$  of size  $\theta$  satisfying

$$(2.1) \quad q_{ij}q_{ji} = q_{ii}^{a_{ij}}, \quad q_{ii} \neq 1, \text{ with } q_{ij} = \chi_j(g_i) \text{ for all } 1 \leq i, j \leq \theta.$$

We call  $\theta$  the *rank* of  $\mathcal{D}$ . A datum  $\mathcal{D}$  of Cartan type will be called of finite Cartan type if  $(a_{ij})$  is of finite type.

**Example 2.2.** A Cartan datum  $(I, \cdot)$  in the sense of Lusztig [L3, 1.1.1] defines a datum of Cartan type for the free abelian group  $ZI$  with  $g_i = \alpha_i$ ,  $\chi_i = \psi_i$ ,  $1 \leq i \leq \theta$ , as in Section 1.1, where

$$q_{ij} = v^{d_i a_{ij}}, \quad d_i = \frac{i \cdot i}{2}, \quad a_{ij} = 2 \frac{i \cdot j}{i \cdot i} \text{ for all } 1 \leq i, j \leq \theta.$$

In Example 2.2,  $d_i a_{ij} = i \cdot j$  is the symmetrized Cartan matrix, and  $q_{ij} = q_{ji}$  for all  $1 \leq i, j \leq \theta$ . In general, the matrix  $(q_{ij})$  of a datum of Cartan type is not symmetric, but by Lemma 1.2 we can reduce to the symmetric case by twisting.

We fix a finite abelian group  $\Gamma$  and a datum

$$\mathcal{D} = \mathcal{D}(\Gamma, (g_i)_{1 \leq i \leq \theta}, (\chi_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta})$$

of finite Cartan type. The Weyl group  $W \subset \text{Aut}(\mathbb{Z}[I])$  of  $(a_{ij})$  is generated by the reflections  $s_i : \mathbb{Z}[I] \rightarrow \mathbb{Z}[I]$  with  $s_i(\alpha_j) = \alpha_j - a_{ij}\alpha_i$  for all  $i, j$ . The root system is  $\Phi = \cup_{i=1}^{\theta} W(\alpha_i)$ , and

$$\Phi^+ = \left\{ \alpha \in \Phi \mid \alpha = \sum_{i=1}^{\theta} n_i \alpha_i, n_i \geq 0 \text{ for all } 1 \leq i \leq \theta \right\}$$

denotes the set of positive roots with respect to the basis of simple roots  $\alpha_1, \dots, \alpha_{\theta}$ . Let  $p$  be the number of positive roots.

For  $\alpha = \sum_{i=1}^{\theta} n_i \alpha_i \in \mathbb{Z}[I]$ ,  $n_i \in \mathbb{Z}$  for all  $i$  we define

$$(2.2) \quad g_{\alpha} = g_1^{n_1} g_2^{n_2} \cdots g_{\theta}^{n_{\theta}} \text{ and } \chi_{\alpha} = \chi_1^{n_1} \chi_2^{n_2} \cdots \chi_{\theta}^{n_{\theta}}.$$

In this section, we assume that the Dynkin diagram of  $(a_{ij})$  is *connected*. In this case we say that  $\mathcal{D}$  is connected.

We fix a reduced decomposition of the longest element

$$w_0 = s_{i_1} s_{i_2} \cdots s_{i_p}$$

of  $W$  in terms of the simple reflections. Then

$$\beta_l = s_{i_1} \cdots s_{i_{l-1}}(\alpha_{i_l}), 1 \leq l \leq p,$$

is a convex ordering of the positive roots.

Let  $d_1, \dots, d_{\theta} \in \{1, 2, 3\}$  such that  $d_i a_{ij} = d_j a_{ji}$  for all  $i, j$ . We assume for all  $1 \leq i, j \leq \theta$ ,

$$(2.3) \quad q_{ij} \text{ has odd order, and}$$

$$(2.4) \quad \text{the order of } q_{ii} \text{ is prime to 3, if } (a_{ij}) \text{ is of type } G_2.$$

Then it follows from (2.1) ([AS2, 4.3]) that the elements  $q_{ii}$  have the same order in  $k^{\times}$ . We define

$$(2.5) \quad N = \text{order of } q_{ii}, 1 \leq i \leq \theta.$$

**Definition 2.3.** Let  $V = V(\mathcal{D})$  be a vector space with basis  $x_1, \dots, x_{\theta}$ , and let  $V \in {}_{\Gamma}^{\Gamma} \mathcal{YD}$  by  $x_i \in V_{g_i}^{x_i}$  for all  $1 \leq i \leq \theta$ . Then  $T(V)$  is a braided Hopf algebra in  ${}_{\Gamma}^{\Gamma} \mathcal{YD}$  as in Section 1.1. Let

$$R(\mathcal{D}) = T(V) / ((\text{ad}_c x_i)^{1-a_{ij}}(x_j) \mid 1 \leq i, j \leq \theta)$$

be the quotient Hopf algebra in  ${}_{\Gamma}^{\Gamma} \mathcal{YD}$ .

It is well-known that the elements  $(\text{ad}_c x_i)^{1-a_{ij}}(x_j)$ ,  $1 \leq i, j \leq \theta$  are primitive in the free algebra  $T(V)$  (see for example [AS2, A.1]), hence they generate a Hopf ideal. By abuse of language, we denote the images of the elements  $x_i$  in  $R(\mathcal{D})$  again by  $x_i$ .

In the situation of Example 2.2, Lusztig [L2] defined root vectors  $x_\alpha$  in  $R(\mathcal{D}) = U^+$  for each positive root  $\alpha$  using the convex ordering of the positive roots. As noted in [AS4], these root vectors can be seen to be iterated braided commutators of the elements  $x_1, \dots, x_\theta$  with respect to the braiding given by the matrix  $(v^{d_i a_{ij}})$ . This follows for example from the inductive definition of the root vectors in [Ri].

In the case of our general braiding given by  $(q_{ij})$  we define root vectors  $x_\alpha \in R(\mathcal{D})$  for each  $\alpha \in \Phi^+$  by the same iterated braided commutator of the elements  $x_1, \dots, x_\theta$  as in Lusztig's case but with respect to the general braiding.

**Definition 2.4.** Let  $K(\mathcal{D})$  be the subalgebra of  $R(\mathcal{D})$  generated by the elements  $x_\alpha^N, \alpha \in \Phi^+$ .

**Theorem 2.5.** Let  $\mathcal{D}$  be a connected datum of finite Cartan type, and assume (2.3), (2.4).

(1) The elements

$$x_{\beta_1}^{a_1} x_{\beta_2}^{a_2} \cdots x_{\beta_p}^{a_p}, a_1, a_2, \dots, a_p \geq 0,$$

form a basis of  $R(\mathcal{D})$ .

(2)  $K(\mathcal{D})$  is a braided Hopf subalgebra of  $R(\mathcal{D})$ .

(3) For all  $\alpha, \beta \in \Phi^+$ ,  $x_\alpha x_\beta^N = \chi_\beta^N(g_\alpha) x_\beta^N x_\alpha$ , that is,  $[x_\alpha, x_\beta^N]_c = 0$ .

*Proof.* (a) In the situation of 2.2, the elements in (1) form Lusztig's PBW-basis of  $U^+$  over  $\mathbb{Z}[v, v^{-1}]$  by [L2, 5.7].

(b) Now we assume that the braiding has the form  $(q_{ij} = q^{d_i a_{ij}})$ , where  $(d_i a_{ij})$  is the symmetrized Cartan matrix, and  $q$  is a non-zero element in  $k$  of odd order, and not divisible by 3 if the Dynkin diagram of  $(a_{ij})$  is  $G_2$ . Then (1) follows from Lusztig's result by extension of scalars, and (2) is shown in [dCP, 19.1] (for another proof see [M2, 3.1]). The algebra  $K(\mathcal{D})$  is commutative since it is a subalgebra of the commutative algebra  $Z_0$  of [dCP, 19.1]. This proves (3) since  $q^N = 1$ , hence  $\chi_\beta^N(g_\alpha) = 1$ .

(c) In the situation of a general braiding matrix  $(q_{ij})_{1 \leq i, j \leq \theta}$  assumed in the theorem, we define a matrix  $(q'_{ij})_{1 \leq i, j \leq \theta}$  by  $q'_{ii} = q_{ii}$  for all  $i$ , and for all  $i \neq j$  we define  $q'_{ij} = q'_{ji}$  to be a square root of  $q_{ij} q_{ji}$ . By [AS2, 4.3],  $q'_{ij} = q^{d_i a_{ij}}$  for all  $i, j$ , and for some  $q \in k$ . Thus by part (b) of the proof, (1), (2) and (3) hold for the braiding  $(q'_{ij})$ , and hence by Lemma 1.2 for  $(q_{ij})$ .  $\square$

**2.2. The Hopf algebra  $K(\mathcal{D}) \# k[\Gamma]$ .** We assume the situation of Section 2.1. By Theorem 2.5 (2),  $K(\mathcal{D})$  is a braided Hopf algebra in  ${}^\Gamma \mathcal{YD}$ ,

and the smash product  $K(\mathcal{D})\#k[\Gamma]$  is a Hopf algebra in the usual sense. We want to describe all Hopf algebra maps

$$K(\mathcal{D})\#k[\Gamma] \rightarrow k[\Gamma]$$

which are the identity on the group algebra  $k[\Gamma]$ .

**Definition 2.6.** For any  $1 \leq l \leq p$  and  $a = (a_1, a_2, \dots, a_p) \in \mathbb{N}^p$  we define

$$\begin{aligned} h_l &= g_{\beta_l}^N, \\ \eta_l &= \chi_{\beta_l}^N, \\ z_l &= x_{\beta_l}^N, \\ z^a &= z_1^{a_1} z_2^{a_2} \cdots z_p^{a_p} \in K(\mathcal{D}), \\ h^a &= h_1^{a_1} h_2^{a_2} \cdots h_p^{a_p} \in \Gamma, \\ \eta^a &= \eta_1^{a_1} \eta_2^{a_2} \cdots \eta_p^{a_p} \in \widehat{\Gamma}, \\ \underline{a} &= a_1 \beta_1 + a_2 \beta_2 + \cdots + a_p \beta_p \in \mathbb{Z}[I]. \end{aligned}$$

For  $\alpha = \sum_{i=1}^{\theta} n_i \alpha_i \in \mathbb{Z}[I]$ ,  $n_i \in \mathbb{Z}$  for all  $i$ , we call  $\text{ht}(\alpha) = \sum_{i=1}^{\theta} n_i$  the *height* of  $\alpha$ . Let  $e_l = (\delta_{kl})_{1 \leq k \leq p} \in \mathbb{N}^p$ , where  $\delta_{kl} = 1$  if  $k = l$  and  $\delta_{kl} = 0$  if  $k \neq l$ .

Note that for all  $a, b, c \in \mathbb{N}^p$ ,

$$(2.6) \quad h^a = h^b h^c, \quad \eta^a = \eta^b \eta^c, \quad \text{if } \underline{a} = \underline{b} + \underline{c},$$

$$(2.7) \quad \text{ht}(\underline{b}) < \text{ht}(\underline{a}), \quad \text{if } \underline{a} = \underline{b} + \underline{c} \text{ and } c \neq 0.$$

As explained in Section 1.1, we view  $T(V)$  as a braided Hopf algebra in  $\frac{\mathbb{Z}[I]}{\mathbb{Z}[I]} \mathcal{YD}$ . Then the quotient Hopf algebra  $R(\mathcal{D})$  and its Hopf subalgebra  $K(\mathcal{D})$  are braided Hopf algebras in  $\frac{\mathbb{Z}[I]}{\mathbb{Z}[I]} \mathcal{YD}$ . In particular, the comultiplication  $\Delta_{K(\mathcal{D})} : K(\mathcal{D}) \rightarrow K(\mathcal{D}) \otimes K(\mathcal{D})$  is  $\mathbb{Z}[I]$ -graded. By construction, for any  $\alpha \in \Phi^+$ , the root vector  $x_\alpha$  in  $R(\mathcal{D})$  is  $\mathbb{Z}[I]$ -homogeneous of  $\mathbb{Z}[I]$ -degree  $\alpha$ . Thus  $x_\alpha \in R(\mathcal{D})_{g_\alpha}^{\chi_\alpha}$ , and for all  $a \in \mathbb{N}^p$ ,  $z^a$  has  $\mathbb{Z}[I]$ -degree  $N\underline{a}$ , and

$$(2.8) \quad z^a \in K(\mathcal{D})_{h^a}^{\eta^a}.$$

For  $z \in K(\mathcal{D})$ ,  $g \in \Gamma$ , we will denote  $z\#g \in K(\mathcal{D})\#k[\Gamma]$  by  $zg$ . By Theorem 2.5 the elements  $z^a g$  with  $a \in \mathbb{N}^p$ ,  $g \in \Gamma$ , form a basis of  $K(\mathcal{D})\#k[\Gamma]$ , and it follows that for all  $a, b = (b_i), c = (c_i) \in \mathbb{N}^p$ ,

$$(2.9) \quad z^b z^c = \gamma_{b,c} z^{b+c}, \quad \text{where } \gamma_{b,c} = \prod_{k>l} \eta_l(h_k)^{b_k c_l},$$

$$(2.10) \quad h^a z^b = \eta^b(h^a) z^b h^a \text{ in } R \# k[\Gamma].$$

**Lemma 2.7.** *For any  $0 \neq a \in \mathbb{N}^p$  there are uniquely determined scalars  $t_{b,c}^a \in k$ ,  $0 \neq b, c \in \mathbb{N}^p$ , such that*

$$(2.11) \quad \Delta_{K(\mathcal{D})}(z^a) = z^a \otimes 1 + 1 \otimes z^a + \sum_{b,c \neq 0, \underline{b} + \underline{c} = \underline{a}} t_{b,c}^a z^b \otimes z^c.$$

*Proof.* Since  $\Delta_{K(\mathcal{D})}$  is  $\mathbb{Z}[I]$ -graded,  $\Delta_{K(\mathcal{D})}(z^a)$  is a linear combination of elements  $z^b \otimes z^c$  where  $\underline{b} + \underline{c} = \underline{a}$ . Hence

$$\Delta_{K(\mathcal{D})}(z^a) = x \otimes 1 + 1 \otimes y + \sum_{b,c \neq 0, \underline{b} + \underline{c} = \underline{a}} t_{b,c}^a z^b \otimes z^c,$$

where  $x, y$  are elements in  $K(\mathcal{D})$ . By applying the augmentation  $\varepsilon$  it follows that  $x = y = z^a$ .  $\square$

We now define recursively a family of elements  $u^a$  in  $k[\Gamma]$  depending on parameters  $\mu_a$  which behave like the elements  $z^a$  with respect to comultiplication.

**Lemma 2.8.** *Let  $n \geq 2$ . For all  $0 \neq b \in \mathbb{N}^p$ ,  $\text{ht}(\underline{b}) < n$ , let  $\mu_b \in k$  and  $u^b \in k[\Gamma]$  such that*

$$(2.12) \quad u^b = \mu_b(1 - h^b) + \sum_{d,e \neq 0, \underline{d} + \underline{e} = \underline{b}} t_{d,e}^b \mu_d u^e,$$

$$(2.13) \quad \Delta(u^b) = h^b \otimes u^b + u^b \otimes 1 + \sum_{d,e \neq 0, \underline{d} + \underline{e} = \underline{b}} t_{d,e}^b u^d h^e \otimes u^e.$$

*Let  $a \in \mathbb{N}^p$  with  $\text{ht}(\underline{a}) = n$ , and  $u^a \in k[\Gamma]$ . Then the following statements are equivalent:*

$$(2.14) \quad u^a = \mu_a(1 - h^a) + \sum_{b,c \neq 0, \underline{b} + \underline{c} = \underline{a}} t_{b,c}^a \mu_b u^c \text{ for some } \mu_a \in k.$$

$$(2.15) \quad \Delta(u^a) = h^a \otimes u^a + u^a \otimes 1 + \sum_{b,c \neq 0, \underline{b} + \underline{c} = \underline{a}} t_{b,c}^a u^b h^c \otimes u^c.$$

*Proof.* Let

$$v_a = u^a - \sum_{b,c \neq 0, \underline{b} + \underline{c} = \underline{a}} t_{b,c}^a \mu_b u^c.$$

Then  $u^a$  can be written as in (2.14) if and only if  $\Delta(v_a) = h^a \otimes v_a + v_a \otimes 1$ . Hence it is enough to prove that

$$\Delta(v_a) - h^a \otimes v_a - v_a \otimes 1 = \Delta(u^a) - h^a \otimes u^a - u^a \otimes 1 - \sum_{b,c \neq 0, \underline{b} + \underline{c} = \underline{a}} t_{b,c}^a u^b h^c \otimes u^c.$$

We compute

$$\begin{aligned}
 \Delta(v_a) - h^a \otimes v_a - v_a \otimes 1 &= \\
 &= \Delta(u^a) - \sum_{b,c \neq 0, \underline{b} + \underline{c} = \underline{a}} t_{b,c}^a \mu_b \Delta(u^c) - h^a \otimes v_a - v_a \otimes 1 \\
 &= \Delta(u^a) - h^a \otimes u^a - u^a \otimes 1 + \sum_{b,c \neq 0, \underline{b} + \underline{c} = \underline{a}} t_{b,c}^a \mu_b (h^a \otimes u^c - h^c \otimes u^c) \\
 &\quad - \sum_{\substack{b,c,f,g \neq 0 \\ \underline{b} + \underline{c} = \underline{a}, \underline{f} + \underline{g} = \underline{c}}} t_{b,c}^a t_{f,g}^c \mu_b u^f h^g \otimes u^g,
 \end{aligned}$$

using the definition of  $v_a$  in the first equation, and the formula for  $\Delta(u^c)$  from (2.13) in the second equation. Note that the term

$$\sum_{b,c \neq 0, \underline{b} + \underline{c} = \underline{a}} t_{b,c}^a \mu_b u^c \otimes 1$$

cancels. Hence we have to show that

$$\begin{aligned}
 &\sum_{\substack{b,c,f,g \neq 0 \\ \underline{b} + \underline{c} = \underline{a}, \underline{f} + \underline{g} = \underline{c}}} t_{b,c}^a t_{f,g}^c \mu_b u^f h^g \otimes u^g = \\
 &= \sum_{b,c \neq 0, \underline{b} + \underline{c} = \underline{a}} t_{b,c}^a (\mu_b h^a \otimes u^c - \mu_b h^c \otimes u^c + u^b h^c \otimes u^c).
 \end{aligned}$$

Since for all  $b, c \neq 0, \underline{b} + \underline{c} = \underline{a}$ , we have  $h^a = h^b h^c$ , it follows that

$$\mu_b h^a \otimes u^c - \mu_b h^c \otimes u^c + u^b h^c \otimes u^c = (\mu_b (h^b - 1) + u^b) h^c \otimes u^c.$$

Using the formula for  $u^b$  from (2.12), we finally have to prove

$$\sum_{\substack{b,c,f,g \neq 0 \\ \underline{b} + \underline{c} = \underline{a}, \underline{f} + \underline{g} = \underline{c}}} t_{b,c}^a t_{f,g}^c \mu_b u^f h^g \otimes u^g = \sum_{\substack{b,c,d,e \neq 0 \\ \underline{b} + \underline{c} = \underline{a}, \underline{d} + \underline{e} = \underline{b}}} t_{b,c}^a t_{d,e}^b \mu_d u^e h^c \otimes u^c.$$

This last equality follows from the coassociativity of  $K(\mathcal{D})$ . Indeed, from

$$(\text{id} \otimes \Delta_{K(\mathcal{D})}) \Delta_{K(\mathcal{D})}(z^a) = (\Delta_{K(\mathcal{D})} \otimes \text{id}) \Delta_{K(\mathcal{D})}(z^a)$$

we obtain with (2.11) after cancelling several terms

$$\sum_{\substack{b,c,f,g \neq 0 \\ \underline{b} + \underline{c} = \underline{a}, \underline{f} + \underline{g} = \underline{c}}} t_{b,c}^a t_{f,g}^c z^b \otimes z^f \otimes z^g = \sum_{\substack{b,c,d,e \neq 0 \\ \underline{b} + \underline{c} = \underline{a}, \underline{d} + \underline{e} = \underline{b}}} t_{b,c}^a t_{d,e}^b z^d \otimes z^e \otimes z^c.$$

Thus mapping  $z^r \otimes z^s \otimes z^t, r, s, t \neq 0, \text{ht}(\underline{r}), \text{ht}(\underline{s}), \text{ht}(\underline{t}) < n$ , onto  $\mu_r u^s h^t \otimes u^t$  proves the claim. Here we are using that the elements  $z^a$  are linearly independent by Theorem 2.5.  $\square$

Let  $K(\mathcal{D})\#k[\Gamma]$  be the Hopf algebra corresponding to the braided Hopf algebra  $K(\mathcal{D})$  by (1.3). Thus by definition and Lemma 2.7, for all  $0 \neq a \in \mathbb{N}^p$ ,

$$(2.16) \quad \Delta_{K(\mathcal{D})\#k[\Gamma]}(z^a) = h^a \otimes z^a + z^a \otimes 1 + \sum_{b,c \neq 0, b+c=a} t_{b,c}^a z^b h^c \otimes z^c.$$

For all  $n \geq 0$ , let  $K(\mathcal{D})_n$  be the vector subspace spanned by all  $z^a, a \in \mathbb{N}^p, \text{ht}(\underline{a}) \leq n$ . Then  $K(\mathcal{D})_n\#k[\Gamma] \subset K(\mathcal{D})\#k[\Gamma]$  is a subcoalgebra.

In the next Lemma we describe all coalgebra maps

$$\varphi : K(\mathcal{D})_n\#k[\Gamma] \rightarrow k[\Gamma] \text{ with } \varphi|_{\Gamma} = \text{id}.$$

Note that such a coalgebra map is given by a family of elements  $\varphi(z^a) =: u^a, 0 \neq a \in \mathbb{N}^p, \text{ht}(\underline{a}) \leq n$ , such that (2.15) holds for all  $0 \neq a, \text{ht}(\underline{a}) \leq n$ . It follows by induction on  $\text{ht}(\underline{a})$  from Lemma 2.8 with (2.14) that  $\varepsilon(u^a) = 0$  for all  $a$ .

**Lemma 2.9.** *Let  $n \geq 1$ .*

(1) *Let  $(\mu_a)_{0 \neq a \in \mathbb{N}^p, \text{ht}(\underline{a}) \leq n}$  be a family of elements in  $k$  such that for all  $a$ , if  $h^a = 1$ , then  $\mu_a = 0$ . Define the family  $(u^a)_{0 \neq a \in \mathbb{N}^p, \text{ht}(\underline{a}) \leq n}$  by induction on  $\text{ht}(\underline{a})$  by (2.14). Then*

$$\varphi : K(\mathcal{D})_n\#k[\Gamma] \rightarrow k[\Gamma], \varphi(z^a g) = u^a g, a \in \mathbb{N}^p, \text{ht}(\underline{a}) \leq n, g \in \Gamma,$$

*is a coalgebra map.*

(2) *The map defined in (1) from the set of all  $(\mu_a)_{0 \neq a \in \mathbb{N}^p, \text{ht}(\underline{a}) \leq n}$  such that for all  $a$ , if  $h^a = 1$ , then  $\mu_a = 0$ , to the set of all coalgebra maps  $\varphi$  with  $\varphi|_{\Gamma} = \text{id}$  is bijective.*

*Proof.* This follows from Lemma 2.8 by induction on  $\text{ht}(\underline{a})$ . Note that the coefficient  $\mu_a$  in (2.14) is uniquely determined if we define  $\mu_a = 0$  if  $h^a = 1$ .  $\square$

**Definition 2.10.** Let  $n \geq 1$ . A coalgebra map  $\varphi : K(\mathcal{D})_n\#k[\Gamma] \rightarrow k[\Gamma]$  with  $\varphi|_{\Gamma} = \text{id}$  is called a *partial Hopf algebra map*, if for all  $x, y \in K(\mathcal{D})_n\#k[\Gamma]$  with  $xy \in K(\mathcal{D})_n\#k[\Gamma]$ , we have  $\varphi(xy) = \varphi(x)\varphi(y)$ .

**Lemma 2.11.** *Let  $n \geq 1$ , and  $\varphi : K(\mathcal{D})_n\#k[\Gamma] \rightarrow k[\Gamma]$  a coalgebra map,  $(\mu_a)_{0 \neq a \in \mathbb{N}^p, \text{ht}(\underline{a}) \leq n}$  the family of scalars corresponding to  $\varphi$  by Lemma 2.9, and  $u^a = \varphi(a)$  for all  $a \in \mathbb{N}^p$  with  $\text{ht}(\underline{a}) \leq n$ . Then the following are equivalent:*

- (1)  $\varphi$  is a partial Hopf algebra map.
- (2) For all  $0 \neq a = (a_1, \dots, a_p) \in \mathbb{N}^p$  with  $\text{ht}(\underline{a}) \leq n$ ,
  - (a)  $u^a = \prod_{a_l > 0} u_l^{a_l}$ , where for all  $1 \leq l \leq p, u_l = u^{e_l}$ , if  $a_l > 0$ ,
  - (b) if  $\eta^a \neq \varepsilon$ , then  $\mu_a = 0$ , and  $u^a = 0$ .



- (3) (a) As (2) (a).  
 (b) For all  $1 \leq l \leq p$  with  $\text{ht}(\underline{e}_l) \leq n$ , if  $\eta_l \neq \varepsilon$ , then  $u^{e_l} = 0$ .

*Proof.* (1)  $\Rightarrow$  (2): If  $\varphi$  is a partial Hopf algebra map, then (a) follows immediately, and to prove (b), let  $0 \neq a \in \mathbb{N}^p$ ,  $\text{ht}(\underline{a}) \leq n$ , and  $g \in \Gamma$ , with  $\eta^a \neq \varepsilon$ . Then

$$\varphi(gz^a) = \eta^a(g)u^a g = u^a g,$$

since  $gz^a = \eta^a(g)z^a g$  by (2.10). Thus  $u^a = 0$ , and it follows by induction on  $\text{ht}(\underline{a})$  from (2.14) that  $\mu_a = 0$ , since for all  $0 \neq b, c \in \mathbb{N}^p$  with  $\text{ht}(\underline{b}) + \text{ht}(\underline{c}) = \text{ht}(\underline{a})$ ,  $\eta^b \neq \varepsilon$ , or  $\eta^c \neq \varepsilon$ .

(2)  $\Rightarrow$  (3) is trivial. (3)  $\Rightarrow$  (1): The coalgebra map  $\varphi$  is a partial Hopf algebra map if and only if for all  $b, c \in \mathbb{N}^p$  with  $\text{ht}(\underline{b}) + \text{ht}(\underline{c}) \leq n$ , and  $g, h \in \Gamma$ ,

$$\varphi(z^b g z^c h) = u^b g u^c h.$$

By (2.9) and (2.10),  $z^b g z^c h = \eta^c(g)\gamma_{b,c}z^{b+c}gh$ . Thus (1) is equivalent to (2.17)  $\eta^c(g)\gamma_{b,c}u^{b+c} = u^b u^c$  for all  $b, c \in \mathbb{N}^p$ ,  $\text{ht}(\underline{b}) + \text{ht}(\underline{c}) \leq n$ ,  $g \in \Gamma$ .

Let  $b, c \in \mathbb{N}^p$ ,  $\text{ht}(\underline{b}) + \text{ht}(\underline{c}) \leq n$ ,  $g \in \Gamma$ . By (a),

$$u^{b+c} = u^b u^c = \prod_{b_l + c_l > 0} u_l^{b_l + c_l}.$$

To prove (2.17) assume that  $u^b u^c \neq 0$ . Then  $u_l \neq 0$  for all  $l$  with  $c_l > 0$ . Hence by (b),  $\eta_l = \varepsilon$  for all  $l$  with  $c_l > 0$ , and  $\eta^c(g) = 1$ ,  $\gamma_{b,c} = 1$ .  $\square$

To formulate the main result of this section, we define  $M(\mathcal{D})$  as the set of all families  $(\mu_l)_{1 \leq l \leq p}$  of elements in  $k$  satisfying the following condition for all  $1 \leq l \leq p$ : If  $h_l = 1$  or  $\eta_l \neq \varepsilon$ , then  $\mu_l = 0$ .

**Theorem 2.12.** (1) Let  $\mu = (\mu_l)_{1 \leq l \leq p} \in M(\mathcal{D})$ . Then there is exactly one Hopf algebra map

$$\varphi_\mu : K(\mathcal{D}) \# k[\Gamma] \rightarrow k[\Gamma], \quad \varphi|_\Gamma = \text{id}$$

such that the family  $(\mu_a)_{0 \neq a \in \mathbb{N}^p}$  associated to  $\varphi_\mu$  by Lemma 2.9 satisfies  $\mu_{e_l} = \mu_l$  for all  $1 \leq l \leq p$ .

(2) The map  $\mu \mapsto \varphi_\mu$  defined in (1) from  $M(\mathcal{D})$  to the set of all Hopf algebra homomorphisms  $\varphi : K(\mathcal{D}) \# k[\Gamma] \rightarrow k[\Gamma]$  with  $\varphi|_\Gamma = \text{id}$  is bijective.

*Proof.* (1) We proceed by induction on  $n$  to construct partial Hopf algebra maps on  $K(\mathcal{D})_n \# k[\Gamma]$ , the case  $n = 0$  being trivial. We assume that we are given a partial Hopf algebra map

$$\varphi : K(\mathcal{D})_{n-1} \# k[\Gamma] \rightarrow k[\Gamma], \quad n \geq 1,$$

such that  $\mu_{e_l} = \mu_l$  for all  $1 \leq l \leq p$  with  $\text{ht}(e_l) \leq n - 1$ . Here  $(\mu_a)_{0 \neq a \in \mathbb{N}^p, \text{ht}(\underline{a}) \leq n-1}$  is the family of scalars associated to  $\varphi$  by Lemma 2.9. We define  $u^b = \varphi(z^b)$  for all  $0 \neq b, \text{ht}(\underline{b}) \leq n - 1$ . It is enough to show that there is exactly one partial Hopf algebra map

$$\psi : K(\mathcal{D})_n \# k[\Gamma] \rightarrow k[\Gamma]$$

extending  $\varphi$ , and such that  $\mu_{e_l} = \mu_l$  for all  $l$  with  $\text{ht}(e_l) \leq n$ .

Let  $a \in \mathbb{N}^p$  with  $\text{ht}(\underline{a}) = n$ . To define  $\psi(z^a) =: u^a$  we distinguish two cases.

If  $a = e_l$  for some  $1 \leq l \leq p$ , we define

$$(2.18) \quad u^a = \mu_l(1 - h^a) + \sum_{b, c \neq 0, \underline{b} + \underline{c} = \underline{a}} t_{b,c}^a \mu_b u^c.$$

Then (2.15) holds by Lemma 2.8.

If  $a = (a_1, \dots, a_l, 0, \dots, 0)$ ,  $a_l \geq 1$ ,  $1 \leq l \leq p$ , and  $a \neq e_l$ , then  $a = r + s$ , where  $0 \neq r, s = e_l$ . We define  $u^a = u^r u^s$ . To see that  $u^a$  satisfies (2.15), using (2.16) we write

$$\Delta(z^c) = h^c \otimes z^c + z^c \otimes 1 + T(c), \text{ for all } 0 \neq c \in \mathbb{N}^p.$$

Since  $z^r z^s = z^a$  because of (2.9) (note that  $\gamma_{r,s} = 1$  in this case) we see that  $\Delta(z^r)\Delta(z^s) = h^a \otimes z^a + z^a \otimes 1 + T(r, s)$ , where

$$T(r, s) = h^r z^s \otimes z^r + z^r h^s \otimes z^s + (h^r \otimes z^r + z^r \otimes 1)T(s) + T(r)(h^s \otimes z^s + z^s \otimes 1),$$

and  $T(r, s) = T(a)$ . Since  $\varphi$  on  $K(\mathcal{D})_{n-1} \# k[\Gamma]$  is a coalgebra map,

$$\Delta(u^c) = h^c \otimes u^c + u^c \otimes 1 + (\varphi \otimes \varphi)(T(c)),$$

for all  $0 \neq c \in \mathbb{N}^p$  with  $\text{ht}(\underline{c}) \leq n - 1$ . In particular,

$$\Delta(u^r)\Delta(u^s) = h^a \otimes u^a + u^a \otimes 1 + (\varphi \otimes \varphi)(T(r, s)).$$

Thus  $\Delta(u^a) = h^a \otimes u^a + u^a \otimes 1 + (\varphi \otimes \varphi)(T(a))$ , that is,  $u^a$  satisfies (2.15).

Thus the extension of  $\varphi$  defined by  $\psi(z^a g) = u^a g$  for all  $g \in \Gamma, a \in \mathbb{N}^p, \text{ht}(\underline{a}) = n$  is a coalgebra map.

To prove that the extension  $\psi$  is a partial Hopf algebra map, we check condition (3) in Lemma 2.11. Since the restriction of  $\psi$  to  $K(\mathcal{D})_{n-1} \# k[\Gamma]$  is a partial Hopf algebra map, (3) (a) is satisfied. To prove (3)(b), let  $1 \leq l \leq p$  with  $\text{ht}(\underline{e}_l) = n$ ,  $a = e_l$ , and assume  $\eta_l \neq \varepsilon$ . Then for all  $0 \neq b, c \in \mathbb{N}^p$  with  $\underline{b} + \underline{c} = \underline{a}$ , we have  $\eta^b \neq \varepsilon$  or  $\eta^c \neq \varepsilon$ . Since  $\varphi$  is a Hopf algebra map, it follows from Lemma 2.11 that  $\mu_b = 0$  or  $u^c = 0$ . By assumption,  $\mu_l = 0$ . Hence by (2.18),  $u^a = 0$ .

This proves (1) since the uniqueness of the extension follows from Lemma 2.8 and Lemma 2.9.

(2) By Lemma 2.9, the map  $\mu \mapsto \varphi_\mu$  is injective. To prove surjectivity, let  $\varphi : K(\mathcal{D})\#k[\Gamma] \rightarrow k[\Gamma]$  be a Hopf algebra map with  $\varphi|_\Gamma = \text{id}$ . By Lemma 2.9,  $\varphi$  is defined by a family  $(\mu_a)_{0 \neq a \in \mathbb{N}^p}$  of scalars. By (1),  $\varphi$  is determined by the values  $\mu_{e_l}, 1 \leq l \leq p$ .  $\square$

**Definition 2.13.** For any  $\mu \in M(\mathcal{D})$  and  $1 \leq l \leq p$ , let  $\varphi_\mu$  be the Hopf algebra map defined in Theorem 2.12, and

$$u_l(\mu) = \varphi_\mu(z_l) \in k[\Gamma].$$

If  $\alpha$  is a positive root in  $\Phi^+$  with  $\alpha = \beta_l$ , we define  $u_\alpha(\mu) = u_l(\mu)$ .

Note that by (2.14), each  $u_\alpha(\mu)$  lies in the augmentation ideal of  $k[g_i^N \mid 1 \leq i \leq \theta]$ .

### 3. LINKING

**3.1. Notations.** In this Section we fix a finite abelian group  $\Gamma$ , and a datum  $\mathcal{D} = \mathcal{D}(\Gamma, (g_i)_{1 \leq i \leq \theta}, (\chi_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta})$  of finite Cartan type. We follow the notations of the previous Section, in particular,  $q_{ij} = \chi_j(g_i)$  for all  $i, j$ .

For all  $1 \leq i, j \leq \theta$  we write  $i \sim j$  if  $i$  and  $j$  are in the same connected component of the Dynkin diagram of  $(a_{ij})$ . Let  $\mathcal{X} = \{I_1, \dots, I_t\}$  be the set of connected components of  $I = \{1, 2, \dots, \theta\}$ . We assume

(3.1)  $q_{ij}$  has odd order for all  $i, j$ , and

(3.2) the order of  $q_{ii}$  is prime to 3, if  $i$  lies in a component  $G_2$ .

For all  $J \in \mathcal{X}$ , let  $N_J$  be the common order of  $q_{ii}, i \in J$ .

As in Section 2.2, for all  $J \in \mathcal{X}$ , we choose a reduced decomposition of the longest element  $w_{0,J}$  of the Weyl group  $W_J$  of the root system  $\Phi_J$  of  $(a_{ij})_{i,j \in J}$ . Then for all  $J, K \in \mathcal{X}$ ,  $w_{0,J}$  and  $w_{0,K}$  commute in the Weyl group  $W$  of the root system  $\Phi$  of  $(a_{ij})_{1 \leq i, j \leq \theta}$ , and

$$w_0 = w_{0,I_1} w_{0,I_2} \cdots w_{0,I_t}$$

gives a reduced representation of the longest element of  $W$ . For all  $J \in \mathcal{X}$ , let  $p_J$  be the number of positive roots in  $\Phi_J^+$ , and

$$\Phi_J^+ = \{\beta_{J,1}, \dots, \beta_{J,p_J}\}$$

the corresponding convex ordering. Then

$$\Phi^+ = \{\beta_{I_1,1}, \dots, \beta_{I_1,p_{I_1}}, \dots, \beta_{I_t,1}, \dots, \beta_{I_t,p_{I_t}}\}$$

is the convex ordering corresponding to the reduced representation of  $w_0 = w_{0,I_1} w_{0,I_2} \cdots w_{0,I_t}$ . We also write

$$\Phi^+ = \{\beta_1, \dots, \beta_p\}, \quad p = \sum_{J \in \mathcal{X}} p_J,$$

for this ordering.

In Section 2.1 we have defined root vectors  $x_\alpha$  in the free algebra  $k\langle x_1, \dots, x_\theta \rangle$  for each positive root in  $\Phi_J^+ \subset \Phi$ ,  $J \in \mathcal{X}$ .

We recall a notion from [AS4].

**Definition 3.1.** A family  $\lambda = (\lambda_{ij})_{1 \leq i < j \leq \theta, i \not\sim j}$  of elements in  $k$  is called a *family of linking parameters for  $\mathcal{D}$*  if the following condition is satisfied for all  $1 \leq i < j \leq \theta$ ,  $i \not\sim j$ : If  $g_i g_j = 1$  or  $\chi_i \chi_j \neq \varepsilon$ , then  $\lambda_{ij} = 0$ . Vertices  $1 \leq i, j \leq \theta$  are called *linkable* if  $i \not\sim j$ ,  $g_i g_j \neq 1$  and  $\chi_i \chi_j = \varepsilon$ .

Any vertex  $i$  is linkable to at most one vertex  $j$ , and if  $i, j$  are linkable, then  $q_{ii} = q_{jj}^{-1}$  [AS4, Section 5.1].

The free algebra  $k\langle x_1, \dots, x_\theta \rangle$  is a braided Hopf algebra in  ${}^\Gamma \mathcal{YD}$  as explained in Section 1.1. Then  $k\langle x_1, \dots, x_\theta \rangle \# k[\Gamma]$  is a Hopf algebra as in 1.2. For simplicity we write  $xg$  instead of  $x \# g$  for elements  $x \in k\langle x_1, \dots, x_\theta \rangle$  and  $g \in \Gamma$ .

**3.2. The Hopf algebra  $U(\mathcal{D}, \lambda)$ .** We assume the situation of Section 3.1.

**Definition 3.2.** Let  $\lambda = (\lambda_{ij})_{1 \leq i < j \leq \theta, i \not\sim j}$  be a family of linking parameters for  $\mathcal{D}$ . Let  $U(\mathcal{D}, \lambda)$  be the quotient Hopf algebra of  $k\langle x_1, \dots, x_\theta \rangle \# k[\Gamma]$  modulo the ideal generated by

$$(3.3) \quad \text{ad}_c(x_i)^{1-a_{ij}}(x_j), \text{ for all } 1 \leq i, j \leq \theta, i \sim j, i \neq j,$$

$$(3.4) \quad x_i x_j - q_{ij} x_j x_i - \lambda_{ij}(1 - g_i g_j), \text{ for all } 1 \leq i < j \leq \theta, i \not\sim j.$$

We denote the images of  $x_i$  and  $g \in \Gamma$  in  $U(\mathcal{D}, \lambda)$  again by  $x_i$  and  $g$ . The elements in (3.3) and (3.4) are skew-primitive. Hence  $U(\mathcal{D}, \lambda)$  is a Hopf algebra with

$$\Delta(x_i) = g_i \otimes x_i + x_i \otimes 1, \quad 1 \leq i \leq \theta.$$

**Theorem 3.3.** *Let  $\Gamma$  be a finite abelian group, and  $\mathcal{D}$  a datum of finite Cartan type satisfying (3.1) and (3.2). Let  $\lambda$  be a family of linking parameters for  $\mathcal{D}$ . Then*

(1) *The elements*

$$x_{\beta_1}^{a_1} x_{\beta_2}^{a_2} \cdots x_{\beta_p}^{a_p} g, \quad a_1, a_2, \dots, a_p \geq 0, g \in \Gamma,$$

*form a basis of the vector space  $U(\mathcal{D}, \lambda)$ .*

(2) *Let  $J \in \mathcal{X}$ , and  $\alpha \in \Phi^+$ ,  $\beta \in \Phi_J^+$ . Then  $[x_\alpha, x_\beta^{N_J}]_c = 0$ , that is,*

$$x_\alpha x_\beta^{N_J} = q_{\alpha, \beta}^{N_J} x_\beta^{N_J} x_\alpha.$$

*Proof.* We adapt the method of proof of [AS4, Section 5.3] and proceed by induction on the number  $t$  of connected components.

If  $I$  is connected, (1) and (2) follow from Theorem 2.5.

If  $t > 1$ , we assume that  $I_1 = \{1, 2, \dots, \tilde{\theta}\}$ ,  $1 \leq \tilde{\theta} < \theta$ . For all  $1 \leq i \leq \tilde{\theta}$ , let  $l_i$  be the least common multiple of the orders of  $g_i$  and  $\chi_i$ ,  $1 \leq i \leq \tilde{\theta}$ . Let  $\tilde{\Gamma} = \langle h_1, \dots, h_{\tilde{\theta}} \mid h_i h_j = h_j h_i, h_i^{l_i} = 1 \text{ for all } i, j \rangle$ , and define for all  $1 \leq i \leq \tilde{\theta}$  the character  $\eta_j$  of  $\tilde{\Gamma}$  by  $\eta_j(h_i) = \chi_j(g_i)$ ,  $1 \leq i, j \leq \tilde{\theta}$ . Then we define

$$\mathcal{D}_1 = \mathcal{D}(\tilde{\Gamma}, (h_i)_{1 \leq i \leq \tilde{\theta}}, (\eta_i)_{1 \leq i \leq \tilde{\theta}}, (a_{ij})_{1 \leq i, j \leq \tilde{\theta}}).$$

Let  $\mathcal{D}_2 = \mathcal{D}(\Gamma, (g_i)_{\tilde{\theta} < i \leq \theta}, (\chi_i)_{\tilde{\theta} < i \leq \theta}, (a_{ij})_{\tilde{\theta} < i, j \leq \theta})$  be the restriction of  $\mathcal{D}$  to  $I_2 \cup \dots \cup I_t$ , and  $\lambda_2 = (\lambda_{ij})_{\tilde{\theta} < i < j \leq \theta, i \neq j}$ . We define  $U = U(\mathcal{D}_1)$  (with empty family of linking parameters) with generators  $x_1, \dots, x_{\tilde{\theta}}$ , and  $h \in \tilde{\Gamma}$ , and  $A = U(\mathcal{D}_2, \lambda_2)$  with generators  $y_{\tilde{\theta}+1}, \dots, y_{\theta}$ , and  $g \in \Gamma$ .

It is shown in [AS4, Lemma 5.19] that there are algebra maps  $\gamma_i$ ,  $(\varepsilon, \gamma)$ -derivations  $\delta_i$  and a Hopf algebra map  $\varphi$ ,

$$\gamma_i : A \rightarrow k, \delta_i : A \rightarrow k, \varphi : U \rightarrow (A^0)^{\text{cop}}, 1 \leq i \leq \tilde{\theta},$$

such that for all  $1 \leq i \leq \tilde{\theta} < j \leq \theta$ ,

$$\begin{aligned} \gamma_i|_{\Gamma} &= \chi_i, \gamma_i(y_j) = 0, \\ \delta_i|_{\Gamma} &= 0, \delta_i(y_j) = -\chi_i(g_j)\lambda_{ij}, \\ \varphi(h_i) &= \gamma_i, \varphi(x_i) = \delta_i. \end{aligned}$$

Then  $\sigma : U \otimes A \otimes U \otimes A \rightarrow U \otimes A$ , defined for all  $u, v \in U, a, b \in A$  by

$$\sigma(u \otimes a, v \otimes b) = \varepsilon(u)\tau(v, a)\varepsilon(b), \tau(v, a) = \varphi(v)(a),$$

is a 2-cocycle on the tensor product Hopf algebra of  $U$  and  $A$ , and  $(U \otimes A)_{\sigma}$  is the Hopf algebra with twisted multiplication defined in (1.9). Multiplication in  $(U \otimes A)_{\sigma}$  is given for all  $u, v \in U, a, b \in A$  by

$$(3.5) \quad (u \otimes a) \cdot_{\sigma} (v \otimes b) = u\tau(v_{(1)}, a_{(1)})v_{(2)} \otimes a_{(2)}\tau^{-1}(v_{(3)}, a_{(3)})b,$$

with  $\tau^{-1}(u, a) = \varphi(u)(S^{-1}(a))$ .

The group-like elements  $h_i \otimes g_i^{-1}$ ,  $1 \leq i \leq \tilde{\theta}$ , are central in  $(U \otimes A)_{\sigma}$ , and as in the last part of the proof of [AS4, Theorem 5.17] it can be seen that the map

$$(U \otimes A)_{\sigma} \rightarrow U(\mathcal{D}, \lambda), x_i \otimes 1 \mapsto x_i, h_i \otimes 1 \mapsto g_i, 1 \otimes y_j \mapsto x_j, 1 \otimes g \mapsto g$$

for all  $1 \leq i \leq \tilde{\theta} < j \leq \theta$ ,  $g \in \Gamma$ , induces an isomorphism of Hopf algebras

$$(3.6) \quad (U \otimes A)_{\sigma} / (h_i \otimes g_i^{-1} - 1 \otimes 1 \mid 1 \leq i \leq \tilde{\theta}) \cong U(\mathcal{D}, \lambda).$$

By induction and Theorem 2.5, the elements

$$x_{\beta_1}^{a_1} \cdots x_{\beta_{p_1}}^{a_{p_1}} h \otimes y_{\beta_{p_1+1}}^{a_{p_1+1}} \cdots y_{\beta_p}^{a_p} g, \quad a_1, \dots, a_p \geq 0, h \in \tilde{\Gamma}, g \in \Gamma,$$

are a basis of  $U \otimes A$ . It follows from (3.5) that for all  $p_1 < l \leq p$  and  $1 \leq i \leq \tilde{\theta}$ ,

$$(1 \otimes y_{\beta_l}) \cdot_{\sigma} (h_i \otimes 1) = \chi_i(g_{\beta_l}) h_i \otimes y_{\beta_l}.$$

Hence

$$(x_{\beta_1}^{a_1} \cdots x_{\beta_{p_1}}^{a_{p_1}} \otimes y_{\beta_{p_1+1}}^{a_{p_1+1}} \cdots y_{\beta_p}^{a_p}) \cdot_{\sigma} (h \otimes g), \quad a_1, \dots, a_p \geq 0, h \in \tilde{\Gamma}, g \in \Gamma,$$

is a basis of  $(U \otimes A)_{\sigma}$ .

Let  $P = \{h \otimes g \in (U \otimes A)_{\sigma} \mid h \in \tilde{\Gamma}, g \in \Gamma\}$ , and let  $\tilde{P} \subset P$  be the subgroup generated by  $h_i \otimes g_i^{-1}$ ,  $1 \leq i \leq \tilde{\theta}$ . Then

$$\Gamma \rightarrow P/\tilde{P}, \quad g \mapsto \overline{1 \otimes g},$$

is a group isomorphism. By (3.6),  $(U \otimes A)_{\sigma} \otimes_{k[P]} k[P/\tilde{P}] \cong U(\mathcal{D})$ . Hence

$$x_{\beta_1}^{a_1} x_{\beta_2}^{a_2} \cdots x_{\beta_p}^{a_p} g, \quad a_1, a_2, \dots, a_p \geq 0, g \in \Gamma,$$

is a basis of  $U(\mathcal{D}, \lambda)$ .

To prove (2), we first show that for all  $\tilde{\theta} < i \leq \theta$ , and  $\beta \in \Phi_{I_1}^+$ , with  $N = N_{I_1}$

$$(3.7) \quad (1 \otimes y_i) \cdot_{\sigma} (x_{\beta}^N \otimes 1) = \chi_{\beta}^N(g_i)(x_{\beta}^N \otimes 1) \cdot_{\sigma} (1 \otimes y_i)$$

in  $(U \otimes A)_{\sigma}$ . We use the notations of Section 2.2 with  $N = N_{I_1}$ ,  $z_{\beta} = x_{\beta}^N$ . By (2.16)

$$\Delta_U(z_{\beta}) = g_{\beta}^N \otimes z_{\beta} + z_{\beta} \otimes 1 + \sum_{b,c \neq 0, \underline{b} + \underline{c} = \beta} t_{b,c}^a z^b h^c \otimes z^c.$$

Since  $\Delta(y_i) = g_i \otimes y_i + y_i \otimes 1$ , and

$$\Delta^2(y_i) = g_i \otimes g_i \otimes y_i + g_i \otimes y_i \otimes 1 + y_i \otimes 1 \otimes 1,$$

we have for all  $u \in U$  by (3.5)

$$\begin{aligned} (1 \otimes y_i) \cdot_{\sigma} (u \otimes 1) &= \varphi(u_{(1)})(g_i) u_{(2)} \otimes g_i \varphi(u_{(3)})(S^{-1}(y_i)) \\ &\quad + \varphi(u_{(1)})(g_i) u_{(2)} \otimes y_i \varphi(u_{(3)})(1) \\ &\quad + \varphi(u_{(1)})(y_i) u_{(2)} \otimes 1 \varphi(u_{(3)})(1). \end{aligned}$$

It follows from the definition of  $\varphi$  that

$$\varphi(x_{\beta_l})(g) = 0 \text{ for all } \beta_l \in \Phi_1^+, g \in \Gamma.$$

Hence to compute  $(1 \otimes y_i) \cdot_\sigma (u \otimes 1)$  with  $u = z_\beta$ , we only need to take into account the term  $g_\beta^N \otimes z_\beta \otimes 1$  of  $\Delta^2(z_\beta)$ , and we obtain

$$\begin{aligned} (1 \otimes y_i) \cdot_\sigma (u \otimes 1) &= \varphi(g_\beta^N)(y_{i(1)})z_\beta \otimes y_{i(2)}\varphi(1)(S^{-1}(y_{i(3)})) \\ &= \varphi(g_\beta^N)(y_{i(1)})z_\beta \otimes y_{i(2)} \\ &= \varphi(g_\beta^N)(g_i)z_\beta \otimes y_i + \varphi(g_\beta^N)(y_i)z_\beta \otimes 1 \\ &= \chi_\beta^N(g_i)(x_\beta^N \otimes 1) \cdot_\sigma (1 \otimes y_i), \end{aligned}$$

since  $\varphi(g_\beta^N) = \chi_\beta^N$  and  $\varphi(g_\beta^N)(y_i) = 0$  by the definition of  $\varphi$ .

From (3.6) and (3.7) we see that for all simple roots  $\alpha \in \Phi_K^+$ ,  $K \in \mathcal{X}$ ,  $K \neq I_1$  and all roots  $\beta \in \Phi_J^+$  with  $J = I_1$

$$(3.8) \quad x_\alpha x_\beta^{N_J} = \chi_\beta^{N_J}(g_\alpha) x_\beta^{N_J} x_\alpha$$

in  $U(\mathcal{D}, \lambda)$ . Since the root vectors  $x_\alpha$  are homogeneous, (3.8) holds for all  $\alpha \in \Phi_K^+$ ,  $K \neq I_1$ , and  $\beta \in \Phi_{I_1}^+$ . Since  $U(\mathcal{D}, \lambda)$  and the root vectors  $x_\alpha$ ,  $\alpha \in \Phi^+$ , do not depend on the order of the connected components, we can reorder the connected components and obtain (3.8) for all positive roots  $\alpha, \beta$  lying in different connected components. For roots in the same connected component, (3.8) follows from Theorem 2.5.  $\square$

#### 4. FINITE-DIMENSIONAL QUOTIENTS

**4.1. A general criterion.** We need a generalization of Theorem [AS5, 6.24].

In this section, let  $\Gamma$  be an abelian group,  $A$  an algebra containing the group algebra  $k[\Gamma]$  as a subalgebra and  $p \geq 1$ . We assume

$$y_1, \dots, y_p \in A, h_1, \dots, h_p \in \Gamma, \psi_1, \dots, \psi_p \in \widehat{\Gamma}, \text{ and } N_1, \dots, N_p \geq 1,$$

such that

$$(4.1) \quad gy_l = \psi_l(g)y_lg, \text{ for all } 1 \leq l \leq p, g \in \Gamma,$$

$$(4.2) \quad y_k y_l^{N_l} = \psi_l^{N_l}(h_k) y_l^{N_l} y_k, \text{ for all } 1 \leq k, l \leq p,$$

$$(4.3) \quad y_1^{a_1} \cdots y_p^{a_p} g, \ a_1, \dots, a_p \geq 0, g \in \Gamma, \text{ form a basis of } A.$$

For all  $a = (a_1, \dots, a_p) \in \mathbb{N}^p$ , we define  $y^a = y_1^{a_1} \cdots y_p^{a_p}$  and

$$\mathbb{L} = \{l = (l_1, \dots, l_p) \in \mathbb{N}^p \mid 0 \leq l_i < N_i \text{ for all } 1 \leq i \leq p\}.$$

Hence any element of  $y \in A$  can be written as

$$y = \sum_{l \in \mathbb{L}, a \in \mathbb{N}^p} y^l y^{aN} w_{l,a}, \ w_{l,a} \in k[\Gamma] \text{ for all } l \in \mathbb{L}, a \in \mathbb{N}^p,$$

where the coefficients  $w_{l,a} \in k[\Gamma]$  are uniquely determined. In [AS5] we assumed that  $A = R \# k[\Gamma]$ , and the subalgebra  $R$  of  $A$  generated by

$y_1, \dots, y_p$  had the basis  $y_1^{a_1} \cdots y_p^{a_p}$ ,  $a_1, \dots, a_p \geq 0$ . Hence for  $y \in R$  we could assume that the  $w_{l,a}$  were scalars.

**Theorem 4.1.** *Assume the situation above, and let  $u_l \in k[\Gamma]$ ,  $1 \leq l \leq p$ . Then the following are equivalent:*

- (1) *The residue classes of  $y^a g$ ,  $a \in \mathbb{L}$ ,  $g \in \Gamma$ , form a basis of the quotient algebra  $A/(y_l^{N_l} - u_l \mid 1 \leq l \leq p)$ .*
- (2) *For all  $1 \leq l \leq p$ ,  $u_l$  is central in  $A$ , and if  $\psi_l^{N_l} \neq \varepsilon$ , then  $u_l = 0$ .*

*Proof.* As in [AS5] this follows from Lemma [AS5, 6.23]. To extend the proof of this Lemma to the more general case considered here, we use the following rule. Assume (2), and let  $u^a = u_1^{a_1} \cdots u_p^{a_p}$ , for all  $a = (a_1, \dots, a_p) \in \mathbb{N}^p$ . For all  $1 \leq l \leq p$ , let  $\tilde{\psi}_l : k[\Gamma] \rightarrow k[\Gamma]$  be the algebra isomorphism with  $\tilde{\psi}_l(g) = \psi_l(g)g$  for all  $g \in \Gamma$ . Then

$$(4.4) \quad u^a \tilde{\psi}^{aN}(w) = u^a w, \text{ for all } w \in k[\Gamma], a \in \mathbb{N}^p,$$

where  $\tilde{\psi}^{aN} = \tilde{\psi}_1^{a_1 N_1} \cdots \tilde{\psi}_p^{a_p N_p}$ .  $\square$

**4.2. The Hopf algebra  $u(\mathcal{D}, \lambda, \mu)$ .** Let  $\Gamma$  be a finite abelian group, and  $\mathcal{D} = \mathcal{D}(\Gamma, (g_i)_{1 \leq i \leq \theta}, (\chi_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta})$  a datum of finite Cartan type. We assume the situation of Section 3.1.

**Definition 4.2.** A family  $\mu = (\mu_\alpha)_{\alpha \in \Phi^+}$  of elements in  $k$  is called a *family of root vector parameters for  $\mathcal{D}$*  if the following condition is satisfied for all  $\alpha \in \Phi_J^+$ ,  $J \in \mathcal{X}$ : If  $g_\alpha^{N_J} = 1$  or  $\chi_\alpha^{N_J} \neq \varepsilon$ , then  $\mu_\alpha = 0$ .

Let  $\mu$  be a family of root vector parameters for  $\mathcal{D}$ . For all  $J \in \mathcal{X}$ , and  $\alpha \in \Phi_J^+$ , we define

$$(4.5) \quad \pi_J(\mu) = (\mu_\beta)_{\beta \in \Phi_J^+}, \text{ and } u_\alpha(\mu) = u_\alpha(\pi_J(\mu)),$$

where  $u_\alpha(\pi_J(\mu))$  is introduced in Definition 2.13. Let  $\lambda$  be a family of linking parameters for  $\mathcal{D}$ . Then we define

$$(4.6) \quad u(\mathcal{D}, \lambda, \mu) = U(\mathcal{D}, \lambda) / (x_\alpha^{N_J} - u_\alpha(\mu) \mid \alpha \in \Phi_J^+, J \in \mathcal{X}).$$

By abuse of language we still write  $x_i$  and  $g$  for the images of  $x_i$  and  $g \in \Gamma$  in  $u(\mathcal{D}, \lambda, \mu)$ . For all  $1 \leq l \leq p$ , we define  $N_l = N_J$ , if  $\beta_l \in \Phi_J^+$ ,  $J \in \mathcal{X}$ .

**Lemma 4.3.** *Let  $\mathcal{D}, \lambda$  and  $\mu$  as above, and  $\alpha \in \Phi^+$ . Then  $u_\alpha(\mu)$  is central in  $U(\mathcal{D}, \Lambda)$ .*

*Proof.* Let  $\alpha \in \Phi_J^+$ , where  $J \in \mathcal{X}$ , and  $N = N_J$ . To simplify the notation, we assume  $J = I_1 = \{1, 2, \dots, \theta\}$ , and  $\Phi_J^+ = \{\beta_1, \beta_2, \dots, \beta_p\}$ . We apply the results and notations of Section 2.2 to the connected



component  $I_1$ . For all  $a = (a_1, \dots, a_{\tilde{\theta}}) \in \mathbb{N}^{\tilde{\theta}}$ , and  $1 \leq i \leq \theta$ , we will show that

$$(4.7) \quad \mu_a h^a x_i = \mu_a x_i h^a.$$

We can assume that  $\mu_a \neq 0$ . Let  $1 \leq l \leq \tilde{\theta}$ , and  $\beta_l = \sum_{j=1}^{\tilde{\theta}} n_j \alpha_j$ , where  $n_j \in \mathbb{N}$  for all  $1 \leq j \leq \tilde{\theta}$ . Then by definition,  $g_{\beta_l} = \prod_{1 \leq j \leq \tilde{\theta}} g_j^{n_j}$ , and  $\chi_{\beta_l} = \prod_{1 \leq j \leq \tilde{\theta}} \chi_j^{n_j}$ . Hence

$$\chi_i(g_{\beta_l}^N) \chi_{\beta_l}^N(g_i) = \prod_{1 \leq j \leq \tilde{\theta}} q_{ii}^{a_{ij} N n_j} = 1,$$

since  $q_{ii}^N = 1$ , if  $i \in I_1$ , and  $a_{ij} = 0$ , if  $i \notin I_1$ . By Lemma 2.11,  $\chi_{\beta_l}^N = \varepsilon$  for all  $1 \leq l \leq \tilde{\theta}$  with  $a_l > 0$ . Hence  $\chi_i(g_{\beta_l}^N) = 1$  for all  $l$  with  $a_l > 0$ . This implies (4.7) since  $h^a x_i = \chi_i(h^a) x_i h^a$ .

Finally we prove by induction on  $\text{ht}(\underline{a})$  using (4.7) and (2.14) that  $u^a$  is central in  $U(\mathcal{D}, \lambda)$  (and in  $k\langle x_1, \dots, x_{\theta} \rangle \# k[\Gamma]$ ).  $\square$

**Theorem 4.4.** *Let  $\mathcal{D}$  be a datum of finite Cartan type satisfying (3.1) and (3.2). Let  $\lambda$  and  $\mu$  be families of linking and root vector parameters for  $\mathcal{D}$ . Then  $u(\mathcal{D}, \lambda, \mu)$  is a quotient Hopf algebra of  $U(\mathcal{D}, \lambda)$  with group-like elements  $G(u(\mathcal{D}, \lambda, \mu)) \cong \Gamma$ , and the elements*

$$x_{\beta_1}^{a_1} x_{\beta_2}^{a_2} \cdots x_{\beta_p}^{a_p} g, \quad 0 \leq a_l < N_l, \quad 1 \leq l \leq p, \quad g \in \Gamma$$

form a basis of  $u(\mathcal{D}, \lambda, \mu)$ . In particular,

$$\dim u(\mathcal{D}, \lambda, \mu) = \prod_{J \in \mathcal{X}} N_J^{|\Phi_J^+|} |\Gamma|.$$

*Proof.* By Theorem 3.3, the elements

$$x_{\beta_1}^{a_1} x_{\beta_2}^{a_2} \cdots x_{\beta_p}^{a_p} g, \quad 0 \leq a_l, \quad 1 \leq l \leq p, \quad g \in \Gamma$$

are a basis of  $U(\mathcal{D}, \lambda)$ . We want to apply Theorem 4.1 with

$$y_l = x_{\beta_l}, \quad \psi_l = \chi_{\beta_l}, \quad u_l = u_{\beta_l}(\mu), \quad 1 \leq l \leq p.$$

For each connected component  $J \in \mathcal{X}$  we apply the results of Section 2.2 with

$$\eta_l = \chi_{\beta_l}^{N_l}, \quad 1 \leq l \leq p, \quad \beta_l \in \Phi_J^+.$$

If  $\chi_{\beta_l}^{N_l} \neq \varepsilon$  for some  $1 \leq l \leq p$ ,  $\beta_l \in \Phi_J^+$ , then by assumption,  $\mu_{\beta_l} = 0$ , and by Lemma 2.11,  $u_{\beta_l}(\mu) = 0$ . By Lemma 4.7,  $u_{\beta_l}(\mu)$  is central in  $U(\mathcal{D}, \lambda)$ . Hence the claim concerning the basis of  $u(\mathcal{D}, \lambda, \mu)$  follows from Theorem 3.3 and Theorem 4.1.

We now show that  $u(\mathcal{D}, \lambda, \mu)$  is a Hopf algebra. Let  $J \in \mathcal{X}$ . We denote the restriction of  $\mathcal{D}$  to the connected component  $J$  by  $\mathcal{D}_J$ . By

Theorem 2.12, the map  $\varphi_\mu : K(\mathcal{D}_J)\#k[\Gamma] \rightarrow k[\Gamma]$  is a Hopf algebra homomorphism. The kernel of  $\varphi_\mu$  is generated by all  $x_\alpha^{N_J} - u_\alpha(\mu)$ ,  $\alpha \in \Phi_J^+$ . Hence the elements  $x_\alpha^{N_J} - u_\alpha(\mu)$ ,  $\alpha \in \Phi_J^+$ , generate a Hopf ideal in  $K(\mathcal{D}_J)\#k[\Gamma]$  and in  $U(\mathcal{D}, \lambda)$ .

The Hopf algebra  $u(\mathcal{D}, \lambda, \mu)$  is generated by the skew-primitive elements  $x_1, \dots, x_\theta$  and the image of  $\Gamma$ . In particular,  $G(u(\mathcal{D}, \lambda, \mu)) \cong \Gamma$ .  $\square$

For explicit examples of the Hopf algebras  $u(\mathcal{D}, \lambda, \mu)$  see [AS5, Section 6] for type  $A_n$ ,  $n \geq 1$ , and [BDR] for type  $B_2$ . In these papers, and for these types, the elements  $u_\alpha(\mu)$  are precisely written down. It is an interesting problem to find an explicit algorithm describing the elements  $u_\alpha(\mu)$  for any connected Dynkin diagram.

### 5. THE ASSOCIATED GRADED HOPF ALGEBRA

**5.1. Nichols algebras.** To determine the structure of a given pointed Hopf algebra, we proceed as in [AS1] and study the associated graded Hopf algebra.

Let  $A$  be a pointed Hopf algebra with group of group-like elements  $G(A) = \Gamma$ . Let

$$A_0 = k[\Gamma] \subset A_1 \subset \dots \subset A, \quad A = \bigcup_{n \geq 0} A_n$$

be the coradical filtration of  $A$ . We define the associated graded Hopf algebra [M, 5.2.8] by

$$\text{gr}(A) = \bigoplus_{n \geq 0} A_n/A_{n-1}, \quad A_{-1} = 0.$$

Then  $\text{gr}(A)$  is a pointed Hopf algebra with the same dimension and coradical as  $A$ . The projection map  $\pi : \text{gr}(A) \rightarrow k[\Gamma]$  and the inclusion  $\iota : k[\Gamma] \rightarrow \text{gr}(A)$  are Hopf algebra maps with  $\iota\pi = \text{id}_{k[\Gamma]}$ . Let

$$(5.1) \quad R = \{x \in \text{gr}(A) \mid (\text{id} \otimes \pi)\Delta(x) = x \otimes 1\}$$

be the algebra of  $k[\Gamma]$ -coinvariant elements. Then  $R = \bigoplus_{n \geq 0} R(n)$  is a graded Hopf algebra in  ${}^\Gamma\mathcal{YD}$ , and by (1.7)

$$(5.2) \quad \text{gr}(A) \cong R\#k[\Gamma].$$

Let  $V = P(R) \in {}^\Gamma\mathcal{YD}$  be the Yetter-Drinfeld module of primitive elements in  $R$ . We call its braiding

$$c : V \otimes V \rightarrow V \otimes V$$

the *infinitesimal braiding* of  $A$ .

Let  $\mathfrak{B}(V)$  be the subalgebra of  $R$  generated by  $V$ . Thus  $B = \mathfrak{B}(V)$  is the *Nichols algebra* of  $V$  [AS2], that is,

$$(5.3) \quad B = \bigoplus_{n \geq 0} B(n) \text{ is a graded Hopf algebra in } {}_{\Gamma}\mathcal{YD},$$

$$(5.4) \quad B(0) = k1, \quad B(1) = V,$$

$$(5.5) \quad B(1) = P(B),$$

$$(5.6) \quad B \text{ is generated as an algebra by } B(1).$$

$\mathfrak{B}(V)$  only depends on the vector space  $V$  with its Yetter-Drinfeld structure (see the discussion in [AS5, Section 2]). As an algebra and coalgebra,  $\mathfrak{B}(V)$  only depends on the braided vector space  $(V, c)$ .

We assume in addition that  $A$  is finite-dimensional and  $\Gamma$  is abelian. Then there are  $g_1, \dots, g_\theta \in \Gamma$ ,  $\chi_1, \dots, \chi_\theta \in \widehat{\Gamma}$  and a basis  $x_1, \dots, x_\theta$  of  $V$  such that  $x_i \in V_{g_i}^{\chi_i}$  for all  $1 \leq i \leq \theta$ . We call

$$(q_{ij} = \chi_j(g_i))_{1 \leq i, j \leq \theta}$$

the *infinitesimal braiding matrix* of  $A$ .

The first step to classify pointed Hopf algebras is the computation of the Nichols algebra.

Using results of Lusztig [L1],[L2], Rosso [Ro] and Müller [M1] and twisting we proved in [AS4, Theorem 4.5] the following description of the Nichols algebra of Yetter-Drinfeld modules of finite Cartan type.

**Theorem 5.1.** *Let  $\mathcal{D} = \mathcal{D}(\Gamma, (g_i)_{1 \leq i \leq \theta}, (\chi_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta})$  be a datum of finite Cartan type with finite abelian group  $\Gamma$ . Assume (3.1) and (3.2). Let  $V \in {}_{\Gamma}\mathcal{YD}$  be a vector space with basis  $x_1, \dots, x_\theta$  and  $x_i \in V_{g_i}^{\chi_i}$  for all  $1 \leq i \leq \theta$ . Then  $\mathfrak{B}(V)$  is the quotient algebra of  $T(V)$  modulo the ideal generated by the elements*

$$(5.7) \quad \text{ad}_c(x_i)^{1-a_{ij}}(x_j) \text{ for all } 1 \leq i, j \leq \theta, i \neq j,$$

$$(5.8) \quad x_\alpha^{N_J} \text{ for all } \alpha \in \Phi_J^+, J \in \mathcal{X}.$$

**Corollary 5.2.** *Assume the situation of Theorem 5.1, and let  $\lambda$  and  $\mu$  be linking and root vector parameters for  $\mathcal{D}$ . Then*

$$\text{gr}(u(\mathcal{D}, \lambda, \mu)) \cong u(\mathcal{D}, 0, 0) \cong \mathfrak{B}(V) \# k[\Gamma].$$

*Proof.* Let  $A = u(\mathcal{D}, \lambda, \mu)$ . There is a well-defined Hopf algebra map

$$u(\mathcal{D}, 0, 0) \rightarrow \text{gr}(u(\mathcal{D}, \lambda, \mu)),$$

mapping  $x_i, 1 \leq i \leq \theta$ , onto the residue class of  $x_i$  in  $A_1/A_0$ , and  $g \in \Gamma$  onto  $g$ . Since  $\dim(u(\mathcal{D}, 0, 0)) = \dim(u(\mathcal{D}, \lambda, \mu)) = \dim(\text{gr}(u(\mathcal{D}, \lambda, \mu)))$  by Theorem 4.4, it follows that  $u(\mathcal{D}, 0, 0) \cong \text{gr}(u(\mathcal{D}, \lambda, \mu))$ . By Theorem 5.1,  $u(\mathcal{D}, 0, 0) \cong \mathfrak{B}(V) \# k[\Gamma]$ .  $\square$

In [AS2] and [AS4] we determined the structure of finite-dimensional Nichols algebras assuming that  $V$  is of Cartan type and satisfies some more assumptions in the case of small orders ( $\leq 17$ ) of the diagonal elements  $q_{ii}$ . Recent results of Heckenberger [H1], [H2], [H3] together with Theorem 5.1 allow to prove the following very general structure theorem on Nichols algebras.

**Theorem 5.3.** *Let  $\Gamma$  be a finite abelian group, and  $V \in {}_{\Gamma}\mathcal{YD}$  a Yetter-Drinfeld module such that  $\mathfrak{B}(V)$  is finite-dimensional. Choose a basis  $x_i \in V$  with  $x_i \in V_{g_i}^{\chi_i}$ ,  $g_i \in \Gamma$ ,  $\chi_i \in \widehat{\Gamma}$ , for all  $1 \leq i \leq \theta$ . For all  $1 \leq i, j \leq \theta$ , define  $q_{ij} = \chi_j(g_i)$ , and assume*

$$(5.9) \quad \text{ord}(q_{ij}) \text{ is odd, and } \text{ord}(q_{ii}) \text{ is not } 3,$$

$$(5.10) \quad \text{ord}(q_{ii}) \text{ is prime to } 3 \text{ if } q_{il}q_{li} \in \{q_{ii}^{-3}, q_{ll}^{-3}\} \text{ for some } l.$$

*Then there is a datum  $\mathcal{D} = \mathcal{D}(\Gamma, (g_i)_{1 \leq i \leq \theta}, (\chi_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta})$  of finite Cartan type such that*

$$\mathfrak{B}(V) \# k[\Gamma] \cong u(\mathcal{D}, 0, 0).$$

*Proof.* For all  $1 \leq i, j \leq \theta, i \neq j$ , let  $V_{ij}$  be the vector subspace of  $V$  spanned by  $x_i, x_j$ . Then  $\mathfrak{B}(V_{ij})$  is isomorphic to a subalgebra of  $\mathfrak{B}(V)$ , hence it is finite-dimensional. Heckenberger [H1], [H2] classified finite-dimensional Nichols algebras of rank 2. By (5.9) it follows from the list in [H1, Theorem 4] that  $V_{ij}$  is of finite Cartan type, that is, there are  $a_{ij}, a_{ji} \in \{0, -1, -2, -3\}$  with  $a_{ij}a_{ji} \in \{0, 1, 2, 3\}$ , and

$$q_{ij}q_{ji} = q_{ii}^{a_{ij}} = q_{jj}^{a_{ji}}.$$

Since  $\mathfrak{B}(V) \# k[\Gamma]$  is finite-dimensional,  $q_{ii} \neq 1$  for all  $1 \leq i \leq \theta$  by [AS1, Lemma 3.1]. Thus  $(q_{ij})_{1 \leq i, j \leq \theta}$  is of Cartan type in the sense of [AS2, page 4] with (generalized) Cartan matrix  $(a_{ij})$ . In [H3, Theorem 4] Heckenberger extended part (ii) of [AS2, Theorem 1.1] (where we had to exclude some small primes) and showed that a diagonal braiding  $(q_{ij})$  of a braided vector space  $V$  is of finite Cartan type if it is of Cartan type and  $\mathfrak{B}(V)$  is finite-dimensional. Hence  $(a_{ij})$  is a Cartan matrix of finite type, and the claim follows from Theorem 5.1.  $\square$

**5.2. Generation in degree one.** We generalize our results in [AS4, Section 7]. Let  $A$  be a finite-dimensional pointed Hopf algebra with  $\Gamma, V$ , and  $R$  as in Section 5.1. To prove that  $\mathfrak{B}(V) = R$ , we dualize. Let  $S = R^*$  the dual Hopf algebra in  ${}_{\Gamma}\mathcal{YD}$  as in [AS2, Lemma 5.5]. Then  $S = \bigoplus_{n \geq 0} S(n)$  is a graded Hopf algebra in  ${}_{\Gamma}\mathcal{YD}$ , and by [AS2, Lemma 5.5],  $R$  is generated in degree one, that is,  $\mathfrak{B}(V) = R$ , if and only if  $P(S) = S(1)$ . The dual vector space  $S(1)$  of  $V = R(1)$  has the same braiding  $(q_{ij})$  (with respect to the dual basis) as  $V$ . Our strategy

to show  $P(S) = S(1)$  is to identify  $S$  as a Nichols algebra. In the next Lemma we use [H1, H2] to prove a very general version of [AS4, Lemma 7.2].

**Lemma 5.4.** *Let  $\mathcal{D} = \mathcal{D}(\Gamma, (g_i)_{1 \leq i \leq \theta}, (\chi_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta})$  be a datum of finite Cartan type with finite abelian group  $\Gamma$ . Let  $S = \bigoplus_{n \geq 0} S_n$  be a finite-dimensional graded Hopf algebra in  ${}_{\Gamma}^{\Gamma} \mathcal{YD}$  with  $S(0) = k1$ , and let  $x_1, \dots, x_{\theta}$  be a basis of  $S(1)$  with  $x_i \in S(1)_{g_i}^{\chi_i}$  for all  $1 \leq i \leq \theta$ . Assume for all  $1 \leq i \leq \theta$  that the order of  $q_{ii} = \chi_i(g_i)$  is odd and  $> 7$ . Then*

$$(5.11) \quad \text{ad}_c(x_i)^{1-a_{ij}}(x_j) = 0 \text{ for all } 1 \leq i, j \leq \theta, i \neq j.$$

*Proof.* We first note that the Nichols algebra of the primitive elements  $P(S) \in {}_{\Gamma}^{\Gamma} \mathcal{YD}$  is finite-dimensional. This can be seen by looking at  $\text{gr}(S \# k[\Gamma])$ .

Assume that there are  $1 \leq i, j \leq \theta, i \neq j$ , with  $\text{ad}_c(x_i)^{1-a_{ij}}(x_j) \neq 0$ . We define

$$y_1 = x_1, y_2 = \text{ad}_c(x_i)^{1-a_{ij}}(x_j).$$

By [AS2, A.1],  $y_2$  is a primitive element. Since  $y_1, y_2$  are non-zero elements of different degree, they are linearly independent. We know that the Nichols algebra of  $W = ky_1 + ky_2$  is finite-dimensional, since  $B(P(S))$  is finite-dimensional. We denote

$$h_1 = g_i, h_2 = g_i^{1-a_{ij}} \in \Gamma, \text{ and } \eta_1 = \chi_i, \eta_2 = \chi_i^{1-a_{ij}} \chi_j \in \widehat{\Gamma}.$$

Thus  $y_i \in S_{h_i}^{\eta_i}, 1 \leq i \leq 2$ . Let  $(Q_{ij} = \eta_j(h_i))_{1 \leq i, j \leq 2}$  be the braiding matrix of  $y_1, y_2$ . We compute

$$Q_{11} = q_{ii}, Q_{22} = q_{ii}^{1-a_{ij}} q_{jj}, Q_{12}Q_{21} = q_{ii}^{2-a_{ij}}.$$

By assumption, the order of  $Q_{11} = q_{ii}$  is odd and  $> 3$ . Since  $B(W)$  is finite-dimensional,  $Q_{22} \neq 1$  by [AS1, Lemma 3.1]. Thus  $Q_{22}$  has odd order, since the orders of  $q_{ii}, q_{jj}$  are odd. By checking Heckenberger's list in [H1, Theorem 4], and thanks to [H2], we see that the braiding  $(Q_{ij})$  is of finite Cartan type or that we are in case (T3) with

$$Q_{12}Q_{21} = Q_{11}^{-1}.$$

Hence there exists  $A_{12} \in \{0, -1, -2, -3\}$  with

$$Q_{12}Q_{21} = Q_{11}^{A_{12}}.$$

Since  $Q_{12}Q_{21} = q_{ii}^{2-a_{ij}}$ , and  $Q_{11} = q_{ii}$ , it follows that the order of  $q_{ii}$  divides  $2 - a_{ij} - A_{12} \in \{2, 3, 4, 5, 6, 7, 8\}$ . This is a contradiction since the order of  $q_{ii}$  is odd and  $> 7$ .  $\square$

The next theorem is one of the main results of this paper.

**Theorem 5.5.** *Let  $A$  be a finite-dimensional pointed Hopf algebra with abelian group  $G(A) = \Gamma$  and infinitesimal braiding matrix  $(q_{ij})_{1 \leq i, j \leq \theta}$ . Assume for all  $1 \leq i, j \leq \theta$ , that the order of  $q_{ij}$  is odd, the order of  $q_{ii}$  is  $> 7$ , and that (5.10) holds. Then  $A$  is generated by group-like and skew-primitive elements, that is,*

$$R = \mathfrak{B}(V),$$

where  $R$  is defined by (5.1), and  $V = R(1)$ .

*Proof.* We argue as in the proof of [AS4, Theorem 7.6]. Let  $S = R^*$  be the dual Hopf algebra in  ${}^{\Gamma}\mathcal{YD}$ . Then  $S(1) = R(1)^*$  has the same braiding  $(q_{ij})$  as  $R(1)$  with respect to the dual basis  $(x_i)$  of the corresponding basis of  $R(1)$ . By Theorem 5.3  $(q_{ij})$  is of finite Cartan type. By Lemma 5.4 the Serre relations (5.7) hold for the elements  $x_i$ . Then the root vector relations (5.8) follow by [AS4, Lemma 7.5]. Hence  $S \cong \mathfrak{B}(S(1))$  by Theorem 5.1, and  $S(1) = P(S)$ . By duality,  $R$  is a Nichols algebra.  $\square$

## 6. LIFTING

From Section 5 we know a presentation of  $\text{gr}(A)$  by generators and relations under the assumptions of Theorems 5.3 and 5.5. To lift this presentation to  $A$  we need the following formulation of [AS1, Lemma 5.4] which is a consequence of the theorem of Taft and Wilson [M, Theorem 5.4.1]. Here it is crucial that the group is abelian.

**Lemma 6.1.** *Let  $A$  be a finite-dimensional pointed Hopf algebra with abelian group  $G(A) = \Gamma$ . Write  $\text{gr}(A) \cong R \# k[\Gamma]$  as in (5.2), and let  $V = R(1)$  with basis  $x_i \in V_{g_i}^{\chi_i}$ ,  $g_i \in \Gamma$ ,  $\chi_i \in \widehat{\Gamma}$ ,  $1 \leq i \leq \theta$ . Let  $A_0 \subset A_1$  be the first two terms of the coradical filtration of  $A$ . Then*

$$(6.1) \quad \bigoplus_{g, h \in \Gamma, \varepsilon \neq \chi \in \widehat{\Gamma}} P_{g, h}^{\chi}(A) \xrightarrow{\cong} A_1/A_0 \xleftarrow{\cong} V \# k[\Gamma].$$

$$(6.2) \quad \text{For all } g \in \Gamma, P_{g, 1}(A)^{\varepsilon} = k(1 - g), \text{ and if } \varepsilon \neq \chi \in \widehat{\Gamma}, \text{ then}$$

$$(6.3) \quad P_{g, 1}(A)^{\chi} \neq 0 \iff g = g_i, \chi = \chi_i, \text{ for some } 1 \leq i \leq \theta.$$

We can now prove our main structure theorem.

**Theorem 6.2.** *Let  $A$  be a finite-dimensional pointed Hopf algebra with abelian group  $G(A) = \Gamma$  and infinitesimal braiding matrix  $(q_{ij})_{1 \leq i, j \leq \theta}$ . Assume for all  $1 \leq i, j \leq \theta$ , that the order of  $q_{ij}$  is odd, the order of  $q_{ii}$  is  $> 7$ , and that (5.10) holds. Then*

$$A \cong u(\mathcal{D}, \lambda, \mu),$$

where  $\mathcal{D} = \mathcal{D}(\Gamma, (g_i)_{1 \leq i \leq \theta}, (\chi_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta})$  is a datum of finite Cartan type, and  $\lambda$  and  $\mu$  are families of linking and root vector parameters for  $\mathcal{D}$ .

*Proof.* By Theorems 5.3 and 5.5, there is a datum  $\mathcal{D}$  of finite Cartan type such that  $\text{gr}(A) \cong u(\mathcal{D}, 0, 0)$ . By Lemma 6.1, for all  $1 \leq i \leq \theta$  we can choose

$$a_i \in P(A)_{g_i, 1}^{x_i} \text{ corresponding to } x_i \text{ in (6.1).}$$

We have shown in Theorem [AS4, 6.8] that

$$\begin{aligned} \text{ad}_c(a_i)^{1-a_{ij}}(a_j) &= 0, \text{ for all } 1 \leq i, j \leq \theta, i \sim j, i \neq j, \\ a_i a_j - q_{ij} a_j a_i - \lambda_{ij}(1 - g_i g_j) &= 0, \text{ for all } 1 \leq i < j \leq \theta, i \not\sim j, \end{aligned}$$

for some family  $\lambda$  of linking parameters. Thus there is a homomorphism of Hopf algebras

$$\varphi : U(\mathcal{D}, \lambda) \rightarrow A, \varphi|_{\Gamma} = \text{id}_{\Gamma}, \varphi(x_i) = a_i, \text{ for all } 1 \leq i \leq \theta.$$

By Theorem 5.5,  $\varphi$  is surjective.

We now use the notation of Section 2.2 and show that

$$(6.4) \quad \varphi(x_{\alpha}^{N_J}) \in k[\Gamma] \text{ for all } \alpha \in \Phi_J^+, J \in \mathcal{X}.$$

We fix  $J \in \mathcal{X}$  with  $p = |\Phi_J^+|$ , and show by induction on  $\text{ht}(\underline{a})$  that

$$(6.5) \quad \varphi(z^a) \in k[\Gamma] \text{ for all } a \in \mathbb{N}^p.$$

Let  $0 \neq a \in \mathbb{N}^p$ . Since  $\varphi$  is a Hopf algebra map, we see from (2.16) that

$$\Delta(\varphi(z^a)) = h^a \otimes \varphi(z^a) + \varphi(z^a) \otimes 1 + w,$$

where by induction

$$w = \sum_{b, c \neq 0, b+c=a} t_{b,c}^a \varphi(z^b) h^c \otimes \varphi(z^c) \in k[\Gamma] \otimes k[\Gamma].$$

In particular,  $\varphi(z^a) \in A_1$  by definition of the coradical filtration. We multiply this equation with  $g \otimes g, g \in \Gamma$ , from the left and  $g^{-1} \otimes g^{-1}$  from the right. Since  $gz^a g^{-1} = \eta^a(g)z^a$ , we obtain  $w = \eta^a(g)w$  for all  $g \in \Gamma$ .

Suppose  $\eta^a \neq \varepsilon$ . Then  $w = 0$ , and  $\varphi(z^a) \in P_{h^a, 1}^{\eta^a}$ . Then  $\varphi(z^a) = 0$  by Lemma 6.1 (6.3), since  $\chi_l(g_l) \neq 1$  for all  $1 \leq l \leq \theta$ , but  $\eta^a(h^a) = 1$  by the Cartan condition (see the proof of [AS2, Lemma 7.5] for a similar computation).

If  $\eta^a = \varepsilon$ , then  $\varphi(z^a) \in A_1^{\varepsilon} = k[\Gamma]$  by Lemma 6.1 (6.2).

This proves (6.5) and (6.4). Then we conclude for each  $J \in \mathcal{X}$  from Theorem 2.12 that the map

$$K(\mathcal{D}_J) \# k[\Gamma] \rightarrow U(\mathcal{D}, \lambda) \xrightarrow{\varphi} A$$

has the form  $\varphi_{\mu^J}$  for some family of scalars  $\mu^J$  as in Theorem 2.12 for the connected component  $J$ . Define  $\mu = (\mu_{\alpha})_{\alpha \in \Phi_+}$  by  $\mu_{\alpha} = \mu_{\alpha}^J$

for all  $\alpha \in \Phi_J^+$ . Then  $\mu$  is a family of root vector parameters for  $\mathcal{D}$ , and the elements  $u_\alpha(\mu) \in k[\Gamma]$  are defined in (4.5) for each  $J \in \mathcal{X}$  and  $\alpha \in \Phi_J^+$ . It follows that  $\varphi(x_\alpha^{N_J}) = u_\alpha(\mu) = \varphi(u_\alpha(\mu))$  for all  $J \in \mathcal{X}, \alpha \in \Phi_J^+$ . Thus  $\varphi$  factorizes over  $u(\mathcal{D}, \lambda, \mu)$ . Since  $\dim(A) = \dim(\text{gr}(A)) = \dim(u(\mathcal{D}, \lambda, 0, 0)) = \dim(u(\mathcal{D}, \lambda, \mu))$  by Theorem 4.4,  $\varphi$  induces an isomorphism  $u(\mathcal{D}, \lambda, \mu) \cong A$ .  $\square$

**Corollary 6.3.** *Let  $A$  be a finite-dimensional pointed Hopf algebra with abelian group  $G(A) = \Gamma$  satisfying the assumptions of Theorem 6.2. Then for each prime divisor  $p$  of the dimension of  $A$  there is a group-like element of order  $p$  in  $A$ .*

*Proof.* This follows from Theorems 6.2 and 4.4.  $\square$

We note that the analog of Cauchy's theorem in group theory is false for arbitrary, non-pointed Hopf algebras. Let  $A$  be a finite-dimensional Hopf algebra with only trivial group-like elements, such as the dual of the group algebra of a finite group  $G$  with  $G = [G, G]$ . Then  $A$  does not contain any Hopf subalgebra of prime dimension, since any Hopf algebra of prime dimension is a group algebra by Zhu's theorem [Z].

## 7. ISOMORPHISM CLASSES

In this last section we determine all isomorphisms between the Hopf algebras  $u(\mathcal{D}, \lambda, \mu)$ .

For convenience we introduce a normalization condition for Cartan matrices and their root systems. Let  $(a'_{ij})_{1 \leq i, j \leq \theta}$  and  $(a_{ij})_{1 \leq i, j \leq \theta}$  be Cartan matrices of finite type. A *diagram isomorphism* between  $(a'_{ij})$  and  $(a_{ij})$  is a permutation  $\tau$  of  $\{1, 2, \dots, \theta\}$  with  $a'_{ij} = a_{\tau(i), \tau(j)}$  for all  $1 \leq i, j \leq \theta$ . We choose from each isomorphism class of connected Cartan matrices of finite type one representative. The chosen representatives are called *standard* Cartan matrices. We fix a reduced representation of the longest element in the Weyl group and the corresponding ordering of the positive roots of the standard matrices as described in Section 2.1.

From now on we assume that any Cartan datum  $\mathcal{D}$  satisfies the following *additional normalizing condition*:

The Cartan matrix  $(a_{ij})_{1 \leq i, j \leq \theta}$  of  $\mathcal{D}$  is a block diagonal matrix, and each matrix on the diagonal is one of the standard connected Cartan matrices. Moreover for each connected component  $J$  of  $I$  we fix the same order of the positive roots as for the chosen representative.

Thus up to a shift of indices we can identify the Cartan matrix of any connected component with a standard Cartan matrix. We use the



fixed ordering of the positive roots in each component to define the root vectors of  $\mathcal{D}$ .

Note that any diagram isomorphism induces an isomorphism of the corresponding Nichols algebras. Hence up to Hopf algebra isomorphisms we can assume by the proof of Theorem 6.2 that the normalizing condition is satisfied for the Hopf algebras  $u(\mathcal{D}, \lambda, \mu)$ .

In the next definition we extend the notation for the linking parameters in Definition 3.1 by defining  $\lambda_{ij} = -q_{ij}\lambda_{ji}$  for all  $i \approx j, i > j$ .

**Definition 7.1.** Let  $\mathcal{D} = \mathcal{D}(\Gamma, (g_i)_{1 \leq i \leq \theta}, (\chi_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta})$  and  $\mathcal{D}' = \mathcal{D}(\Gamma', (g'_i)_{1 \leq i \leq \theta'}, (\chi'_i)_{1 \leq i \leq \theta'}, (a'_{ij})_{1 \leq i, j \leq \theta'})$  be Cartan data of finite type satisfying (3.1) and (3.2) and with finite abelian groups  $\Gamma, \Gamma'$ . Assume  $\theta = \theta'$ . Let  $\lambda$  and  $\lambda'$  be linking parameters, and  $\mu$  and  $\mu'$  root vector parameters for  $\mathcal{D}$  and  $\mathcal{D}'$ .

Let  $\varphi : \Gamma' \rightarrow \Gamma$  be a group isomorphism,  $\sigma \in S_\theta$  a permutation and  $(s_i)_{1 \leq i \leq \theta}$  a family of non-zero elements in  $k$ . The triple  $(\varphi, \sigma, (s_i))$  is called an *isomorphism* from  $(\mathcal{D}', \lambda', \mu')$  to  $(\mathcal{D}, \lambda, \mu)$  if the following five conditions are satisfied:

$$(7.1) \quad \varphi(g'_i) = g_{\sigma(i)} \text{ for all } 1 \leq i \leq \theta.$$

$$(7.2) \quad \chi'_i = \chi_{\sigma(i)} \varphi \text{ for all } 1 \leq i \leq \theta.$$

$$(7.3) \quad a'_{ij} = a_{\sigma(i), \sigma(j)} \text{ for all } 1 \leq i, j \leq \theta.$$

$$(7.4) \quad \lambda'_{ij} = s_i s_j \lambda_{\sigma(i)\sigma(j)} \text{ for all } 1 \leq i, j \leq \theta, i \approx j$$

To formulate the fifth condition we have to introduce more notations for any connected component  $J$  of  $\mathcal{D}$ .

- (1) Let  $q_J = (\chi_j(g_i))_{i, j \in J}$  be the braiding matrix of the restriction  $\mathcal{D}_J$  of  $\mathcal{D}$  to  $J$ .
- (2) Using (7.3) we identify the root systems of  $J$  and of  $\sigma^{-1}(J)$  with the corresponding root system of the standard Cartan matrix. Then the restriction of  $\sigma$  to  $\sigma^{-1}(J)$  becomes a diagram automorphism  $\sigma_J$  of the corresponding standard Cartan matrix.
- (3) For any  $\beta \in \Phi_J^+$  let  $u'_\beta(\mu')$  and  $u_\beta(\mu)$  be the elements in the group algebras  $k[\Gamma']$  and  $k[\Gamma]$  defined in (4.5). For any family  $a = (a_\beta)_{\beta \in \Phi_J^+}$  of natural numbers  $a_\beta \geq 0$  we define the product  $u(\mu)^a = \prod_{\beta \in \Phi_J^+} u_\beta(\mu)^{a_\beta}$ .
- (4) For any  $\alpha \in \Phi_J^+$  with  $\alpha = \sum_{i \in J} n_i \alpha_i, n_i \geq 0$  for all  $1 \leq i \leq \theta$ , let  $s_\alpha = \prod_{i \in J} s_i^{n_i}$ .
- (5) For any  $\alpha \in \Phi_J^+$  and any family  $a = (a_\beta)_{\beta \in \Phi_J^+}$  of natural numbers  $a_\beta \geq 0$  let  $t_{\alpha, q_J, \sigma_J}^a$  be the element in  $k$  defined below in Theorem 7.5 applied to  $\mathcal{D}_J$ .

Then the last condition is the following identity in the group algebra  $k[\Gamma]$ :

$$(7.5) \quad \varphi(u'_\alpha(\mu')) = s_\alpha^{N_J} \sum_a t_{\alpha, q_J, \sigma_J}^a u(\mu)^a \text{ for all } \alpha \in \Phi_J^+, J \in \mathcal{X}.$$

Finally let  $\text{Isom}(\mathcal{D}', \lambda', \mu'), (\mathcal{D}, \lambda, \mu)$  be the set of all isomorphisms from  $(\mathcal{D}', \lambda', \mu')$  to  $(\mathcal{D}, \lambda, \mu)$ .

For Hopf algebras  $A', A$  we denote by  $\text{Isom}(A', A)$  the set of all Hopf algebra isomorphisms from  $A'$  to  $A$ .

We now can state the main result of this section.

**Theorem 7.2.** *Let  $\mathcal{D}$  and  $\mathcal{D}'$  be Cartan data of finite type with finite abelian groups  $\Gamma$  and  $\Gamma'$  and rank  $\theta$  and  $\theta'$ . Assume that  $\mathcal{D}$  and  $\mathcal{D}'$  satisfy (3.1), (3.2). In addition assume the following condition on the braiding matrix of  $\mathcal{D}$ :*

$$(7.6) \quad \text{ord}(q_{ij}) > 4 \text{ for all } 1 \leq i, j \leq \theta.$$

*Let  $\lambda$  and  $\lambda'$  be linking parameters, and  $\mu$  and  $\mu'$  root vector parameters for  $\mathcal{D}$  and  $\mathcal{D}'$ .*

*If the Hopf algebras  $u(\mathcal{D}', \lambda', \mu')$  and  $u(\mathcal{D}, \lambda, \mu)$  are isomorphic, then  $\theta' = \theta$ .*

*Assume  $\theta' = \theta$ . Then the map*

$$\text{Isom}((\mathcal{D}', \lambda', \mu'), (\mathcal{D}, \lambda, \mu)) \rightarrow \text{Isom}(u(\mathcal{D}', \lambda', \mu'), u(\mathcal{D}, \lambda, \mu))$$

*given by  $(\varphi, \sigma, (s_i)) \mapsto F$ , where  $F(x'_i) = s_i x_{\sigma(i)}$  and  $F(g') = \varphi(g')$  for all  $1 \leq i \leq \theta$  and  $g' \in \Gamma'$ , is bijective.*

Before we begin with the proof of Theorem 7.2 we need some preparations.

First we see that condition (7.3) is in most cases redundant.

**Lemma 7.3.** *In the situation of Definition 7.1 assume  $\theta' = \theta$ , (7.1), (7.2) and (7.6). Then (7.3) holds.*

*Proof.* For all  $1 \leq i, j \leq \theta$  let  $q'_{ij} = \chi'_j(g'_i)$ ,  $q_{ij} = \chi_j(g_i)$ . Then for all  $i, j$  (7.1) and (7.2) imply that  $q'_{ij} = q_{\sigma(i)\sigma(j)}$ . Hence  $a_{ij} = a'_{\sigma(i)\sigma(j)}$ , since  $q_{ii}^{a_{ij}} = q_{ii}^{a'_{\sigma(i)\sigma(j)}}$ , and  $a_{ij} - a'_{\sigma(i)\sigma(j)} \in \{0, \pm 1, \pm 2, \pm 3\}$ .  $\square$

We need an extra information in the situation of Theorem 2.5.

**Lemma 7.4.** *Let  $\mathcal{D}$  be a connected Cartan datum of finite type with root system  $\Phi$ , finite abelian group  $\Gamma$  and  $N = \text{ord}(q_{ii})$  for all  $i$ . Assume (3.1) and (3.2). By Theorem 5.1 there is a canonical projection  $\pi : R(\mathcal{D}) \rightarrow \mathfrak{B}(V)$  whose kernel is the ideal generated by all  $x_\alpha^N, \alpha \in \Phi^+$ .*

We denote the coalgebra structure of  $R(\mathcal{D})$  by  $\Delta(x) = x^{(1)} \otimes x^{(2)}$  for all  $x \in R(\mathcal{D})$ . Then

$$K(\mathcal{D}) = R(\mathcal{D})^{\text{co}\pi} = \{x \in R(\mathcal{D}) \mid x^{(1)} \otimes \pi(x^{(2)}) = x \otimes 1\}.$$

*Proof.* This follows by bosonization from the corresponding result for pointed Hopf algebras in [Ma].  $\square$

In the following theorem we define the constants in (7.5).

**Theorem 7.5.** *Let  $\mathcal{D}(\Gamma, (g_i)_{1 \leq i \leq \theta}, (\chi_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta})$  be a connected Cartan datum of finite type with root system  $\Phi$ , finite abelian group  $\Gamma$ , and  $N = \text{ord}(g_{ii}), 1 \leq i \leq \theta$ . Assume that  $(a_{ij})_{1 \leq i, j \leq \theta}$  is a standard Cartan matrix, and let  $\sigma$  be a diagram automorphism of  $(a_{ij})_{1 \leq i, j \leq \theta}$ . Define*

$$\mathcal{D}^\sigma = \mathcal{D}(\Gamma, (g'_i)_{1 \leq i \leq \theta}, (\chi'_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta})$$

with  $g'_i = g_{\sigma(i)}, \chi'_i = \chi_{\sigma(i)}$  for all  $1 \leq i \leq \theta$ . Let  $V \in {}^\Gamma \mathcal{YD}$  with basis  $x_i \in V_{g_i}^{\chi_i}$ , and  $V^\sigma \in {}^\Gamma \mathcal{YD}$  with basis  $x_i^\sigma \in (V^\sigma)_{g'_i}^{\chi'_i}$  for all  $1 \leq i \leq \theta$ . Then there is an algebra map

$$F^\sigma : R(\mathcal{D}^\sigma) \rightarrow R(\mathcal{D}), x_i^\sigma \mapsto x_{\sigma(i)} \text{ for all } 1 \leq i \leq \theta.$$

Then for each  $\alpha \in \Phi^+$ , and each family  $a = (a_\beta)_{\beta \in \Phi^+}$  of natural numbers  $a_\beta \geq 0$  there are uniquely determined elements  $t_\alpha^a \in k$  depending on the braiding matrix of  $\mathcal{D}$  and the diagram automorphism  $\sigma$  such that

$$F^\sigma(x_\alpha^\sigma)^N = \sum_a t_\alpha^a z^a,$$

where  $z^a = x_{\beta_1}^{a_1 N} \cdots x_{\beta_p}^{a_p N}$  as in Definition 2.6 with the fixed ordering  $\beta_1, \dots, \beta_p$  of the positive roots, and where  $x_\alpha^\sigma$  denotes the root vector of  $\alpha$  in  $R(\mathcal{D}_\sigma)$ .

*Proof.* The Cartan condition is satisfied for  $\mathcal{D}^\sigma$  since  $\sigma$  is a diagram automorphism. The linear map  $f^\sigma : V^\sigma \rightarrow V, x_i^\sigma \mapsto x_{\sigma(i)}, 1 \leq i \leq \theta$ , is an isomorphism of Yetter-Drinfeld modules. Then  $f^\sigma$  induces isomorphisms  $F^\sigma : R(\mathcal{D}^\sigma) \rightarrow R(\mathcal{D})$  and  $\overline{F}^\sigma : \mathfrak{B}(V^\sigma) \rightarrow \mathfrak{B}(V)$ . Let  $\pi : R(V) \rightarrow \mathfrak{B}(V)$  and  $\pi^\sigma : R(V^\sigma) \rightarrow \mathfrak{B}(V^\sigma)$  be the natural projections. Since  $K(\mathcal{D}) = R(\mathcal{D})^{\text{co}\pi}$  and  $K(\mathcal{D}^\sigma) = R(\mathcal{D}^\sigma)^{\text{co}\pi^\sigma}$  by Lemma 7.4, it follows that  $F^\sigma$  maps  $K(\mathcal{D}^\sigma)$  into  $K(\mathcal{D})$ . This proves the claim by Theorem 2.5.  $\square$

The meaning of the elements  $F^\sigma(x_\alpha^\sigma)$  in the previous theorem can be explained as follows. Let  $x_\alpha$  be represented as iterated skew-commutator of simple root vectors  $x_{i_1}, \dots, x_{i_s}$  in this order. Then  $F^\sigma(x_\alpha^\sigma)$  is the same iterated skew-commutator of the sequence  $x_{\sigma(i_1)}, \dots, x_{\sigma(i_s)}$ .

As an example, take the Dynkin diagram  $A_2$  with the non-simple root  $\alpha = \alpha_1 + \alpha_2$  and the diagram automorphism  $\sigma$  with  $\sigma(1) = 2, \sigma(2) = 1$ . Then  $x_\alpha = x_1x_2 - q_{12}x_2x_1$  and  $F^\sigma(x_\alpha^\sigma) = x_2x_1 - q_{21}x_1x_2$ .

Finally we note

**Lemma 7.6.** *Let  $\mathcal{D} = \mathcal{D}(\Gamma, (g_i)_{1 \leq i \leq \theta}, (\chi_i)_{1 \leq i \leq \theta}, (a_{ij})_{1 \leq i, j \leq \theta})$  be a datum of finite Cartan type and assume (7.6). Then for all  $1 \leq i, j \leq \theta, i \neq j$ ,  $g_i \neq g_j$  or  $\chi_i \neq \chi_j$ .*

*Proof.* Assume there are  $i \neq j$  with  $g_i = g_j, \chi_i = \chi_j$ . Then  $q_{ii} = q_{jj}$ , and  $q_{ii}^2 = q_{ii}^{a_{ij}} = q_{ii}^{a_{ji}}$ . Hence in contradiction to our assumption we have  $q_{ii}^{2-a_{ij}} = 1$  for  $a_{ij} \in \{0, -1, -2\}$ , and  $q_{ii}^{a_{ij}-a_{ji}} = 1$  for  $a_{ij} = -3$ .  $\square$

We can now prove Theorem 7.2:

*Proof.* Assume that there is a Hopf algebra isomorphism

$$F : A' = u(\mathcal{D}', \lambda', \mu') \rightarrow A = u(\mathcal{D}, \lambda, \mu).$$

Then  $F$  preserves the coradical filtration and induces an isomorphism  $A'_0 = k[\Gamma'] \cong A_0 = k[\Gamma]$ , given by a group isomorphism  $\varphi : \Gamma' \rightarrow \Gamma$ , and by Corollary 5.2 an isomorphism

$$A'_1 = k[\Gamma'] \oplus \bigoplus_{\substack{g' \in \Gamma', \\ 1 \leq i \leq \theta'}} kx'_i g' \cong A_1 = k[\Gamma] \oplus \bigoplus_{\substack{g \in \Gamma, \\ 1 \leq i \leq \theta}} kx_i g.$$

Hence it follows from Lemma 7.6 (see [AS2, 6.3]) that  $\theta = \theta'$ , and that there are a permutation  $\sigma \in S_\theta$  and elements  $0 \neq s_i \in k, 1 \leq i \leq \theta$  such that (7.1) and (7.2) hold, and  $F(x'_i) = s_i x_{\sigma(i)}$  for all  $1 \leq i \leq \theta$ . Then Lemma 7.3 implies (7.3), and  $F([x'_i, x'_j]_{c'}) = s_i s_j [x_{\sigma(i)}, x_{\sigma(j)}]_c$  for all  $1 \leq i \leq \theta$ . Now (7.4) follows from the linking relations.

To establish (7.5) we fix a connected component  $J$  of  $\mathcal{D}$ , and we identify  $J$  and  $J' = \sigma^{-1}(J)$  with the index set  $\{1, \dots, \theta_J\}$  of the corresponding standard Cartan matrix. Then the restriction of  $\sigma$  to  $J'$  becomes the diagram automorphism  $\sigma_J$ . Let  $V_J \in {}_\Gamma \mathcal{YD}$  with basis  $x_i \in (V_J)_{g_i}^{x_i}, i \in J$ , and  $V'_J \in {}_{\Gamma'} \mathcal{YD}$  with basis  $x'_i \in (V'_J)_{g'_i}^{x'_i}, i \in J$ . Let  $f_J^{\sigma_J} : V'_J \rightarrow V_J$  be the map of Theorem 7.5 for  $\mathcal{D}_J$  and  $V_J$  instead of  $\mathcal{D}$  and  $V$ .

We define linear maps  $f_J : V'_J \rightarrow V_J, x'_i \mapsto s_i x_{\sigma(i)}, 1 \leq i \leq \theta_J$ , and  $f'_J : V'_J \rightarrow V_J, x'_i \mapsto s_i x_i^{\sigma_J}, 1 \leq i \leq \theta_J$ . Then  $f_J = f_J^{\sigma_J} f'_J$ . The maps  $f_J$  and  $f'_J$  are  $\Gamma'$ -linear and  $\Gamma$ -colinear, where the action of  $\Gamma'$  on  $V'_J$  and the coaction of  $\Gamma$  on  $V_J$  are defined via  $\varphi$ . Hence by (7.1)–(7.3) they induce algebra maps  $F_J : R(\mathcal{D}'_J) \rightarrow R(\mathcal{D}_J)$  and  $F'_J : R(\mathcal{D}'_J) \rightarrow R(\mathcal{D}_J^{\sigma_J})$ , and

$F_J = F_J^{\sigma_J} F'_J$ . By Theorem 7.5 we then obtain for any  $\alpha \in \Phi_J^+$

$$(7.7) \quad F_J(x'_\alpha)^{N_J} = s_\alpha^{N_J} F_J^{\sigma_J}(x_\alpha^{\sigma_J})^{N_J} = s_\alpha^{N_J} \sum_a t_{\alpha, q_J, \sigma_J}^a z^a.$$

The canonical maps  $\pi_J : R(\mathcal{D}_J) \rightarrow u(\mathcal{D}, \lambda, \mu)$  and  $\pi_{J'} : R(\mathcal{D}'_{J'}) \rightarrow u(\mathcal{D}', \lambda', \mu')$  map root vectors to root vectors. Since  $F\pi_{J'} = \pi_J F_J$ , we see from (7.7) that for all  $\alpha \in \Phi_J^+$

$$F\pi_{J'}(x'^{N_J}_\alpha) = \pi_J F_J(x'^{N_J}_\alpha) = s_\alpha^{N_J} \sum_a t_{\alpha, q_J, \sigma_J}^a u(\mu)^a.$$

On the other hand  $x'^{N_J}_\alpha = u'_\alpha(\mu')$  in  $u(\mathcal{D}', \lambda', \mu')$ , hence  $F\pi_{J'}(x'^{N_J}_\alpha) = \varphi(u'_\alpha(\mu'))$ , and (7.5) follows.

It is easy to see that conversely any isomorphism  $(\varphi, \sigma, (s_i))$  defines an isomorphism of Hopf algebras, and that two such triples coincide if they define the same Hopf algebra map.  $\square$

We remark that the situation greatly simplifies if the diagram automorphism  $\sigma_J$  in Definition 7.1 is the identity. This happens in particular if the Dynkin diagram of  $(a_{ij})_{i,j \in J}$  is not of Type  $A, D$  or  $E_6$ . In this case it follows from the inductive definition of the  $u_\alpha(\mu)$  that (7.5) is equivalent to

$$(7.8) \quad \mu'_\alpha = s_\alpha^{N_J} \mu_\alpha \text{ for all } \alpha \in \Phi_J^+, J \in \mathfrak{X}.$$

## REFERENCES

- [AS1] N. Andruskiewitsch and H.-J. Schneider, *Lifting of Quantum Linear Spaces and Pointed Hopf Algebras of order  $p^3$* , J. Algebra **209** (1998), 658–691.
- [AS2] ———, *Finite quantum groups and Cartan matrices*, Adv. in Math. **154** (2000), 1–45.
- [AS3] ———, *Lifting of Nichols algebras of type  $A_2$  and Pointed Hopf Algebras of order  $p^4$* , in "Hopf algebras and quantum groups", Proceedings of the Colloquium in Brussels 1998, ed. S. Caeneppel (2000), 1–16.
- [AS4] ———, *Finite quantum groups over abelian groups of prime exponent*, Ann. Sci. Ec. Norm. Super. **35** (2002), 1–26.
- [AS5] ———, *Pointed Hopf Algebras*, in: Recent developments in Hopf algebra Theory, MSRI Publications **43** (2002), 168, Cambridge Univ. Press.
- [AS6] ———, *A characterization of quantum groups*, J. reine angew. Math. **577** (2004), 81–104.
- [BDR] M. Beattie, S. Dăscălescu and S. Raianu, *Lifting of Nichols algebras of type  $B_2$* , Israel J. Math. **132** (2002), 1–28.
- [dCK] C. De Concini and V. G. Kac, *Representations of quantum groups at roots of 1*, in "Operator Algebras, Unitary Representations, Enveloping Algebras, and Invariant Theory", ed. A. Connes *et al* (2000); Birkhäuser, 471–506.
- [dCP] C. de Concini and C. Procesi, *Quantum Groups*, in "D-modules, Representation theory and Quantum Groups", 31–140, Lecture Notes in Maths. **1565** (1993), Springer-Verlag .

- [D1] D. Didt, *Linkable Dynkin diagrams*, J. Algebra **255** (2002), 373–391.
- [D2] ———, *Linkable Dynkin diagrams and Quasi-isomorphisms for finite dimensional pointed Hopf algebras*, PhD thesis, Ludwig-Maximilians-Universität München, 2002.
- [EO] P. Etingof and V. Ostrik, *Finite tensor categories*, Mosc. Math. J. **4** (2004), no. 3, 627–654, 782–783. math.QA/0301027.
- [Ga] Matthias R. Gaberdiel, *An algebraic approach to logarithmic conformal field theory*, Int.J.Mod.Phys. **A18** (2003) 4593–4638, hep-th/0111260.
- [H1] I. Heckenberger, *Finite dimensional rank 2 Nichols algebras of diagonal type I: Examples*, Preprint math.QA/0402350v2, 2004.
- [H2] I. Heckenberger, *Finite dimensional rank 2 Nichols algebras of diagonal type II: Classification*, Preprint math.QA/0404008, 2004.
- [H3] I. Heckenberger, *The Weyl-Brandt groupoid of a Nichols algebra of diagonal type*, Preprint math.QA/0411477, 2004.
- [K] V. Kharchenko, *A quantum analog of the Poincaré-Birkhoff-Witt theorem*, Algebra and Logic **38** (1999), 259–276.
- [KS] A. Klimyk and K. Schmüdgen, *Quantum Groups and Their Representations*, Springer, Texts and Monographs in Physics, 1997.
- [L1] G. Lusztig, *Finite dimensional Hopf algebras arising from quantized universal enveloping algebras*, J. of Amer. Math. Soc. **3** 257–296.
- [L2] G. Lusztig, *Quantum groups at roots of 1*, Geom. Dedicata **35** (1990), 89–114.
- [L3] G. Lusztig, *Introduction to quantum groups*, Birkhäuser, 1993.
- [Ma] A. Masuoka, *On Hopf algebras with cocommutative coradicals*, J. Algebra **144** (1991), 415–466.
- [M] S. Montgomery, *Hopf Algebras and Their Actions on Rings*, CBMS Conf. Series in Math., vol. 82, Amer. Math. Soc., Providence, RI, 1993.
- [M1] E. Müller, *Some Topics on Frobenius-Lusztig Kernels, I*, J. Algebra **206** (1998), 624–658.
- [M2] E. Müller, *The Coradical Filtration of  $U_q(\mathfrak{g})$  at Roots of Unity*, Comm. Algebra **28** (2000), 1029–1044.
- [Ri] C. Ringel, *Hall algebras and quantum groups*, Inventiones Math. **101** (1990), 583–591.
- [Ro] M. Rosso, *Quantum groups and quantum shuffles*, Inventiones Math. **133** (1998), 399–416.
- [S] Moss E. Sweedler, *Hopf algebras*, Mathematics Lecture Note Series W. A. Benjamin, New York, 1969.
- [Z] Y. Zhu, *Hopf algebras of prime dimension*, Int. Math. Res. Notes **1** (1994), 53–59.

FACULTAD DE MATEMÁTICA, ASTRONOMÍA Y FÍSICA, UNIVERSIDAD NACIONAL DE CÓRDOBA, CIEM - CONICET, (5000) CIUDAD UNIVERSITARIA, CÓRDOBA, ARGENTINA

*E-mail address:* andrus@mate.uncor.edu

MATHEMATISCHES INSTITUT, UNIVERSITÄT MÜNCHEN, THERESIENSTR. 39, D-80333 MUNICH, GERMANY

*E-mail address:* Hans-Juergen.Schneider@mathematik.uni-muenchen.de