

ON NONSINGULAR TWO-STEP NILPOTENT LIE ALGEBRAS

JORGE LAURET AND DAVID OSCARI

ABSTRACT. A 2-step nilpotent Lie algebra \mathfrak{n} is called *nonsingular* if $\text{ad } X : \mathfrak{n} \rightarrow [\mathfrak{n}, \mathfrak{n}]$ is onto for any $X \notin [\mathfrak{n}, \mathfrak{n}]$. We explore nonsingular algebras in several directions, including the classification problem (isomorphism invariants), the existence of canonical inner products (nilsolitons) and their automorphism groups (maximality properties). Our main tools are the moment map for certain real reductive representations, and the Pfaffian form of a 2-step algebra, which is a positive homogeneous polynomial in the nonsingular case.

CONTENTS

1. Introduction	1
2. Invariants of nonsingular algebras	4
2.1. Pfaffian form	6
2.2. Positive polynomials	8
2.3. Real geometric invariant theory	9
2.4. Nilsolitons	11
3. Nonsingular algebras of type $(2, m)$	14
4. Nonsingular algebras of type $(3, 8)$	18
4.1. Existence of nilsolitons	18
4.2. Non-existence of nilsolitons	19
5. Automorphisms of nonsingular algebras	20
5.1. 2-step algebras of <i>Rep</i> -type	23
References	23

1. INTRODUCTION

We study, for a given pair $\mathfrak{n}_1, \mathfrak{n}_2$ of real vector spaces, bilinear skew-symmetric maps

$$\mu : \mathfrak{n}_1 \times \mathfrak{n}_1 \longrightarrow \mathfrak{n}_2,$$

such that the function $(X, Y) \mapsto \alpha(\mu(X, Y))$ is a non-degenerate 2-form on \mathfrak{n}_1 for any nonzero $\alpha \in \mathfrak{n}_2^*$. This is precisely the condition the O'Neill tensor of a Riemannian submersion must necessarily satisfy at each point in order to get that all vertical 2-planes have positive sectional curvature, in which case the bundle is called *fat* (see [FZ11]). On the other hand, each of such μ 's defines a 2-step nilpotent Lie algebra $(\mathfrak{n} = \mathfrak{n}_1 \oplus \mathfrak{n}_2, \mu)$, often called *nonsingular* (also *regular*, or *fat*) in the literature under the above condition. The class of nonsingular nilpotent Lie groups strictly contains the well-known *H-type* groups introduced by A. Kaplan [K81], and their left-invariant metrics enjoy very nice properties

This research was partially supported by grants from CONICET, FonCyT (Argentina) and SeCyT (Universidad Nacional de Córdoba).

concerning curvature and totally geodesic submanifolds which were proved by P. Eberlein in [E94].

If $\dim \mathfrak{n}_1 = m$ and $\dim \mathfrak{n}_2 = n$, then the *isomorphism* relation between two nonsingular algebras is given by the natural $\mathrm{GL}_m \times \mathrm{GL}_n$ -action on the vector space $V_{n,m} := \Lambda^2 \mathfrak{n}_1^* \otimes \mathfrak{n}_2$ and coincides with Lie algebra isomorphism. Only very special values for the *type* (n, m) of a nonsingular algebra are allowed, which coincide with the types of *H-type* algebras and include $n = 1, m = 2k$; $n = 2, 3, m = 4k$; $n = 4, \dots, 7, m = 8k$; etc. The following natural questions arise from both the algebraic and geometric points of view:

- (i) How ‘wild’ is the classification problem for nonsingular algebras up to isomorphism?
- (ii) Given a nonsingular algebra $\mu : \mathfrak{n}_1 \times \mathfrak{n}_1 \longrightarrow \mathfrak{n}_2$, is there a canonical inner product on $\mathfrak{n}_1 \oplus \mathfrak{n}_2$ attached to μ ?
- (iii) Is any *H-type* algebra the ‘most symmetric’ one among all nonsingular algebras of the same type? This in the sense that its automorphism group has maximal dimension, or equivalently, its $\mathrm{GL}_m \times \mathrm{GL}_n$ -orbit in $V_{n,m}$ is of minimal dimension.

The classification of 2-step algebras is considered ‘hopeless’ from a rigorous point of view (see [BLS05]). A complete classification is only known for the types $(1, m)$ (all Heisenberg algebras up to abelian factors), $(2, m)$ (see [G73, LT99] or Section 3), $(5, 5)$ and (n, m) with $n + m \leq 9$ (over \mathbb{C} , see [GT99]). By using pencil invariants, one can show that nonsingular algebras of type $(2, m)$ are parameterized by sets of the form $\mathcal{S} = \{(\alpha_1, k_1), \dots, (\alpha_r, k_r)\}$, where $\alpha_1, \dots, \alpha_r \in \mathbb{C} \setminus \mathbb{R}$, and isomorphism classes correspond to $\mathrm{GL}_2(\mathbb{R})$ -orbits of such sets relative to the $\mathrm{GL}_2(\mathbb{R})$ -action on \mathbb{C} by Möbius transformations (see Section 3).

For a given inner product $\langle \cdot, \cdot \rangle$ on $\mathfrak{n} = \mathfrak{n}_1 \oplus \mathfrak{n}_2$ (with $\mathfrak{n}_1 \perp \mathfrak{n}_2$), one can encode the structural constants of μ in a map $J_\mu : \mathfrak{n}_2 \longrightarrow \mathfrak{so}(\mathfrak{n}_1)$ defined by

$$\langle J_\mu(Z)X, Y \rangle = \langle \mu(X, Y), Z \rangle, \quad \forall X, Y \in \mathfrak{n}_1, Z \in \mathfrak{n}_2.$$

There is a nice and useful isomorphism invariant for 2-step algebras (with m even) called the *Pfaffian form*, which is the projective equivalence class of the homogeneous polynomial f_μ of degree $m/2$ in n variables defined by

$$f_\mu(Z)^2 = \det J_\mu(Z), \quad \forall Z \in \mathfrak{n}_2,$$

for each μ of type (n, m) (see Section 2.1). It is an immediate, though quite an intriguing fact, that μ is nonsingular if and only if f_μ is a *positive polynomial* (i.e. $f_\mu(x) > 0$ for any nonzero $x \in \mathbb{R}^n$). Even though the theory of positive polynomials has a very rich presence in the literature, a complete classification of them, up to projective equivalence, seems to be as difficult as the classification of all forms and thus is also unknown and mostly untractable, even in low dimensional cases like ternary quartics (i.e. $(n, m) = (3, 8)$). Nevertheless, in this paper, the Pfaffian form has proved to be a powerful tool to exhibit continuous families of pairwise non-isomorphic nonsingular algebras. It is proved in [O12c] that any positive ternary quartic form is the Pfaffian form of at least one nonsingular algebra of type $(3, 8)$, showing that a reasonable classification of nonsingular algebras of type $(3, 8)$ is hopeless.

Concerning question (ii) above, the meaning of the word ‘canonical’ is part of the problem. The maps $J_\mu(Z)$ provide a nice tool to consider compatibility conditions between μ and $\langle \cdot, \cdot \rangle$. For instance, when $J_\mu(Z)^2 = -\|Z\|^2 I$ for any $Z \in \mathfrak{n}_2$, the metric Lie algebra $(\mathfrak{n}, \mu, \langle \cdot, \cdot \rangle)$ is called *H-type* (see [K81, BTV95]), and if more in general, $Z \mapsto (-\det J_\mu(Z)^2)^{1/m}$ is a quadratic form, then the algebra is said to be of *\tilde{H} -type* (see [LT99]).

We may also consider the following condition involving only μ and $\langle \cdot, \cdot \rangle$, as another generalization of H -type: for any (or some) orthonormal basis $\{Z_i\}$ of \mathfrak{n}_2 ,

$$(1) \quad \sum J_\mu(Z_i)^2 = aI, \quad \text{tr } J_\mu(Z_i)J_\mu(Z_j) = b\delta_{ij}, \quad \text{for some } a, b < 0.$$

It follows from well-known results in geometric invariant theory (see Section 2.3) that if condition (1) holds, then:

- μ is a minimal vector for the $\text{SL}_m \times \text{SL}_n$ -action on $V_{n,m}$ (i.e. $\|\mu\| \leq \|h \cdot \mu\|$ for any $h \in \text{SL}_m \times \text{SL}_n$).
- The $\text{SL}_m \times \text{SL}_n$ -orbit of μ is closed in $V_{n,m}$.
- The set of minimal vectors in $(\text{SL}_m \times \text{SL}_n) \cdot \mu$ consists of a single $\text{SO}(m) \times \text{SO}(n)$ -orbit. We will strongly use this uniqueness property to distinguish, up to isomorphism, those special nonsingular algebras satisfying (1) by using $\text{SO}(m) \times \text{SO}(n)$ -invariants, which are much more abundant.

From a different geometric point of view, condition (1) implies that the left invariant metric g determined by $(\mathfrak{n}, \mu, \langle \cdot, \cdot \rangle)$ on the corresponding simply connected nilpotent Lie group N maximizes the scalar curvature among all left-invariant metrics on N with $\mathfrak{n}_1 \perp \mathfrak{n}_2$ and the same volume. It is also a minimum of the square norm of the Ricci tensor among all left-invariant metrics on N with the same scalar curvature. On the other hand, (N, g) is known to be a *Ricci soliton*, i.e. the Ricci flow solution starting at g evolves only by time-dependent scaling and pull-back by diffeomorphisms. Such metrics are called *nilsolitons* in the literature and have been extensively studied in the last decade (see e.g. the survey [L09]). Additionally, a metric satisfying (1) is called a *metric with optimal Ricci tensor*, and in particular, has a geodesic-flow invariant Ricci tensor (see [E04, Section 7]). An inner product for which (1) holds is unique up to scaling and automorphisms of μ (i.e. up to homothety), which also supports the presentation of these inner products as canonical or distinguished for a given μ .

We prove in this paper that condition (1) is actually also necessary for any nilsoliton on a nonsingular algebra (see Theorem 2.15 for a more complete statement).

Theorem 1.1. *If μ is a nonsingular algebra such that $\langle \cdot, \cdot \rangle$ is a nilsoliton, then condition (1) holds.*

In [N08], Y. Nikolayevsky gave a complete classification of 2-step nilpotent Lie algebras of type $(2, m)$ admitting a nilsoliton. We give in Section 3 an alternative proof in the nonsingular case.

Theorem 1.2. *A nonsingular algebra $\mu_{\mathcal{S}}$ of type $(2, m)$, where $\mathcal{S} = \{(\alpha_1, k_1), \dots, (\alpha_r, k_r)\}$, admits a nilsoliton if and only if $k_1 = \dots = k_r = 1$.*

We use this result to obtain k -parameter families of nonsingular algebras of type $(2, m)$ which do not admit nilsoliton metrics, for any $k \geq 1$. For the type $(2, 8)$, $k \geq 2$, the curve $\mu_t = \mu_{\mathcal{S}_t}$, where $\mathcal{S}_t := \{(\mathbf{i}, 1), (\sqrt{t}\mathbf{i}, 1)\}$, $t \geq 1$, is pairwise non-isomorphic and consists of nonsingular algebras which do admit a nilsoliton. Their Pfaffian forms equal $f_{\mu_t}(x, y) = (x^2 + y^2)(x^2 + ty^2)$. On the other hand, notice that $\mu_{\mathcal{S}}$ with $\mathcal{S} := \{(\mathbf{i}, 2)\}$ has identical Pfaffian than μ_1 , given by $(x^2 + y^2)^2$, but $\mu_{\mathcal{S}}$ does not admit a nilsoliton. We also use Nikolayevsky's classification to obtain curves of 2-step algebras of type $(2, m)$ (not necessarily nonsingular) which do not admit nilsoliton metrics. Our examples cover any dimension ≥ 14 and dimension 12 (compare with [J11, P11, O12a]).

In Section 4, we give many curves of nonsingular algebras of type $(3, 8)$ admitting a nilsoliton metric. We have also found the first examples of nonsingular algebras of type

(3, 8) which do not admit nilsolitons. They consist of two curves having identical Pfaffian, but with different dimensions for their respective automorphism groups.

Let us now consider question (iii) above. It has recently been proved by A. Kaplan and A. Tiraboschi in [KT12] that for any nonsingular algebra μ , the projection of the automorphism group $\text{Aut}(\mu)$ on $\mathfrak{gl}_n = \mathfrak{gl}(\mathfrak{n}_2)$ satisfies that

$$(2) \quad \dim \text{Aut}(\mu)|_{\mathfrak{n}_2} \leq 1 + \frac{n(n-1)}{2},$$

by showing the existence of an $\text{Aut}(\mu)$ -invariant (up to scaling) inner product on \mathfrak{n}_2 (which generalizes L. Saal's result in the case of H -type algebras, see [S96]). It is well known that equality holds for H -type algebras (see [Ri82]), and it was believed for a long time that this might be a way to characterize H -type algebras among nonsingular ones. In Section 5, we show that this is not true by proving that the nonsingular algebra $\mu_{\mathcal{S}}$ of type $(2, 4k)$, where $\mathcal{S} = \{(i, k)\}$, which is of H -type if and only if $k = 1$, satisfies the equality in condition (2) for any $k \in \mathbb{N}$ (we have recently become aware that this result was independently obtained in [KT12]).

On the other hand, it is natural to consider the stronger condition

$$(3) \quad \dim K(\mu)|_{\mathfrak{n}_2} = \frac{n(n-1)}{2}, \quad \text{or equivalently,} \quad \text{SO}(n) \subset K(\mu)|_{\mathfrak{n}_2},$$

where $K(\mu)$ denotes the subgroup of $\text{Aut}(\mu)$ of orthogonal automorphisms. Clearly, this also holds for H -type algebras, and it is claimed in [KT12, Theorem 2.5] that H -type algebras are the only nonsingular algebras for which (3) holds. However, we give many examples of nonsingular algebras in each type $(3, 4k)$, $k \geq 2$, satisfying (3), none of which is H -type. The simplest one is of type $(3, 8)$ and is obtained by taking $J_{\mu}(\mathfrak{n}_2) \subset \mathfrak{so}(8)$ to be any Lie subalgebra of $\mathfrak{so}(8)$ isomorphic to $\mathfrak{su}(2)$ and acting irreducibly on \mathbb{R}^8 .

Acknowledgements. This research is part of the Ph.D. thesis (Universidad Nacional de Córdoba) by the second author, under the supervision of the first one (see [O12b]). We are very grateful to F. Cukierman, M. Jablonski and W. Ziller for fruitful discussions on the topic of this paper.

2. INVARIANTS OF NONSINGULAR ALGEBRAS

We consider a real vector space \mathfrak{n} and fix a direct sum decomposition

$$\mathfrak{n} = \mathfrak{n}_1 \oplus \mathfrak{n}_2, \quad \dim \mathfrak{n}_1 = m, \quad \dim \mathfrak{n}_2 = n.$$

Every 2-step nilpotent Lie algebra of dimension $m + n$ with derived algebra of dimension $\leq n$ can be represented by a bilinear skew-symmetric map

$$\mu : \mathfrak{n}_1 \times \mathfrak{n}_1 \longrightarrow \mathfrak{n}_2.$$

The set of all such maps is the vector space $V_{n,m} := \Lambda^2 \mathfrak{n}_1^* \otimes \mathfrak{n}_2$ of dimension $n \binom{m}{2}$, and an element $\mu \in V_{n,m}$ is said to be of *type* (n, m) if $\mu(\mathfrak{n}_1, \mathfrak{n}_1) = \mathfrak{n}_2$. We note that these elements form an open and dense subset of $V_{n,m}$, and two of them are isomorphic as Lie algebras if and only if they lie in the same $\text{GL}_m \times \text{GL}_n$ -orbit with respect to the natural action:

$$(4) \quad (\psi, \varphi) \cdot \mu := \varphi \mu(\psi^{-1} \cdot, \psi^{-1} \cdot), \quad (\psi, \varphi) \in \text{GL}_m \times \text{GL}_n, \quad \mu \in V_{n,m}.$$

Here and throughout the rest of the paper, GL_n will denote the group of invertible linear operators of an n -dimensional vector space, and also the real general linear group $\text{GL}_n(\mathbb{R})$ if there is a fixed basis in the context.

If we fix bases $\{X_1, \dots, X_m\}$ and $\{Z_1, \dots, Z_n\}$ of \mathfrak{n}_1 and \mathfrak{n}_2 , respectively, then each $\mu \in V_{n,m}$ is determined by its structural constants $\mu_{ij}^k \in \mathbb{R}$ defined by

$$\mu(X_i, X_j) = \sum_k \mu_{ij}^k Z_k.$$

An alternative way to arrange the structural constants of μ is by fixing the inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{n} that makes the above basis orthonormal in order to define $J_\mu : \mathfrak{n}_2 \rightarrow \mathfrak{so}(\mathfrak{n}_1)$ by

$$\langle J_\mu(Z)X, Y \rangle = \langle \mu(X, Y), Z \rangle, \quad \forall X, Y \in \mathfrak{n}_1, Z \in \mathfrak{n}_2.$$

Indeed, the ij -entry of the matrix of $J_\mu(Z_k)$ is precisely given by $-\mu_{ij}^k$.

Example 2.1. The 11-dimensional 2-step algebra $\mu \in V_{3,8}$ defined by

$$\begin{aligned} \mu(X_1, X_3) &= -Z_1, & \mu(X_1, X_5) &= -Z_2, & \mu(X_1, X_7) &= -2Z_2 + \sqrt{2}Z_3, \\ \mu(X_2, X_4) &= -Z_1, & \mu(X_2, X_6) &= -Z_2, & \mu(X_2, X_8) &= -Z_2 + \frac{1}{\sqrt{2}}Z_3, \\ \mu(X_3, X_6) &= -Z_3, & \mu(X_3, X_8) &= -\frac{1}{\sqrt{2}}Z_2, & \mu(X_4, X_6) &= -Z_3, \\ \mu(X_4, X_8) &= -\frac{1}{\sqrt{2}}Z_2, & \mu(X_5, X_8) &= -Z_1, & \mu(X_6, X_7) &= -Z_1, \end{aligned}$$

can be represented by its structural map J_μ given by

$$J_\mu(xZ_1 + yZ_2 + zZ_3) = \begin{bmatrix} 0 & 0 & x & 0 & y & 0 & 2y - \sqrt{2}z & 0 \\ 0 & 0 & 0 & x & 0 & y & 0 & y + \frac{1}{\sqrt{2}}z \\ -x & 0 & 0 & 0 & z & 0 & \sqrt{2}y & 0 \\ 0 & -x & 0 & 0 & 0 & z & 0 & -\frac{1}{\sqrt{2}}y \\ -y & 0 & -z & 0 & 0 & 0 & 0 & x \\ 0 & -y & 0 & -z & 0 & 0 & x & 0 \\ -2y + \sqrt{2}z & 0 & -\sqrt{2}y & 0 & 0 & -x & 0 & 0 \\ 0 & -y - \frac{1}{\sqrt{2}}z & 0 & \frac{1}{\sqrt{2}}y & -x & 0 & 0 & 0 \end{bmatrix}.$$

We focus in this paper on the following special class of algebras.

Definition 2.2. A 2-step algebra $(\mathfrak{n} = \mathfrak{n}_1 \oplus \mathfrak{n}_2, \mu)$, $\mu \in V_{n,m}$, is called *nonsingular* if any of the following conditions holds:

- The map $\mathfrak{n}_1 \rightarrow \mathfrak{n}_2$, $Y \mapsto \mu(X, Y)$ is onto for any nonzero $X \in \mathfrak{n}_1$.
- For each nonzero $Z \in \mathfrak{n}_2$, the map $(X, Y) \mapsto \langle \mu(X, Y), Z \rangle$ is a non-degenerate 2-form on \mathfrak{n}_1 .
- $J_\mu(Z)$ is invertible for any nonzero $Z \in \mathfrak{n}_2$.

It is easy to see that these conditions are indeed equivalent and do not depend on the chosen inner product. It follows from the strong restrictions for the maximal number of linearly independent vector fields on spheres that nonsingular algebras can only exist in very special dimensions. Indeed, if $m = (2a + 1)2^{4b+c}$, where $0 \leq c \leq 3$, then $n \leq 2^c + 8b - 1$. In particular, for $n \leq 11$ one has

$$\frac{n}{m} \mid \frac{1}{2k} \mid \frac{2-3}{4k} \mid \frac{4-7}{8k} \mid \frac{8}{16k} \mid \frac{9}{32k} \mid \frac{10-11}{64k}, \quad k \in \mathbb{N}.$$

The subset $V_{n,m}^+ \subset V_{n,m}$ of nonsingular algebras is clearly open in $V_{n,m}$, although it is not always dense. For instance, $V_{2,4}^+$ consists of a single $\mathrm{GL}_4 \times \mathrm{GL}_2$ -orbit of dimension 12, the one corresponding to the (real) H-type algebra $\mathfrak{h}_3 \otimes \mathbb{C}$ (where \mathfrak{h}_3 denotes the 3-dimensional real Heisenberg algebra), but the orbit of $\mathfrak{h}_3 \oplus \mathfrak{h}_3$ is also 12-dimensional (and open) and so $V_{2,4}^+$ is not dense in $V_{2,4}$.

There is a great lack of known invariants to distinguish isomorphism classes in $V_{n,m}$, even for nonsingular elements. A complete classification is only known for the types $(1, m)$ (all Heisenberg algebras up to abelian factors), $(2, m)$ (see [G73] or Section 3), $(5, 5)$ and

One can therefore use invariants of forms to distinguish algebras in $V_{n,m}$ up to isomorphism. Consider the ring $\mathbb{R}[P_{n,d}]^{\text{SL}_n}$ of SL_n -invariant polynomials on $P_{n,d}$, that is, the set of all polynomial functions $I: P_{n,d} \rightarrow \mathbb{R}$ such that

$$I(\varphi \cdot f) = I(f) \quad \forall \varphi \in \text{SL}_n, \quad f \in P_{n,d}.$$

It is well known that I is invariant if and only if each of its homogeneous components are so. However, a set of generators and their relations for such a ring is only known for small values of n and d , including $d = 2$ and any n , $n = 2$ and $d \leq 8$, $n = 3$ and $d \leq 3$. We refer to [Do03, M03] for many explicit classification results on forms and invariants of forms.

Lemma 2.5. *For $f, g \in P_{n,d}$, n odd, assume that there exist homogeneous invariants $I, I' \in \mathbb{R}[P_{n,d}]^{\text{SL}_n}$ of the same degree such that $I'(f), I'(g) \neq 0$ and*

$$\frac{I(f)}{I'(f)} \neq \frac{I(g)}{I'(g)}.$$

Then f is not projectively equivalent to g .

The proof easily follows by using that $I(c\varphi \cdot f) = c^k I(f)$ for any $c \in \mathbb{R}$, $\varphi \in \text{SL}_n$, $f \in P_{n,d}$, where k is the degree of the invariant I , and the fact that for n odd one has $\text{GL}_n = \mathbb{R}^* \text{SL}_n$.

Example 2.6. Suppose we have a one-parameter family $\mu_t \in V_{3,8}$ of 2-step algebras with Pfaffian forms

$$f_{\mu_t}(x, y, z) = (x^2 + y^2 + z^2)^2 + tx^2y^2 \in P_{3,4}.$$

It follows that μ_t is nonsingular for any $t \geq 0$ (recall that $\det J_{\mu_t}(Z) = f_{\mu_t}(Z)^2$), but the question is: is this really a ‘curve’ of algebras, in the sense that μ_t is not isomorphic to μ_s for all $t \neq s$? If we write any ternary quartic $f \in P_{3,4}$ as

$$(6) \quad f(x, y, z) = ax^4 + 4bx^3y + 6cx^2y^2 + 4dxy^3 + ey^4 + 4fx^3z + 12gx^2yz + 12hxy^2z \\ + 4iy^3z + 6jx^2z^2 + 12kxyz^2 + 6ly^2z^2 + 4mxz^3 + 4nyz^3 + pz^4,$$

then from [Di87] we get the following SL_3 -invariant homogeneous polynomials on $P_{3,4}$ of degree 3 and 6, respectively:

$$I_3(f) := aep + 3(al^2 + ej^2 + pc^2) + 4(bim + fdn) \\ - 4(ain + efm + pbd) + 6cjl + 12(ck^2 + jh^2 + lg^2) - 12ghk \\ - 12(bkl + fhl + dkj + igj + mhc + ngc) + 12(gdm + hnb + kfi),$$

and

$$(7) \quad I_6(f) := \det H(f), \quad \text{where } H(f) := \begin{vmatrix} a & c & j & g & f & b \\ c & e & l & i & h & d \\ j & l & p & n & m & k \\ g & i & n & l & k & h \\ f & h & m & k & j & g \\ b & d & k & h & g & c \end{vmatrix}.$$

The invariant $I_6(f)$ is called the *calecticant* and $H(f)$ the *Hankel matrix* (or quadratic form) of f . There are other five invariants described in [Di87] of degree 9, 12, 15, 18 and 27, respectively, which are more difficult to handle. Anyway, by a straightforward computation, we get that

$$I_6(f_{\mu_t}) = -\frac{1}{1944}t^3 - \frac{7}{2916}t^2 + \frac{8}{729}t + \frac{20}{729}, \quad I_3(f_{\mu_t}) = \frac{1}{12}t^2 + \frac{4}{9}t + \frac{20}{9},$$

and so

$$\frac{I_6(f_{\mu_t})}{I_3(f_{\mu_t})^2} = -\frac{2(3t^3 + 14t^2 - 64t - 160)}{9(3t^2 + 16t + 80)^3},$$

which is easily seen to be an injective function for $t \in [12, \infty)$. Thus f_{μ_t} , $t \geq 12$, belong to pairwise different projective equivalence classes by Lemma 2.5, which implies that μ_t , $t \geq 12$, represents a family of pairwise non-isomorphic nonsingular algebras of type (3, 8). Other intervals for t giving rise to the same conclusion are $(-\infty, -4]$, $[-4, 0]$ and $[0, 12]$.

2.2. Positive polynomials. The following elementary but still intriguing fact follows from the third condition in Definition 2.2,

$\mu \in V_{n,2d}$ is nonsingular if and only if f_μ (or $-f_\mu$) is a *positive* polynomial, i.e. $f_\mu(x_1, \dots, x_n) > 0$ for any nonzero $(x_1, \dots, x_n) \in \mathbb{R}^n$.

For instance, the 2-step algebras of type (3, 8) given in Examples 2.1-2.3 and 2.4 are both nonsingular. Notice that one of the forms $\pm f \in P_{n,d}$ is positive if and only if f vanishes only at $0 \in \mathbb{R}^n$, and so for the existence of a nonzero positive d -ic form, d must be even. The subset $P_{n,d}^+ \subset P_{n,d}$ of positive forms is open, it is precisely the interior of the closed convex cone of all *nonnegative* (i.e. $f \geq 0$) n -ary d -ic forms (see e.g. [R10, Theorem 3.14]).

In spite the theory of positive polynomials has a long and rich history in the literature, including Hilbert's 17th Problem (see [R10] for further information), a complete classification of them up to projective equivalence seems to be as difficult as the classification of all forms and thus also unknown and mostly untractable, even in low dimensional cases like ternary quartics (i.e. for the type $(n, m) = (3, 8)$).

Example 2.7. There is only one positive quadratic form $f \in P_{n,2}$ up to projective equivalence; namely, $f(x_1, \dots, x_n) = x_1^2 + \dots + x_n^2$. The family $(x_1^2 + \dots + x_n^2)^{d/2}$ provides examples for any even degree $d \geq 2$ of positive polynomials whose SL_n -orbit is closed and are however *singular* (i.e. $\frac{\partial}{\partial x_i} f = 0$, $i = 1, \dots, n$ has a nonzero solution in \mathbb{C}^n) for $d \geq 4$. Their SL_n -orbits are not either of maximal dimension, as their isotropy subgroups all equal $\text{O}(n)$.

Example 2.8. A complex binary d -ic form $f \in P_{2,d}(\mathbb{C})$ can be identified (up to scaling) with its set of zeroes in $\mathbb{P}^1 = \mathbb{P}\mathbb{C}^2$ as follows:

$$f(x, y) = \prod_{i=1}^d (a_i y - b_i x) \quad \leftrightarrow \quad \mathcal{Z}(f) = \{(a_1 : b_1), \dots, (a_d : b_d)\} \subset \mathbb{P}^1.$$

The $\text{SL}_2(\mathbb{C})$ -action on the projective space of $P_{2,d}(\mathbb{C})$ is therefore equivalent to the action on subsets of unordered d points in \mathbb{P}^1 (counting multiplicities). The following facts are well known in invariant theory (see e.g. [Do03, M03]).

- f is *semistable* (i.e. $0 \notin \overline{\text{SL}_2(\mathbb{C}) \cdot f}$) if and only if any element in $\mathcal{Z}(f)$ has multiplicity $\leq d/2$.
- f is *stable* (i.e. $\text{SL}_2(\mathbb{C}) \cdot f$ is closed and the isotropy subgroup at f is finite) if and only if any element in $\mathcal{Z}(f)$ has multiplicity $< d/2$.

It is easy to see that f is real and positive if and only if d is even and

$$\mathcal{Z}(f) = \{(1 : \alpha_1), (1 : \overline{\alpha_1}), \dots, (1 : \alpha_{d/2}), (1 : \overline{\alpha_{d/2}})\}, \quad \text{with } \alpha_i \in \mathbb{C} \setminus \mathbb{R}.$$

Positive binary forms are therefore all stable with the only exception of $(x^2 + y^2)^{d/2}$, which still has anyway closed $\text{SL}_2(\mathbb{C})$ -orbit. As for a real form f , its $\text{SL}_2(\mathbb{C})$ -orbit is closed if and only if its $\text{SL}_2(\mathbb{R})$ -orbit is so (see e.g. [BH62, Proposition 2.3] and [B71, Corollary 5.3]), it follows that $\text{SL}_2(\mathbb{R}) \cdot f$ is closed for any positive binary form f .

Example 2.9. It was proved by Hilbert [H88] that any nonnegative ternary quartic, as well as any binary or quadratic form, is a sum of squares (Hilbert also showed that this does not hold in any other case). It is also known that $f \in P_{3,4}$ (or f a binary or quadratic form) is a sum of 4-th powers of linear forms (and consequently $f \geq 0$) if and only if its Hankel matrix $H(f)$ (see (7)) is positive semidefinite (see the second sentence after [R10, (5.25)] and correct a typo by replacing the first Q by P). This is not longer true in the other cases. However, it is proved in [C07] that for any form $f \in P_{n,d}$, we have that if $f > 0$ ($f \geq 0$) then $H(f) > 0$ ($H(f) \geq 0$).

It is worth noticing at this point that positivity is a notion which only makes sense over \mathbb{R} , and it is therefore difficult to study properties of positive forms from the point of view of invariant theory, a subject mostly attached to algebraic closed fields. We have seen in Examples 2.7 and 2.8 that positive quadratic and binary forms have closed SL_n -orbits. We do not know if this is also true for any form. It certainly fails in the nonnegative case, e.g. $0 \in \overline{\mathrm{SL}_n \cdot x_1^d}$ for any $n \geq 2$ (and d even). We now prove that a positive form is at least always semistable.

Lemma 2.10. *If $f \in P_{n,d}$ is positive, then $0 \notin \overline{\mathrm{SL}_n \cdot f}$.*

Proof. Assume that $0 \in \overline{\mathrm{SL}_n \cdot f}$, that is, there exists a sequence $\varphi_k \in \mathrm{SL}_n$ such that $f \circ \varphi_k \rightarrow 0$, as $k \rightarrow \infty$. If $S \subset \mathbb{R}^n$ is the sphere of radius 1 and $m := \min(f|_S)$, which is a positive number due to the positivity of f , then

$$m \|\varphi_k(Z)\|^d \leq f(\varphi_k(Z)), \quad \forall Z \in \mathfrak{n}_2.$$

This implies that

$$m \|\varphi_k\|^d \leq \max(f \circ \varphi_k|_S) \rightarrow 0,$$

as $k \rightarrow \infty$, and thus $\varphi_k \rightarrow 0$, which is a contradiction since $\det \varphi_k = 1$ for all k . This concludes the proof of the lemma. \square

2.3. Real geometric invariant theory. Let G be a real reductive group acting linearly on a finite dimensional real vector space V . The precise definition of our setting is the one considered in [RS90], or the more general one in [HSS08] (see also [EJ09]), where many results from geometric invariant theory are adapted and proved over \mathbb{R} .

The derivative of the above action defines a representation of the Lie algebra \mathfrak{g} of G in V , which will be denoted by $\pi : \mathfrak{g} \rightarrow \mathrm{End}(V)$. We consider a Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, where \mathfrak{k} is the Lie algebra of a maximal compact subgroup K of G . Endow V with a fixed from now on K -invariant inner product $\langle \cdot, \cdot \rangle$ such that \mathfrak{p} acts on V by symmetric operators, and endow \mathfrak{p} with an $\mathrm{Ad}(K)$ -invariant inner product also denoted by $\langle \cdot, \cdot \rangle$.

The function $m : V \rightarrow \mathfrak{p}$ implicitly defined by

$$\langle m(v), \alpha \rangle = \frac{1}{\|v\|^2} \langle \pi(\alpha)v, v \rangle, \quad \forall \alpha \in \mathfrak{p}, v \in V \setminus \{0\}, \quad m(0) = 0,$$

is called the *moment map* for the representation V of G . Since $m(cv) = m(v)$ for any nonzero $c \in \mathbb{R}$, we may also view the moment map as defined on the projective space $\mathbb{P}V$ of V . It is easy to see that m is K -equivariant: $m(k.v) = \mathrm{Ad}(k)m(v)$ for all $k \in K$.

Let $\mathcal{M} = \mathcal{M}(G, V)$ denote the set of *minimal vectors*, that is,

$$\mathcal{M} = \{v \in V : \|v\| \leq \|g.v\| \quad \forall g \in G\}.$$

In [RS90], the following results were obtained:

- an orbit $G.v$ is closed if and only if $G.v$ meets \mathcal{M} ;
- $G.v \cap \mathcal{M}$ is either empty or consists of a single K -orbit;
- the closure of any G -orbit contains a unique closed G -orbit;

- $\mathcal{M} = \{v \in V : m(v) = 0\}$.

The functional square norm of the moment map

$$(8) \quad F_m : V \longrightarrow \mathbb{R}, \quad F_m(v) := \|m(v)\|^2,$$

is scaling invariant and so it can actually be viewed as a function on any sphere of V or on $\mathbb{P}V$. The remaining critical points of F_m other than minimal vectors (i.e. those for which $F_m(v) > 0$) are all *unstable* (i.e. $0 \in \overline{G \cdot v}$) but still enjoy most of the nice properties of minimal vectors stated above. Let $\mathcal{C}(F_m)$ denote the critical set of $F_m : V \longrightarrow \mathbb{R}$.

Theorem 2.11. [Mr01, HSS08] *The following conditions are equivalent:*

- (i) $v \in \mathcal{C}(F_m)$.
- (ii) The functional $F_m|_{G \cdot v}$ attains its minimum value at v .
- (iii) $\pi(m(v))v = cv$ for some $c \in \mathbb{R}$.

Moreover, the following uniqueness property holds:

- (iv) The intersection of $\mathcal{C}(F_m)$ with any G -orbit is either empty or consists of a single K -orbit (up to scaling).

It follows from part (iv) that two G -orbits containing critical points of F_m can be distinguished by using K -invariants, which are always much more abundant. This method will be illustrated in the examples below.

Example 2.12. Let us consider the natural linear action of $G = \mathrm{GL}_n$ on $V = P_{n,d}$ given by $\varphi \cdot f = f \circ \varphi^{-1}$. We have that $\mathfrak{g} = \mathfrak{gl}_n$, $K = \mathrm{O}(n)$, $\mathfrak{k} = \mathfrak{so}(n)$ and $\mathfrak{p} = \mathrm{sym}(n)$. As an $\mathrm{Ad}(K)$ -invariant inner product on \mathfrak{p} we take $\langle \alpha, \beta \rangle = \mathrm{tr} \alpha \beta$, and it is easy to see that the inner product $\langle \cdot, \cdot \rangle$ on V for which the basis of monomials

$$\{x^D := x_1^{d_1} \dots x_n^{d_n} : d_1 + \dots + d_n = d, D = (d_1, \dots, d_n)\}$$

is orthogonal and

$$\|x^D\|^2 = \frac{d_1! \dots d_n!}{d!}, \quad \forall D = (d_1, \dots, d_n),$$

satisfies the required conditions (indeed, $\langle f \circ \alpha, g \rangle = \langle f, g \circ \alpha^t \rangle$ for all $f, g \in P_{n,d}$, $\alpha \in \mathfrak{gl}_n$, see e.g. [R10, (1.30)]). Let E_{ij} denote as usual the $n \times n$ matrix whose only nonzero coefficient is a 1 in the entries ij . Since

$$\pi(E_{ij})f = \frac{d}{dt} \Big|_0 f \circ e^{-tE_{ij}} = -x_j \frac{\partial f}{\partial x_i},$$

we obtain that the moment map $m : P_{n,d} \longrightarrow \mathrm{sym}(n)$ is given by

$$(9) \quad m(f) = -\frac{1}{\|f\|^2} \left[\left\langle x_j \frac{\partial f}{\partial x_i}, f \right\rangle \right].$$

We are using here that $\langle x_j \frac{\partial f}{\partial x_i}, f \rangle = \langle x_i \frac{\partial f}{\partial x_j}, f \rangle$ for all i, j . It is also easy to see that the action of a diagonal matrix $\alpha \in \mathfrak{gl}_n$ with entries a_1, \dots, a_n is given by

$$(10) \quad \pi(\alpha)x^D = -\left(\sum_{i=1}^n a_i d_i \right) x^D, \quad \forall D = (d_1, \dots, d_n),$$

and since

$$m(x^D) = \begin{bmatrix} -d_1 & & \\ & \ddots & \\ & & -d_n \end{bmatrix},$$

we get that x^D is an eigenvector of $m(x^D)$ with eigenvalue $F_m(x^D) = \sum d_i^2$. Every monomial is therefore a critical point of F_m by Theorem 2.11, although $\mathcal{C}(F_m)$ is known to be much larger (see [N84, Section 10] and [L09, Example 11.5]).

According to the above example, it follows from Theorem 2.11, (iv) that $O(n)$ -invariants of n -ary forms play a key role in distinguishing critical points of $F_m : P_{n,d} \rightarrow \mathbb{R}$. Let us therefore consider the *Laplacian* $\Delta : P_{n,d} \rightarrow P_{n,d-2}$ defined by

$$\Delta(f) := \frac{\partial^2 f}{\partial x_1^2} + \cdots + \frac{\partial^2 f}{\partial x_n^2},$$

which is well known to be $O(n)$ -equivariant:

$$\Delta(\varphi \cdot f) = \varphi \cdot \Delta(f), \quad \forall \varphi \in O(n), \quad f \in P_{n,d}.$$

Example 2.13. Is the curve of ternary quartics $f_t = x^4 + y^4 + z^4 + tx^2y^2$ pairwise non-equivalent? The argument used in Example 2.6 in terms of SL_3 -invariants does not work here because $I_6(f_t) = 0$ for all t . However, if we consider

$$\varphi_t \cdot f_t = x^4 + y^4 + (1 + t^2/12)^{1/2} z^4 + tx^2y^2, \quad \text{where } \varphi_t := \begin{bmatrix} 1 & & \\ & 1 & \\ & & (1+t^2/12)^{-1/8} \end{bmatrix},$$

then it is straightforward to check by using (9) that $m(\varphi_t \cdot f_t) = -\frac{4}{3}I$, which implies that $\varphi_t \cdot f_t \in \mathcal{C}(F_m)$ for all t by Theorem 2.11, (iii). According to Theorem 2.11, (iv) we now only need to differentiate $\varphi_t \cdot f_t$ with $O(3)$ -invariants (up to scaling). A computation of the Laplacian gives

$$\Delta(\varphi_t \cdot f_t) = (12 + 2t)x^2 + (12 + 2t)y^2 + 12(1 + t^2/12)^{1/2}z^2.$$

As the function $\frac{12+2t}{12(1+t^2/12)^{1/2}}$ is the quotient of the two different eigenvalues of these ternary quadratic forms, and since it is a strictly decreasing function for $t \geq 2$, it follows that $\varphi_t \cdot f_t$ (or equivalently, f_t), $t \geq 2$, is a family of pairwise non-equivalent ternary quartics.

We now consider the representation where 2-step algebras live.

Example 2.14. Let $V = V_{n,m}$ be the representation of the real reductive group $G = GL_m \times GL_n$ given by (4), for which we have $\mathfrak{g} = \mathfrak{gl}_m \oplus \mathfrak{gl}_n$, $K = O(m) \times O(n)$, $\mathfrak{k} = \mathfrak{so}(m) \oplus \mathfrak{so}(n)$ and $\mathfrak{p} = \text{sym}(m) \oplus \text{sym}(n)$. Recall that we have fixed an inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{n} . As an $\text{Ad}(K)$ -invariant inner product on \mathfrak{p} we can take $\langle \alpha, \beta \rangle = \text{tr } \alpha\beta$ on each factor and the inner product $\langle \cdot, \cdot \rangle$ on V is defined by

$$(11) \quad \langle \mu, \lambda \rangle := \sum_{ij} \langle \mu(X_i, X_j), \lambda(X_i, X_j) \rangle = \sum_{ijk} \mu_{ij}^k \lambda_{ij}^k.$$

The corresponding representation of \mathfrak{g} on V is

$$(12) \quad \pi(\alpha, \beta)\mu = \beta\mu(\cdot, \cdot) - \mu(\alpha, \cdot) - \mu(\cdot, \alpha),$$

and the moment map $m : V_{n,m} \rightarrow \text{sym}(m) \oplus \text{sym}(n)$ is given by $m(\mu) = (m_1(\mu), m_2(\mu))$ (see e.g. [E08]), where

$$(13) \quad m_1(\mu) = \frac{2}{\|\mu\|^2} \sum_i J_\mu(Z_i)^2, \quad \langle m_2(\mu)Z, W \rangle = -\frac{1}{\|\mu\|^2} \text{tr } J_\mu(Z)J_\mu(W), \quad \forall Z, W \in \mathfrak{n}_2.$$

2.4. Nilsolitons. We consider in this subsection the problem of the existence of a ‘canonical’ inner product for a given nonsingular algebra. Our approach will follow the lines of real geometric invariant theory described in Section 2.3, and it turns out that the uniqueness (up to isometry) of such distinguished inner product will give rise to new invariants.

Recall the maps $J_\mu(Z) \in \mathfrak{so}(\mathfrak{n}_1)$, $Z \in \mathfrak{n}_2$, defined at the beginning of Section 2 by fixing an inner product $\langle \cdot, \cdot \rangle$ on $\mathfrak{n} = \mathfrak{n}_1 \oplus \mathfrak{n}_2$ (with $\mathfrak{n}_1 \perp \mathfrak{n}_2$). These maps provide a nice tool to consider compatibility conditions between an algebra μ and an inner product $\langle \cdot, \cdot \rangle$. For instance, when $J_\mu(Z)^2 = -\|Z\|^2 I$ for any $Z \in \mathfrak{n}_2$, the metric Lie algebra $(\mathfrak{n}, \mu, \langle \cdot, \cdot \rangle)$ is called *H-type* (see [K81, BTV95]), and if more in general, $Z \mapsto (-\det J_\mu(Z)^2)^{1/m}$ is a positive

quadratic form, then the algebra is said to be of \tilde{H} -type (see [LT99]). Equivalently, in terms of its Pfaffian form, $(\mathfrak{n}, \mu, \langle \cdot, \cdot \rangle)$ is \tilde{H} -type if and only if, up to projective equivalence,

$$f_\mu(x_1, \dots, x_n) = (x_1^2 + \dots + x_n^2)^{m/4}.$$

For instance, the algebra in Example 2.4 is of \tilde{H} -type but not of H -type.

We may also consider the following condition involving only μ and $\langle \cdot, \cdot \rangle$, as another generalization of H -type: for any (or some) orthonormal basis $\{Z_i\}$ of \mathfrak{n}_2 ,

$$(14) \quad \sum_i J_\mu(Z_i)^2 = aI, \quad \text{tr } J_\mu(Z_i)J_\mu(Z_j) = b\delta_{ij}, \quad \text{for some } a, b < 0.$$

In terms of the moment map, this condition on $(\mu, \langle \cdot, \cdot \rangle)$ is equivalent to have $m(\mu) = (2aI, -bI)$ (see (13)), from which easily follows that part (iii) in Theorem 2.11 holds and thus μ is a critical point of F_m . From the uniqueness statement in part (iv) of the same theorem, we obtain that among the isomorphism class $\text{GL}_m \times \text{GL}_n \cdot \mu$, only the elements in the subset $\mathbb{R}^* \text{O}(n) \cdot \mu$ satisfy condition (14). By using that such condition holds for the pair $(\varphi \cdot \mu, \langle \cdot, \cdot \rangle)$, $\varphi \in \text{GL}_m \times \text{GL}_n$ if and only if does so for $(\mu, \langle \varphi \cdot, \varphi \cdot \rangle)$, this uniqueness property can be rephrased as follows:

Given an algebra $\mu \in V_{n,m}$, there exists at most one inner product on \mathfrak{n} (with $\mathfrak{n}_1 \perp \mathfrak{n}_2$) for which (14) holds, up to scaling and automorphisms of μ .

This strongly supports the proposal of these inner products as canonical or distinguished for a given μ .

It also follows from invariant theory that if condition (14) holds, then the $\text{SL}_m \times \text{SL}_n$ -orbit of μ is closed in $V_{n,m}$. Indeed, the orthogonal projection of the moment map $m(\mu)$ onto $\mathfrak{sl}_m \oplus \mathfrak{sl}_n$ vanishes, and since this is precisely the moment map for the $\text{SL}_m \times \text{SL}_n$ -action on $V_{n,m}$ we get that μ is a minimal vector and so $\text{SL}_m \times \text{SL}_n \cdot \mu$ is closed (see Section 2.3). Furthermore, the following conditions are equivalent for a 2-step algebra $\mu \in V_{n,m}$:

- there exists an inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{n} such that condition (14) holds for $(\mu, \langle \cdot, \cdot \rangle)$ (in particular, $\mu \in \mathcal{C}(F_m)$);
- for any fixed inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{n} , there exists an isomorphic algebra $\mu_0 \in \text{GL}_m \times \text{GL}_n \cdot \mu$ such that condition (14) holds for $(\mu_0, \langle \cdot, \cdot \rangle)$ (or equivalently, μ_0 is a minimal vector for the $\text{SL}_m \times \text{SL}_n$ -action);
- the orbit $\text{SL}_m \times \text{SL}_n \cdot \mu$ is closed.

In particular, the existence for a given nonsingular μ of an inner product satisfying (14) provides in a way a candidate for a ‘normal form’ for μ ; namely, the minimal vector μ_0 above, which is unique up to scaling and the action of $\text{O}(m) \times \text{O}(n)$.

From a geometric point of view, we know that μ is a critical point of F_m if and only if the left invariant Riemannian metric g determined by $(\mathfrak{n}, \mu, \langle \cdot, \cdot \rangle)$ on the corresponding simply connected nilpotent Lie group N is a *Ricci soliton*, i.e. the Ricci flow solution starting at g evolves only by time-dependent scaling and pull-back by diffeomorphisms (see [L01]). Such metrics are called *nilsolitons* in the literature and have been extensively studied in the last decade (see the survey [L09] for further information). The core of this interplay relies in the fact that

$$(15) \quad m(\mu) = \frac{4}{\|\mu\|^2} \text{Ric}(g),$$

where $m(\mu)$ is the moment map and $\text{Ric}(g)$ is the Ricci operator of (N, g) .

We now prove that condition (14) must actually hold for any nilsoliton on a nonsingular algebra.

Theorem 2.15. *Let μ be a nonsingular algebra of type (n, m) and set $\mathfrak{n} = \mathfrak{n}_1 \oplus \mathfrak{n}_2$. Then the following conditions are equivalent:*

- (i) \mathfrak{n} admits an inner product for which condition (14) holds.
- (ii) The Lie algebra (\mathfrak{n}, μ) admits a nilsoliton inner product (or equivalently, $\mu \in \mathcal{C}(F_m)$ for some $\langle \cdot, \cdot \rangle$ on \mathfrak{n}).
- (iii) The $\mathrm{SL}_m \times \mathrm{SL}_n$ -orbit of μ is closed in $V_{n,m}$.

Remark 2.16. This theorem also follows by using [N06, Lemma 5] and [N11, Theorem 2].

Proof. The equivalence between parts (i) and (iii) has been observed above (cf. [L03, Proposition 9.1]), as it has the fact that part (i) implies (ii). Let us therefore assume part (ii) and prove part (i). The nonsingularity of μ implies that

$$(16) \quad 0 \notin \overline{\mathrm{SL}_m \times \mathrm{SL}_n \cdot \mu},$$

the closure of the orbit relative to the vector space topology. Indeed, if $\mu_k := (\psi_k, \varphi_k) \cdot \mu \rightarrow 0$, as $k \rightarrow \infty$, with $(\psi_k, \varphi_k) \in \mathrm{SL}_m \times \mathrm{SL}_n$, then their Pfaffian forms $f_{\mu_k} \rightarrow 0$, which are given by $f_{\mu_k} = f_{\mu} \circ \varphi_k^t \in \mathrm{SL}_n \cdot f_{\mu}$ for all k , and so this contradicts Lemma 2.10 since f_{μ} is positive.

If we use the nilsoliton inner product on \mathfrak{n} (with $\mathfrak{n}_1 \perp \mathfrak{n}_2$) to define the moment map m , then $\mu \in \mathcal{C}(F_m)$ by [L01, Theorem 4.2]. Let $\beta \in \mathfrak{gl}_{m+n}$ be the unique matrix of the form $\beta = \begin{bmatrix} cI_m & \\ & dI_n \end{bmatrix}$ such that $\mathrm{tr} \beta = -1$ (i.e. $cm + dn = -1$) and $\beta + \|\beta\|^2 I = \begin{bmatrix} eI_m & \\ & 2eI_n \end{bmatrix}$ for some $e \in \mathbb{R}$ (i.e. $d + c^2 m + d^2 n = 2(c + c^2 m + d^2 n)$).

In what follows, we will use the notation in [L10, pp. 1869-1870]. We have that $\mu \in Z_{\beta}$ by the fact that $\beta + \|\beta\|^2 I \in \mathrm{Der}(\mu)$, and since $H_{\beta} = \mathrm{SL}_m \times \mathrm{SL}_n$, it follows from (16) and [L10, Proposition 2.14,(i)] that μ belongs to the stratum \mathcal{S}_{β} . Finally, by using [LW11, Theorem 2.3] and [LW11, Proposition 3.4,(ii)], we get from $\mu \in \mathcal{C}(F_m)$ that $m(\mu)$ is conjugate to β (see (15)). This implies that condition (14) holds for the pair $(\varphi \cdot \mu, \langle \cdot, \cdot \rangle)$ for some $\varphi \in \mathrm{O}(m) \times \mathrm{O}(n)$, and hence it does so for $(\mu, \langle \cdot, \cdot \rangle)$, from which part (i) follows. \square

By fixing an inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{n} as in previous sections, we deduce the following from Theorem 2.15.

Corollary 2.17. *If μ is nonsingular, then the following conditions are equivalent:*

- (i) The nilmanifold (N, g) attached to $(\mathfrak{n}, \mu, \langle \cdot, \cdot \rangle)$ is a nilsoliton.
- (ii) Condition (14) holds for $(\mathfrak{n}, \mu, \langle \cdot, \cdot \rangle)$.
- (iii) The Ricci operator $\mathrm{Ric}(g)$ has exactly two different eigenvalues.

Example 2.18. H -type Lie groups are all nonsingular nilsolitons. On the other hand, it is easy to check that the 2-step algebra of type (3, 8) given in Example 2.4 is a nonsingular nilsoliton which is not H -type.

In order to get explicit continuous families of nonsingular algebras of type (n, m) admitting a nilsoliton metric, we may consider those $\mu \in V_{n,m}$ such that each entries of $J_{\mu}(x_1 Z_1 + \cdots + x_n Z_n)$ is a scalar multiple of some x_i and there is no any repetition of the x_i 's appearing in any given row (or column). This is equivalent to say that $\{X_1, \dots, X_m, Z_1, \dots, Z_n\}$ is a *nice* basis for the algebra μ (see e.g. [LW11, Lemma 3.9] or [N11, Definition 3]). In the presence of a nice basis, Y. Nikolayevsky has proved a very useful criterium for the existence of a nilsoliton; namely, when a certain symmetric $q \times q$ matrix U which only depends on the set of indices (i, j, k) with $\mu_{ij}^k \neq 0$, has a positive solution $v \in \mathbb{R}^q$ to the equation $Uv = [1]$, where $[1] \in \mathbb{R}^q$ denotes the vector with all its entries 1 (see [N11, Theorem 3]). This approach will be used in Section 4.

If in addition to the nice basis condition we ask for each row (or column) of $J_{\mu}(x_1 Z_1 + \cdots + x_n Z_n)$ to have the same number of nonzero entries, then the metric algebra is called *uniform* (see [D79, Wo91]). In this case, it is easy to see that the matrix U from Nikolayevky's criterium has all 3's as diagonal entries (as always) and only 1's out of the diagonal with

an identical number of 1's on each row (or column). It follows that there exists a positive $v \in \mathbb{R}^q$ to $Uv = [1]$ (just take $v_1 = \dots = v_q = a$ for the right a), and thus μ admits a nilsoliton inner product. It is important to note that one obtains this independently on the precise value of the structure constants μ_{ij}^k of the algebra μ , as the uniformity condition is only in terms of the set of structure constants which are nonzero.

Let us summarize the above discussion in a lemma for future reference.

Lemma 2.19. [D79, Wo91] *Any uniform $\mu \in V_{n,m}$ admits a nilsoliton inner product.*

In particular, according to Theorem 2.15, any uniform nonsingular algebra admits an inner product for which condition (14) holds.

3. NONSINGULAR ALGEBRAS OF TYPE $(2, m)$

The set of isomorphism classes in $V_{2,m}$, that is, when $\dim \mathfrak{n}_2 = 2$, can be nicely parameterized by using pencil invariants. We refer to [G73, LT99] and references there in for more detailed treatments and proofs.

To each set

$$\mathcal{S} = \{(\alpha_1, k_1), \dots, (\alpha_r, k_r), \epsilon_1, \dots, \epsilon_s\}, \quad \alpha_i \in \mathbb{C} \cup \{\infty\}, \quad k_i, \epsilon_i \in \mathbb{N},$$

we associate the element $\mu_{\mathcal{S}} \in V_{2,m}$ of type $(2, m)$ such that the matrix of $J_{\mu_{\mathcal{S}}}(xZ_1 + yZ_2)$ with respect to the fixed basis $\{X_1, \dots, X_m\}$ is made of blocks attached to each of the elements of \mathcal{S} , which are defined according to the following rules:

$$(\infty, k) \rightsquigarrow \left[\begin{array}{c|ccc} & 0 & & x \\ & & & x \\ & & \ddots & y \\ & x & y & 0 \\ \hline * & & & \end{array} \right] (2k \times 2k),$$

$$(a, k) \rightsquigarrow \left[\begin{array}{c|ccc} & 0 & & y - ax \\ & & & y - ax \\ & & \ddots & x \\ & y - ax & x & \ddots \\ \hline * & & & 0 \end{array} \right] (2k \times 2k),$$

$$(17) (a+ib, k) \rightsquigarrow \left[\begin{array}{c|cccc} & 0 & & & -bx & y - ax \\ & & & & y - ax & bx \\ & & & -bx & y - ax & 0 \\ & & & y - ax & bx & x \\ & & & & & 0 \\ & & \ddots & & \ddots & \\ & -bx & y - ax & 0 & x & \\ y - ax & bx & x & 0 & & 0 \\ \hline * & & & & & \end{array} \right] (4k \times 4k),$$

$$\epsilon \rightsquigarrow \left[\begin{array}{ccc|ccc} & & & y & & 0 \\ & & & x & y & \\ & & & & \ddots & \ddots \\ & & & & & x & y \\ \hline & & & 0 & & x \\ \hline * & & & & & \end{array} \right] \quad ((2\epsilon + 1) \times (2\epsilon + 1)).$$

The upper right block in the last matrix has $\epsilon + 1$ rows and ϵ columns. Notice that

$$m = \sum_{\alpha_i \in \mathbb{R} \cup \{\infty\}} 2k_i + \sum_{\alpha_i \in \mathbb{C} \setminus \mathbb{R}} 4k_i + \sum_{i=1}^s (2\epsilon_i + 1).$$

Every $\mu \in V_{2,m}$ of type $(2, m)$ (i.e. $\mu(\mathbf{n}_1, \mathbf{n}_1) = \mathbf{n}_2$) is isomorphic to a $\mu_{\mathcal{S}}$ for at least one of such sets \mathcal{S} , and two $\mu_{\mathcal{S}}, \mu_{\mathcal{S}'}$ are isomorphic if and only if $s = s'$, $\{\epsilon_1, \dots, \epsilon_s\} = \{\epsilon'_1, \dots, \epsilon'_s\}$ (counting multiplicities), $r = r'$, and there exists a real Möbius transformation $T: \mathbb{C} \cup \{\infty\} \rightarrow \mathbb{C} \cup \{\infty\}$,

$$Tz := \frac{az + b}{cz + d}, \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{GL}_2(\mathbb{R}),$$

such that $\{(\alpha'_1, k'_1), \dots, (\alpha'_r, k'_r)\} = \{(T\alpha_1, k_1), \dots, (T\alpha_r, k_r)\}$ (counting multiplicities).

It is straightforward to see that the Pfaffian form of $\mu_{\mathcal{S}}$ is given by

$$f_{\mu_{\mathcal{S}}}(x, y) = x^{\sum_{\alpha_i = \infty} k_i} \cdot \prod_{\alpha_i \in \mathbb{R}} (y - \alpha_i x)^{k_i} \cdot \prod_{\alpha_i \in \mathbb{C} \setminus \mathbb{R}} (y - \alpha_i x)^{k_i} (y - \overline{\alpha_i} x)^{k_i},$$

from which we deduce that $\mu_{\mathcal{S}}$ is nonsingular if and only if

$$\mathcal{S} = \{(\alpha_1, k_1), \dots, (\alpha_r, k_r)\}, \quad \text{with } \alpha_1, \dots, \alpha_r \in \mathbb{C} \setminus \mathbb{R}.$$

In that case, if $\alpha_i = a_i + ib_i$ then

$$f_{\mu_{\mathcal{S}}}(x, y) = \prod_i \left((y - a_i x)^2 + b_i^2 x^2 \right)^{k_i}.$$

Example 3.1. If $\mathcal{S} = \{(\mathbf{i}, 1), (\mathbf{i}, 1)\}$ and $\mathcal{S}' = \{(\mathbf{i}, 2)\}$ then $\mu_{\mathcal{S}}, \mu_{\mathcal{S}'} \in V_{2,8}$ are two non-isomorphic nonsingular algebras with identical Pfaffian form given by $(x^2 + y^2)^2$. By adding the family $\mu_{\mathcal{S}_t}$, $\mathcal{S}_t = \{(\mathbf{i}, 1), (t\mathbf{i}, 1)\}$, $t > 1$, of pairwise non-isomorphic algebras with Pfaffian forms $f_{\mu_{\mathcal{S}_t}} = (x^2 + y^2)(t^2 x^2 + y^2)$, one obtains the classification of all nonsingular algebras of type $(2, 8)$ (cf. [LT99, Corollary 4.9, (ii)]). The non-isomorphism within the family follows by using that the hyperbolic distance between \mathbf{i} and $t\mathbf{i}$ strictly increases with t , or alternatively, from the non-equivalence between their Pfaffian forms.

Y. Nikolayevsky has obtained a complete classification of all algebras of type $(2, m)$ admitting a nilsoliton inner product.

Theorem 3.2. [N08] *Let $\mu_{\mathcal{S}}$ be a 2-step algebra of type $(2, m)$, where*

$$\mathcal{S} = \{(\alpha_1, k_1), \dots, (\alpha_r, k_r), (a_1, j_1), \dots, (a_u, j_u), \epsilon_1, \dots, \epsilon_s\}, \\ \alpha_i \in \mathbb{C} \setminus \mathbb{R}, \quad a_i \in \mathbb{R} \cup \{\infty\}, \quad \epsilon_i, k_i, j_i \in \mathbb{N},$$

and assume that either $r > 0$ or $u \geq 3$. Then $\mu_{\mathcal{S}}$ admits a nilsoliton inner product if and only if

- $k_1 = \dots = k_r = 1$ and $j_1 = \dots = j_u = 1$, and
- $\#\{i : a_i = a\} < r + \frac{1}{2}u$, for all $i = 1, \dots, u$.

In particular, λ is also nonsingular. Moreover, it follows from the equivalence between f_λ and $f_{\mu_{\mathcal{S}'}}$ that λ must be isomorphic to an algebra $\mu_{\mathcal{S}'}$ for some set \mathcal{S}' of the form

$$\mathcal{S}' = \{(\alpha_1, k_1), \dots, (\alpha_r, k_r)\},$$

such that for any fixed $j = 1, \dots, r$,

$$\sum_{\alpha_i = \alpha_j} k_i = \#\{\alpha_i : \alpha_i = \alpha_j\}.$$

It is easy to see by using Corollary 3.5 that $\mu_{\mathcal{S}} \in \overline{\mathrm{SL}_{4r} \times \mathrm{SL}_2 \cdot \lambda}$, and since $\lambda \in \overline{\mathrm{SL}_{4r} \times \mathrm{SL}_2 \cdot \mu_{\mathcal{S}'}}$, we obtain that $\lambda \in \mathrm{SL}_{4r} \times \mathrm{SL}_2 \cdot \mu_{\mathcal{S}'}$. This implies that $\mathrm{SL}_{4r} \times \mathrm{SL}_2 \cdot \mu_{\mathcal{S}'}$ is indeed closed, as was to be shown. \square

It follows from Corollaries 3.5 and 3.3 that for any given positive binary form $f \in P_{2,d}$, there exists a unique nonsingular algebra $\mu \in V_{2,2d}$ (up to isomorphism) with Pfaffian form $f_\mu = f$ admitting a nilsoliton inner product. We will see in the next section that this is not longer true for ternary quartics.

We now use Theorem 3.2 to get explicit continuous families of pairwise non-isomorphic 2-step nilpotent Lie algebras of type $(2, m)$ which do not admit nilsoliton metrics. Our examples cover any dimension ≥ 14 and dimension 12. Families of this kind have been exhibited by M. Jablonski in [J11] for any dimension ≥ 24 , and in the 3-step nilpotent case in any dimension ≥ 8 by T. Payne in [P11] (see also [O12a]).

Proposition 3.6. *For any $r \geq 1$, each of the following sets \mathcal{S}_t , $t > 1$, provides a one-parameter family $\mu_{\mathcal{S}_t}$ of pairwise non-isomorphic indecomposable 2-step algebras of type $(2, m)$ which do not admit any nilsoliton metric:*

- $m = 8 + 2r$, $\mathcal{S}_t = \{(\mathbf{i}t, 1), (0, 2), \overbrace{(1, 1), \dots, (1, 1)}^r\}$.
- $m = 11 + 2r$, $\mathcal{S}_t = \{(\mathbf{i}t, 1), (0, 2), \overbrace{(1, 1), \dots, (1, 1)}^r, 1\}$.
- $m = 8 + 4r$, $\mathcal{S}_t = \{(\mathbf{i}t, 2), \overbrace{(\mathbf{i}, 1), \dots, (\mathbf{i}, 1)}^r\}$.
- $m = 8 + 4r + 5$, $\mathcal{S}_t = \{(\mathbf{i}t, 2), \overbrace{(\mathbf{i}, 1), \dots, (\mathbf{i}, 1)}^r, (0, 1), 1\}$.
- $m = 8 + 4r + 2$, $\mathcal{S}_t = \{(\mathbf{i}t, 2), \overbrace{(\mathbf{i}, 1), \dots, (\mathbf{i}, 1)}^r, (0, 1)\}$.
- $m = 8 + 4r + 3$, $\mathcal{S}_t = \{(\mathbf{i}t, 2), \overbrace{(\mathbf{i}, 1), \dots, (\mathbf{i}, 1)}^r, 1\}$.

Proof. In all cases, the non-existence of a nilsoliton follows at once by applying Theorem 3.2, and the pairwise non-isomorphism can be easily deduced from elementary properties of Möbius transformations. \square

By arguing exactly as above, one can also easily obtain k -parameter families of pairwise non-isomorphic indecomposable 2-step algebras of type $(2, m)$ which do not admit any nilsoliton metric for any $k \geq 1$, even in the nonsingular case. For example, take $\mu_{\mathcal{S}_{t_1, \dots, t_k}}$ for

$$\mathcal{S}_{t_1, \dots, t_k} = \{(\mathbf{i}, 2), (\mathbf{i}t_1, 1), \dots, (\mathbf{i}t_k, 1)\}, \quad 1 \leq t_1 \leq \dots \leq t_k,$$

in which case $m = 8 + 4k$.

4. NONSINGULAR ALGEBRAS OF TYPE (3, 8)

In view of the results obtained in Section 3, we have by now a quite clear picture of the nonsingular algebras of type $(2, m)$, including of those admitting a nilsoliton metric. As it is well known that there is only one nonsingular algebra of type $(3, 4)$ (namely, the quaternionic H -type algebra $\mathfrak{n} = \mathbb{H} \oplus \text{Im}\mathbb{H}$), the next type to study in degree of difficulty is $(3, 8)$, which is the aim of this section.

A first problem we have to face in type $(3, 8)$ is that the Pfaffian forms are ternary quartics, a very subtle topic from many points of view. The classification of ternary quartics up to equivalence is an open classical problem in invariant theory, even the ring of invariants $\mathbb{C}[P_{3,4}]^{\text{SL}_n(\mathbb{C})}$ is not yet completely understood. Over the real numbers, it may be inferred that an explicit classification for positive ternary quartics is hopeless. According to a result in [O12c], there is at least one nonsingular $\mu \in V_{3,8}^+$ with Pfaffian form a given $f \in P_{3,4}^+$, so that in some sense, a classification of nonsingular algebras of type $(3, 8)$ can also be considered out of reach. One may get an astounding indication of this by looking at the following existence result due to J. Heber (see [H98, Theorem 6.19]): there exists a 12-parameter continuous family of pairwise non-isomorphic 2-step algebras of type $(3, 8)$ around the H -type one, which are all nonsingular in a neighborhood by continuity. Moreover, they all admit a nilsoliton metric.

4.1. Existence of nilsolitons. The following three propositions provide explicit examples of continuous families of nonsingular algebras admitting a nilsoliton inner product with different types of Pfaffian forms. The existence of a nilsoliton follows in all cases from Lemma 2.19, as they are all uniform algebras. The computation of the Pfaffian forms is straightforward, and in Propositions 4.2 and 4.3, the fact that the algebras in the families are pairwise non-isomorphic follows from Examples 2.13 and 2.6, respectively.

Proposition 4.1. *Let $\mu_{t_1, t_2, t_3} \in V_{3,8}$ be the Lie algebra defined by*

$$J_\mu(xZ_1 + yZ_2 + zZ_3) = \begin{bmatrix} 0 & 0 & x & 0 & y & 0 & z & 0 \\ 0 & 0 & 0 & t_1x & 0 & t_2y & 0 & t_3z \\ -x & 0 & 0 & 0 & z & 0 & -y & 0 \\ 0 & -t_1x & 0 & 0 & 0 & z & 0 & -y \\ -y & 0 & -z & 0 & 0 & 0 & x & 0 \\ 0 & -t_2y & 0 & -z & 0 & 0 & 0 & x \\ -z & 0 & y & 0 & -x & 0 & 0 & 0 \\ 0 & -t_3z & 0 & y & 0 & -x & 0 & 0 \end{bmatrix}.$$

Then μ_{t_1, t_2, t_3} with

$$t_1^2 + t_2^2 + t_3^2 = 1, \quad 0 < t_3 \leq t_2 \leq t_1,$$

is a 2-parameter family of pairwise non-isomorphic nonsingular algebras admitting nilsolitons with Pfaffian forms given by

$$f_{\mu_{t_1, t_2, t_3}}(x, y, z) = (x^2 + y^2 + z^2)(t_1x^2 + t_2y^2 + t_3z^2).$$

Proof. It only remains to prove that they are pairwise non-isomorphic, which we shall do via their Pfaffian forms. We use that an equivalence map between $f_{\mu_{t_1, t_2, t_3}}$ and $f_{\mu_{s_1, s_2, s_3}}$ must either preserve or interchange their irreducible factors and that the isotropy subgroup of $x^2 + y^2 + z^2$ is $O(3)$. It is therefore easy to prove that for $t_i, s_i > 0$, $i = 1, 2, 3$, we have that $f_{\mu_{t_1, t_2, t_3}} \simeq f_{\mu_{s_1, s_2, s_3}}$ if and only if there exist $c > 0$ and a permutation $\sigma \in S_3$ such that either $t_i = cs_{\sigma(i)}$ or $t_i = \frac{c}{s_{\sigma(i)}}$ for all $i = 1, 2, 3$, which is impossible for two different triples satisfying the conditions required by the proposition. \square

The algebra μ_{t_1, t_2, t_3} is isomorphic to the H -type algebra $\mathfrak{n} = \mathbb{H} \oplus \mathbb{H} \oplus \text{Im}\mathbb{H}$ if and only if $t_1 = t_2 = t_3 = 1/\sqrt{3}$.

Proposition 4.2. *Let $\mu_t \in V_{3,8}$ be defined by*

$$J_\mu(xZ_1 + yZ_2 + zZ_3) = \begin{bmatrix} 0 & 0 & ax & 0 & 0 & y & 0 & z \\ 0 & 0 & 0 & bx & y & 0 & -z & 0 \\ -ax & 0 & 0 & 0 & 0 & z & y & 0 \\ 0 & -bx & 0 & 0 & z & 0 & 0 & y \\ 0 & -y & 0 & -z & 0 & 0 & 0 & x \\ -y & 0 & -z & 0 & 0 & 0 & x & 0 \\ 0 & z & -y & 0 & 0 & -x & 0 & 0 \\ -z & 0 & 0 & -y & -x & 0 & 0 & 0 \end{bmatrix},$$

where $a = \frac{-t-\sqrt{t^2-4}}{2}$, $b = \frac{-t+\sqrt{t^2-4}}{2}$. Then μ_t with $t \geq 2$ is a 1-parameter family of pairwise non-isomorphic nonsingular algebras admitting nilsolitons with Pfaffian forms given by

$$f_{\mu_t}(x, y, z) = x^4 + y^4 + z^4 + tx^2y^2.$$

Proposition 4.3. *Let $\mu_t \in V_{3,8}$ be defined by*

$$J_\mu(xZ_1 + yZ_2 + zZ_3) = \begin{bmatrix} 0 & 0 & -t^2x & 0 & y & 0 & 0 & \alpha_t z \\ 0 & 0 & 0 & -t^2x & 0 & y & z & 0 \\ t^2x & 0 & 0 & 0 & 0 & -z & y & 0 \\ 0 & t^2x & 0 & 0 & -\frac{z}{\alpha_t} & 0 & 0 & y \\ -y & 0 & 0 & \frac{z}{\alpha_t} & 0 & 0 & x & 0 \\ 0 & -y & z & 0 & 0 & 0 & 0 & x \\ 0 & -z & -y & 0 & -x & 0 & 0 & 0 \\ -\alpha_t z & 0 & 0 & -y & 0 & -x & 0 & 0 \end{bmatrix},$$

where $\alpha_t := \frac{2t^2}{2t^2+1-\sqrt{4t^2+1}}$. Then μ_t with $t > 1$ is a 1-parameter family of pairwise non-isomorphic nonsingular algebras admitting nilsolitons with Pfaffian forms given by

$$f_{\mu_t}(x, y, z) = (x^2 + y^2 + z^2)^2 + tx^2z^2.$$

We note that the families given in the above three propositions are pairwise different, which easily follows from the non-equivalence between their Pfaffian forms.

4.2. Non-existence of nilsolitons. The following two propositions provide explicit examples of continuous families of nonsingular algebras of type (3, 8) which do not admit a nilsoliton inner product. The algebras in these families are pairwise non-isomorphic since their Pfaffian forms are pairwise non-equivalent, as we have seen in the proof of Proposition 4.1.

Proposition 4.4. *Let $\mu_t \in V_{3,8}$ be defined by*

$$J_{\mu_t}(xZ_1 + yZ_2 + zZ_3) = \begin{bmatrix} 0 & 0 & -x & -y & 0 & 0 & 0 & -z \\ 0 & 0 & yt & -x & 0 & 0 & -z & 0 \\ x & -yt & 0 & -x & 0 & -z & 0 & 0 \\ y & x & x & 0 & -z & 0 & 0 & 0 \\ 0 & 0 & 0 & z & 0 & 0 & -x & -y \\ 0 & 0 & z & 0 & 0 & 0 & y & -x \\ 0 & z & 0 & 0 & x & -y & 0 & -x \\ z & 0 & 0 & 0 & y & x & x & 0 \end{bmatrix}.$$

Then μ_t with $t > 1$ is a 1-parameter family of pairwise non-isomorphic nonsingular algebras with Pfaffian forms given by

$$f_{\mu_t}(x, y, z) = (x^2 + y^2 + z^2)(x^2 + ty^2 + z^2),$$

which do not admit a nilsoliton inner-product.

Proof. If we consider $\lambda_s := \varphi_s \cdot \mu_t$, where

$$\varphi_s := \text{Diag}(e^s, e^s, e^{-s}, e^{-s}, e^s, e^s, e^{-s}, e^{-s}, 1, 1, 1) \in \text{SL}_8 \times \text{SL}_3,$$

then it is easy to prove that $\lambda_s \rightarrow \tilde{\mu}_t$, as $s \rightarrow \infty$, where $\tilde{\mu}_t \in V_{3,8}$ is defined by

$$J_{\tilde{\mu}_t}(xZ_1 + yZ_2 + xZ_3) = \begin{bmatrix} 0 & 0 & -x & -y & 0 & 0 & 0 & -z \\ 0 & 0 & yt & -x & 0 & 0 & -z & 0 \\ x & -yt & 0 & 0 & 0 & -z & 0 & 0 \\ y & x & 0 & 0 & -z & 0 & 0 & 0 \\ 0 & 0 & 0 & z & 0 & 0 & -x & -y \\ 0 & 0 & z & 0 & 0 & 0 & y & -x \\ 0 & z & 0 & 0 & x & -y & 0 & 0 \\ z & 0 & 0 & 0 & y & x & 0 & 0 \end{bmatrix},$$

for all t . This implies that $\tilde{\mu}_t \in \overline{\mathrm{SL}_8 \times \mathrm{SL}_3 \cdot \mu_t}$, and since $\dim \mathrm{Der}(\mu_t) = 26$ and $\dim \mathrm{Der}(\tilde{\mu}_t) = 29$ (which can be easily computed by using Maple) we have that μ_t is not isomorphic to $\tilde{\mu}_t$. It follows that $\mathrm{SL}_8 \times \mathrm{SL}_3 \cdot \mu_t$ is not closed and thus μ_t does not admit a nilsoliton by Theorem 2.15. \square

Proposition 4.5. *Let $\lambda_t \in V_{3,8}$ be defined by*

$$J_{\lambda_t}(xZ_1 + yZ_2 + zZ_3) = \begin{bmatrix} 0 & 0 & -x & -y & 0 & 0 & 0 & -z \\ 0 & 0 & yt & -x & 0 & 0 & -z & 0 \\ x & -yt & 0 & -y & 0 & -z & -y & -y \\ y & x & y & 0 & -z & 0 & -y & -y \\ 0 & 0 & 0 & z & 0 & 0 & -x & -y \\ 0 & 0 & z & 0 & 0 & 0 & y & -x \\ 0 & z & y & y & x & -y & 0 & -y \\ z & 0 & y & y & y & x & y & 0 \end{bmatrix}.$$

Then λ_t with $t > 1$ is a 1-parameter family of pairwise non-isomorphic nonsingular algebras with Pfaffian forms given by

$$f_{\lambda_t}(x, y, z) = (x^2 + y^2 + z^2)(x^2 + ty^2 + z^2),$$

which do not admit a nilsoliton inner-product.

The proof follows in much the same way as the proof of Proposition 4.4, even with the same φ_s and $\tilde{\mu}_t$, and we now use that $\dim \mathrm{Der}(\lambda_t) = 27$ for all t . This also shows that the families in the above two propositions are different.

5. AUTOMORPHISMS OF NONSINGULAR ALGEBRAS

We study in this section automorphisms and derivations of 2-step algebras which are nonsingular. Our main contribution here is to provide examples other than H -type algebras with certain maximality properties concerning the space of derivations.

The Lie algebra of derivations of a 2-step algebra $\mu \in V_{n,m}$ of type (n, m) decomposes as

$$(18) \quad \mathrm{Der}(\mu) = \mathbb{R} \begin{bmatrix} I & 0 \\ 0 & 2I \end{bmatrix} \oplus \begin{bmatrix} 0 & 0 \\ * & 0 \end{bmatrix} \oplus \mathrm{Der}_{gr}(\mu),$$

where $\mathrm{Der}_{gr}(\mu)$ is the subalgebra of graded derivations given by

$$\mathrm{Der}_{gr}(\mu) := \left\{ \begin{bmatrix} B & 0 \\ 0 & A \end{bmatrix} : B \in \mathfrak{sl}_m, A \in \mathfrak{gl}_n, B^t J_\mu(Z) + J_\mu(Z) B = J_\mu(A^t Z), \forall Z \in \mathfrak{n}_2 \right\}.$$

This is the Lie algebra of the Lie subgroup of $\mathrm{Aut}(\mu)$ given by

$$\mathrm{Aut}_{gr}(\mu) := \left\{ \begin{bmatrix} \psi & 0 \\ 0 & \varphi \end{bmatrix} : \psi \in \mathrm{SL}_m, \varphi \in \mathrm{GL}_n, \psi^t J_\mu(Z) \psi = J_\mu(\varphi^t Z), \forall Z \in \mathfrak{n}_2 \right\}.$$

It follows that $f_\mu(\varphi^t Z) = f_\mu(Z)$ for all Z , that is, φ^t is in the automorphism group $\mathrm{Aut}(f_\mu)$ of the Pfaffian form f_μ . When μ is nonsingular, one has that $f_\mu > 0$, and consequently $\mathrm{Aut}(f_\mu)$ is compact. The closure of the projection of $\mathrm{Aut}_{gr}(\mu)$ in $\mathrm{SL}_n = \mathrm{SL}(\mathfrak{n}_2)$ is therefore compact, from which follows the following.

Theorem 5.1. [KT12, Theorem 2.1] *If μ is nonsingular, then there exists an inner product on the center \mathfrak{n}_2 which is invariant under $\text{Aut}_{gr}(\mu)$.*

Let us assume from now on that our fixed inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{n}_2 is $\text{Aut}_{gr}(\mu)$ -invariant for any nonsingular μ we consider. By defining the ideal of $\text{Der}_{gr}(\mu)$,

$$\text{Der}_0(\mu) := \left\{ \begin{bmatrix} B & 0 \\ 0 & A \end{bmatrix} \in \text{Der}_{gr}(\mu) : A = 0 \right\},$$

and the corresponding normal subgroup of $\text{Aut}(\mu)$,

$$\text{Aut}_0(\mu) := \left\{ \begin{bmatrix} \psi & 0 \\ 0 & \varphi \end{bmatrix} \in \text{Aut}_{gr}(\mu) : \varphi = I \right\},$$

one deduces the following.

Corollary 5.2. [KT12, Corollary 2.4] *For any nonsingular μ of type (n, m) ,*

$$\dim \text{Der}_{gr}(\mu) / \text{Der}_0(\mu) = \dim \text{Aut}_{gr}(\mu) / \text{Aut}_0(\mu) \leq \frac{n(n-1)}{2}.$$

Example 5.3. It is well known that $\dim \text{Der}_{gr}(\mu) / \text{Der}_0(\mu) = \frac{n(n-1)}{2}$ for any H -type algebra μ (see [Ri82, S96]). Indeed,

$$\begin{bmatrix} J_\mu(Z) & 0 \\ 0 & R_Z \end{bmatrix} \in \text{Aut}(\mu), \quad \forall Z \in \mathfrak{n}_2, \quad \|Z\| = 1,$$

where $R_Z(W) = 2\langle Z, W \rangle Z - W$ is the reflection with respect to the hyperplane $\{Z\}^\perp$ in the center \mathfrak{n}_2 .

A natural question arises: which nonsingular algebras of type (n, m) satisfies any of the following equivalent conditions? Do these conditions hold only for H -type algebras?

- $\dim \text{Der}_{gr}(\mu) / \text{Der}_0(\mu) = \frac{n(n-1)}{2}$.
- $\text{Der}_{gr}(\mu) / \text{Der}_0(\mu) \simeq \mathfrak{so}(n)$.
- For any $A \in \mathfrak{so}(n)$ there exists $B \in \mathfrak{sl}_m$ such that $\begin{bmatrix} B & 0 \\ 0 & A \end{bmatrix} \in \text{Der}(\mu)$.
- For any $\varphi \in \text{SO}(n)$ there exists $\psi \in \text{SL}_m$ such that $\begin{bmatrix} \psi & 0 \\ 0 & \varphi \end{bmatrix} \in \text{Aut}(\mu)$.

It is worth to note that these conditions imply that the Pfaffian form is necessarily equal, up to scaling, to $f_\mu = (x_1^2 + \dots + x_n^2)^{m/4}$, as this is the only n -ary $m/2$ -ic form which is $\text{SO}(n)$ -invariant. In particular, μ must be of \tilde{H} -type.

We now show that already for type $(2, m)$, there are examples of nonsingular algebras satisfying the above maximality conditions which are not H -type.

Proposition 5.4. *If $\mathcal{S} = \{(i, k_1), \dots, (i, k_r)\}$, then the corresponding nonsingular 2-step algebra $\mu_{\mathcal{S}}$ of type $(2, m)$ (see Section 3) satisfies that $\text{Der}_{gr}(\mu_{\mathcal{S}}) / \text{Der}_0(\mu_{\mathcal{S}}) \simeq \mathfrak{so}(2)$.*

Remark 5.5. This result has been independently proved in [KT12, Theorem 3.5], where it is in addition obtained that $\text{Der}_{gr}(\mu_{\mathcal{S}}) / \text{Der}_0(\mu_{\mathcal{S}}) = 0$ for any other nonsingular algebra of type $(2, m)$.

Proof. Suppose first that $\mathcal{S} = \{(i, k)\}$. It is an involved but straightforward computation to see that

$$\begin{bmatrix} M_1 & & & \\ & M_2 & & \\ & & 0 & 1 \\ & & -1 & 0 \end{bmatrix} \in \text{Der}(\mu_{\mathcal{S}}),$$

and

$$K_0(\mu) := \left\{ \begin{bmatrix} \psi & 0 \\ 0 & \varphi \end{bmatrix} \in K(\mu) : \varphi = I \right\}.$$

5.1. 2-step algebras of *Rep*-type. We now consider a class of 2-step algebras attached to finite-dimensional real representations of compact Lie groups. Our aim is to show that condition (19) can also hold for nonsingular algebras which are not of *H*-type.

Definition 5.6. A 2-step algebra $\mu \in V_{n,m}$ is said to be of *Rep*-type if the following holds:

- \mathfrak{n}_2 is a compact Lie algebra (say with Lie bracket $[\cdot, \cdot]$).
- $J_\mu : \mathfrak{n}_2 \rightarrow \mathfrak{so}(\mathfrak{n}_1)$ is a representation of \mathfrak{n}_2 , i.e.

$$J_\mu([Z, W]) = [J_\mu(Z), J_\mu(W)], \quad \forall Z, W \in \mathfrak{n}_2.$$

We refer to [EH96, L99, E05] for more detailed expositions on *Rep*-type algebras. Some of their nice and interesting properties are now listed, all of which can be easily checked.

- μ is of type (n, m) if and only if the representation J_μ is faithful.
- We can assume from now on that the fixed inner product we have on \mathfrak{n}_2 is $\text{ad } \mathfrak{n}_2$ -invariant, i.e. any adjoint map $\text{ad } Z$ of the Lie algebra \mathfrak{n}_2 is skew-symmetric.
- Each $Z \in \mathfrak{n}_2$ defines a skew-symmetric derivation of μ by

$$\begin{bmatrix} J_\mu(Z) & 0 \\ 0 & \text{ad } Z \end{bmatrix} \in \text{Der}(\mu).$$

(see [L99, Theorem 3.12]).

- μ is nonsingular if and only if $\mathfrak{n}_2 = \mathfrak{su}(2)$ ($= \mathfrak{so}(3)$) and any irreducible subrepresentation of (\mathfrak{n}_1, J_μ) is even dimensional (or equivalently, a copy of the $2(k+1)$ -dimensional $\mathfrak{su}(2)$ -representation $P_{2,k}(\mathbb{C})$, k odd, of binary k -ic complex forms viewed as real, see e.g. [BtD85]). This can be deduced as follows: in order to have that $J_\mu(Z)$ is invertible for any nonzero Z it is necessary and sufficient that $\text{rank}(\mathfrak{n}_2) = 1$ and the weights of the representation (\mathfrak{n}_1, J_μ) being all nonzero.

We therefore obtain the following result by combining the above observations.

Proposition 5.7. *If μ is a nonsingular μ of *Rep*-type, then*

$$\mathfrak{k}(\mu)/\mathfrak{k}_0(\mu) \simeq \mathfrak{so}(3).$$

Therefore, any nonsingular μ of *Rep*-type, which is not of *H*-type, provides a counterexample to [KT12, Theorem 2.5].

It is easy to see that a *Rep*-type μ is of *H*-type if and only if $\mathfrak{n}_2 = \mathfrak{su}(2)$ and (\mathfrak{n}_1, J_μ) is a sum of copies of the standard representation \mathbb{C}^2 of $\mathfrak{su}(2)$. The simplest nonsingular μ of *Rep*-type not of *H*-type is $\mathfrak{n}_2 = \mathfrak{su}(2)$ and the representation $(\mathfrak{n}_1, J_\mu) = P_{2,3}(\mathbb{C}) = \mathbb{C}^4$ viewed as real. This is precisely the example of type $(3, 8)$ considered in Example 2.4, and an elementary way to see that it is not of *H*-type is by noticing that $J_\mu(\mathfrak{n}_2)$ acts irreducibly on \mathfrak{n}_1 .

Any *Rep*-type 2-step algebra admits a nilsoliton metric (see [EH96, L09]). Under the assumptions that \mathfrak{n}_2 is simple and J_μ is irreducible, we have that $(\mathfrak{n}, \mu, \langle \cdot, \cdot \rangle)$ is a nilsoliton itself. Indeed, since $(Z, W) := \text{tr } J_\mu(Z)J_\mu(W)$ is $\text{ad } \mathfrak{n}_2$ -invariant and the Casimir operator $\sum J_\mu(Z_i)^2$ intertwines the representation, we obtain that condition (14) holds by Schur's Lemma.

REFERENCES

- [BLS05] G. BELITSKII, R. LIPYANSKI, V. SERGEICHUK, Problems of classifying associative or Lie algebras and triples of symmetric or skew-symmetric matrices are wild, *Lin. Alg. Appl.* **407** (2005), 249-262.

- [BTV95] J. BERNDT, F. TRICERRI, L. VANHECKE, Generalized Heisenberg groups and Damek-Ricci harmonic spaces, *Lect. Notes in Math.* **1598** (1995) Springer-Verlag Berlin Heidelberg.
- [B71] D. BIRKES, Orbits of Linear Algebraic Groups, *Ann. of Math.* (2) **93** (1971), 459-475.
- [BH62] A. BOREL, HARISH-CHANDRA, Arithmetic subgroups of algebraic groups, *Ann. of Math.* (2) **75** (1962), 485-535.
- [BtD85] T. BROCKER, T. TOM DIECK, Representations of compact Lie groups, *Springer-Verlag*, New York, 1985.
- [C07] F. CUKIERMAN, Positive polynomials and hyperdeterminants, *Collectanea Math.* **58** (2007), 279-289.
- [D79] E. DELOFF, Naturally reductive metrics and metrics with volume preserving geodesic symmetries on NC algebras, Dissertation, Rutgers 1979.
- [Di87] J. DIXMIER, On the projective invariants of quartic plane curves, *Adv. in Math.* **64**, 279-304 (1987).
- [Do03] I. DOLGACHEV, Lectures on invariant theory, *London Math. Soc. Lect. Notes* **296** (2003), Cambridge Univ. Press.
- [E94] P. EBERLEIN, Geometry of 2-step nilpotent Lie groups with a left invariant metric II, *Trans. Amer. Math. Soc.* **343** (1994), 805-828.
- [E04] ———, Geometry of 2-step nilpotent Lie groups, *Modern Dynamical Systems*, Cambridge Univ. Press, 2004, 67-101.
- [E05] ———, Two step nilpotent Lie groups arising from semisimple modules, preprint 2005 (arXiv).
- [E08] ———, Riemannian 2-step nilmanifolds with prescribed Ricci tensor, *Contemp. Math.* **469** (2008), 167-195.
- [EH96] P. EBERLIEN, J. HEBER, Quarter pinched homogeneous spaces of negative curvature, *Internat. J. Math.* **7**(1996) 441-500.
- [EJ09] P. EBERLEIN, M. JABLONSKI, Closed orbits of semisimple group actions and the real M function, *Contemp. Math.* **491** (2009), 283-321.
- [FZ11] L. FLORIT, W. ZILLER, Topological obstructions to fatness, *Geom. Top.* **15** (2011), 891-925.
- [GT99] L. GALITSKI, D. TIMASHEV, On classification of metabelian Lie algebras, *J. Lie Theory* **9** (1999), 125-156.
- [G73] M. GAUGER, On the classification of metabelian Lie algebras, *Trans. Amer. Math. Soc.* **179** (1973), 293-329.
- [H98] J. HEBER, Noncompact homogeneous Einstein spaces, *Invent. math.* **133** (1998), 279-352.
- [HSS08] P. HEINZNER, G. W. SCHWARZ, H. STÖTZEL, Stratifications with respect to actions of real reductive groups, *Compositio Math.* **144** (2008), 163-185.
- [H88] D. HILBERT, Über die darstellung definiter formen als summe von formenquadraten, *Math. Ann.* **32** (1888), 342-350.
- [J11] M. JABLONSKI, Moduli of Einstein and non-Einstein nilradicals, *Geom. Ded.* **152** (2011), 63-84.
- [K81] A. KAPLAN, Riemannian nilmanifolds attached to Clifford modules, *Geom. Dedicata* **11**(1981), 127-136.
- [KT12] A. KAPLAN, A. TIRABOSCHI, Automorphisms of non-singular nilpotent Lie algebras, preprint 2012 (arXiv).
- [L99] J. LAURET, Homogeneous nilmanifolds attached to representations of compact Lie groups, *Manusc. Math.* **99** (1999), 287-309.
- [L01] ———, Ricci soliton homogeneous nilmanifolds, *Math. Annalen* **319** (2001), 715-733.
- [L03] ———, Degenerations of Lie algebras and geometry of Lie groups. *Diff. Geom. Appl.* **18**(2) (2003), 177194.
- [L08] ———, Rational forms of nilpotent Lie algebras and Anosov diffeomorphisms, *Monatsh. Math.* **155**, 15-30 (2008).
- [L09] ———, Einstein solvmanifolds and nilsolitons, *Contemp. Math.* **491** (2009), 1-35.
- [L10] ———, Einstein solvmanifolds are standard, *Ann. of Math.* **172** (2010), 1859-1877.
- [LW11] J. LAURET, C.E. WILL, Einstein solvmanifolds: existence and non-existence questions, *Math. Annalen* **350** (2011), 199-225.
- [LT99] F. LEVSTEIN, A. TIRABOSCHI, Classes of 2-step nilpotent Lie algebras, *Comm. Alg.* **27** (1999), 2425-2440.
- [Mr01] A. MARIAN, On the real moment map, *Math. Res. Lett.* **8** (2001), 779-788.
- [M03] S. MUKAI, An introduction to invariants and moduli, *Cambridge Studies in adv. math* **81** (2003), Cambridge Univ. Press.

- [N84] L. NESS, A stratification of the null cone via the momentum map, *Amer. J. Math.* **106** (1984), 1281-1329 (with an appendix by D. Mumford).
- [N06] Y. NIKOLAYEVSKY, Nilradicals of Einstein solvmanifolds, *arXiv: math.DG/ 0612117v1*.
- [N11] ———, Einstein solvmanifolds and the pre-Einstein derivation, *Trans. Amer. Math. Soc.* **363** (2011), 3935-3958.
- [N08] ———, Einstein solvmanifolds attached to two-step nilradicals, *Math. Z.*, in press.
- [O12a] D. OSCARI, On the existence of nilsolitons on 2-step nilpotent Lie groups, *Adv. Geom.*, in press.
- [O12b] ———, Álgebras de Lie 2-pasos nilpotentes nosingulares, Ph.D. thesis (in spanish), FaMAF, Universidad Nacional de Córdoba, Argentina (2012).
- [O12c] ———, On the surjectivity of the Pfaffian form for 2-step nilpotent Lie algebras, in preparation.
- [P11] T. PAYNE, The topology of the set of nonsoliton Lie algebras in the moduli space of nilpotent Lie algebras, preprint 2011 (arXiv).
- [R10] B. REZNICK, Sum of even powers of real linear forms, *Memoirs Amer. Math. Soc.* **463** (1992).
- [R10] B. REZNICK, On Hilbert's construction of positive polynomials, preprint 2007 (arXiv).
- [RS90] R.W. RICHARDSON, P.J. SŁODOWY, Minimum vectors for real reductive algebraic groups, *J. London Math. Soc. (2)* **42** (1990), 409-429.
- [Ri82] C. RIEHM, The automorphism group of a composition of quadratic forms, *Trans. Amer. Math. Soc.* **269** (1982), 403-414.
- [S96] L. SAAL, The automorphism group of a Lie algebra of Heisenberg type, *Rend. Sem. Mat. Univ. Pol. Torino* **54** (1996), 101-113.
- [Sc67] J. SCHEUNEMAN, Two-step nilpotent Lie algebras, *J. Algebra* **7** (1967), 152-159.
- [W82] E. WILSON, Isometry groups on homogeneous nilmanifolds, *Geom. Ded.* **12** (1982), 337-346.
- [Wo91] T. H. WOLTER, Einstein metrics on solvable groups, *Math. Z.* **206** (1991), 457-471.

FAMAF AND CIEM, UNIVERSIDAD NACIONAL DE CÓRDOBA, 5000 CÓRDOBA, ARGENTINA
E-mail address: lauret@famaf.unc.edu.ar, oscari@famaf.unc.edu.ar